



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 5, 2026 – 12:51 PM UTC

PDB ID : 9J11 / pdb_00009j11
Title : Structure of mEos3.2 in the green fluorescent state
Authors : Zheng, S.P.; Shi, X.R.
Deposited on : 2024-08-03
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	 81% 6% 13%
1	B	248	 79% 8% 13%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3946 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green to red photoconvertible GFP-like protein EosFP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	216	Total	C	N	O	S	0	3	0
			1755	1121	298	326	10			
1	B	216	Total	C	N	O	S	0	2	0
			1752	1119	298	325	10			

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	initiating methionine	UNP Q5S6Z9
A	-22	GLY	-	expression tag	UNP Q5S6Z9
A	-21	SER	-	expression tag	UNP Q5S6Z9
A	-20	SER	-	expression tag	UNP Q5S6Z9
A	-19	HIS	-	expression tag	UNP Q5S6Z9
A	-18	HIS	-	expression tag	UNP Q5S6Z9
A	-17	HIS	-	expression tag	UNP Q5S6Z9
A	-16	HIS	-	expression tag	UNP Q5S6Z9
A	-15	HIS	-	expression tag	UNP Q5S6Z9
A	-14	HIS	-	expression tag	UNP Q5S6Z9
A	-13	SER	-	expression tag	UNP Q5S6Z9
A	-12	GLN	-	expression tag	UNP Q5S6Z9
A	-11	ASP	-	expression tag	UNP Q5S6Z9
A	-10	PRO	-	expression tag	UNP Q5S6Z9
A	-9	LEU	-	expression tag	UNP Q5S6Z9
A	-8	GLU	-	expression tag	UNP Q5S6Z9
A	-7	VAL	-	expression tag	UNP Q5S6Z9
A	-6	LEU	-	expression tag	UNP Q5S6Z9
A	-5	PHE	-	expression tag	UNP Q5S6Z9
A	-4	GLN	-	expression tag	UNP Q5S6Z9
A	-3	GLY	-	expression tag	UNP Q5S6Z9
A	-2	PRO	-	expression tag	UNP Q5S6Z9
A	-1	GLU	-	expression tag	UNP Q5S6Z9
A	0	PHE	-	expression tag	UNP Q5S6Z9
A	11	LYS	ASN	engineered mutation	UNP Q5S6Z9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	64	5SQ	HIS	chromophore	UNP Q5S6Z9
A	64	5SQ	TYR	chromophore	UNP Q5S6Z9
A	64	5SQ	GLY	chromophore	UNP Q5S6Z9
A	70	LYS	GLU	engineered mutation	UNP Q5S6Z9
A	74	ASN	HIS	engineered mutation	UNP Q5S6Z9
A	102	ASN	ILE	engineered mutation	UNP Q5S6Z9
A	121	TYR	HIS	engineered mutation	UNP Q5S6Z9
A	123	THR	VAL	engineered mutation	UNP Q5S6Z9
A	158	GLU	THR	engineered mutation	UNP Q5S6Z9
A	189	ALA	TYR	engineered mutation	UNP Q5S6Z9
B	-23	MET	-	initiating methionine	UNP Q5S6Z9
B	-22	GLY	-	expression tag	UNP Q5S6Z9
B	-21	SER	-	expression tag	UNP Q5S6Z9
B	-20	SER	-	expression tag	UNP Q5S6Z9
B	-19	HIS	-	expression tag	UNP Q5S6Z9
B	-18	HIS	-	expression tag	UNP Q5S6Z9
B	-17	HIS	-	expression tag	UNP Q5S6Z9
B	-16	HIS	-	expression tag	UNP Q5S6Z9
B	-15	HIS	-	expression tag	UNP Q5S6Z9
B	-14	HIS	-	expression tag	UNP Q5S6Z9
B	-13	SER	-	expression tag	UNP Q5S6Z9
B	-12	GLN	-	expression tag	UNP Q5S6Z9
B	-11	ASP	-	expression tag	UNP Q5S6Z9
B	-10	PRO	-	expression tag	UNP Q5S6Z9
B	-9	LEU	-	expression tag	UNP Q5S6Z9
B	-8	GLU	-	expression tag	UNP Q5S6Z9
B	-7	VAL	-	expression tag	UNP Q5S6Z9
B	-6	LEU	-	expression tag	UNP Q5S6Z9
B	-5	PHE	-	expression tag	UNP Q5S6Z9
B	-4	GLN	-	expression tag	UNP Q5S6Z9
B	-3	GLY	-	expression tag	UNP Q5S6Z9
B	-2	PRO	-	expression tag	UNP Q5S6Z9
B	-1	GLU	-	expression tag	UNP Q5S6Z9
B	0	PHE	-	expression tag	UNP Q5S6Z9
B	11	LYS	ASN	engineered mutation	UNP Q5S6Z9
B	64	5SQ	HIS	chromophore	UNP Q5S6Z9
B	64	5SQ	TYR	chromophore	UNP Q5S6Z9
B	64	5SQ	GLY	chromophore	UNP Q5S6Z9
B	70	LYS	GLU	engineered mutation	UNP Q5S6Z9
B	74	ASN	HIS	engineered mutation	UNP Q5S6Z9
B	102	ASN	ILE	engineered mutation	UNP Q5S6Z9
B	121	TYR	HIS	engineered mutation	UNP Q5S6Z9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	123	THR	VAL	engineered mutation	UNP Q5S6Z9
B	158	GLU	THR	engineered mutation	UNP Q5S6Z9
B	189	ALA	TYR	engineered mutation	UNP Q5S6Z9

- Molecule 2 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (CCD ID: DTT) (formula: $C_4H_{10}O_2S_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	O	S	0	0
			8	4	2	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	223	Total	O	0	0
			223	223		
3	B	208	Total	O	0	0
			208	208		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Green to red photoconvertible GFP-like protein EosFP

Chain A: 



- Molecule 1: Green to red photoconvertible GFP-like protein EosFP

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.36Å 82.39Å 61.77Å 90.00° 101.93° 90.00°	Depositor
Resolution (Å)	28.64 – 1.85 28.64 – 1.85	Depositor EDS
% Data completeness (in resolution range)	95.3 (28.64-1.85) 95.3 (28.64-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 1.85Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.153 , 0.184 0.153 , 0.184	Depositor DCC
R_{free} test set	1903 reflections (4.42%)	wwPDB-VP
Wilson B-factor (Å ²)	20.0	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 50.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3946	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5SQ, DTT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/1783	0.57	0/2399
1	B	0.31	0/1777	0.54	0/2391
All	All	0.32	0/3560	0.55	0/4790

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1755	0	1680	10	0
1	B	1752	0	1675	16	0
2	B	8	0	8	0	0
3	A	223	0	0	2	2
3	B	208	0	0	4	2
All	All	3946	0	3363	26	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 26 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:166:ASN:O	1:B:166:ASN:ND2	2.14	0.79
1:A:145:LYS:NZ	3:A:301:HOH:O	2.18	0.76
1:A:104:ARG:NH2	1:A:106:ASP:OD2	2.19	0.75
1:B:124:ASN:ND2	3:B:401:HOH:O	2.12	0.75
1:B:149:ARG:NH2	3:B:403:HOH:O	2.20	0.72

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:465:HOH:O	3:B:597:HOH:O[2_545]	1.98	0.22
3:A:460:HOH:O	3:B:556:HOH:O[2_545]	2.03	0.17

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/248 (86%)	213 (100%)	1 (0%)	0	100	100
1	B	213/248 (86%)	210 (99%)	3 (1%)	0	100	100
All	All	427/496 (86%)	423 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/212 (88%)	186 (100%)	0	100	100
1	B	185/212 (87%)	185 (100%)	0	100	100
All	All	371/424 (88%)	371 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	206	ASN
1	B	81	GLN
1	B	124	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	5SQ	A	64	1	26,27,28	2.52	8 (30%)	34,37,39	5.49	16 (47%)
1	5SQ	B	64	1	26,27,28	2.49	9 (34%)	34,37,39	5.63	16 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	5SQ	A	64	1	-	4/12/31/32	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	5SQ	B	64	1	-	4/12/31/32	0/3/3/3

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	64	5SQ	C1-N2	7.32	1.43	1.32
1	B	64	5SQ	C1-N2	7.03	1.42	1.32
1	B	64	5SQ	CA2-C2	6.23	1.55	1.48
1	A	64	5SQ	CA2-C2	5.99	1.55	1.48
1	B	64	5SQ	CA2-N2	5.12	1.49	1.38

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	5SQ	CB2-CA2-C2	18.72	145.04	122.36
1	B	64	5SQ	CB2-CA2-C2	18.47	144.75	122.36
1	A	64	5SQ	CB2-CA2-N2	-15.60	107.58	128.76
1	B	64	5SQ	O2-C2-CA2	-15.55	121.09	131.02
1	B	64	5SQ	CB2-CA2-N2	-15.25	108.06	128.76

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	64	5SQ	C3-CA3-N3-C2
1	B	64	5SQ	C3-CA3-N3-C2
1	A	64	5SQ	CA1-CB1-CG1-C2H
1	B	64	5SQ	CA1-CB1-CG1-C2H
1	A	64	5SQ	CA1-CB1-CG1-N1H

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DTT	B	301	1	7,7,7	0.93	0	4,8,8	0.79	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DTT	B	301	1	-	0/8/8/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/248 (86%)	-0.45	1 (0%) 87 90	13, 23, 40, 57	3 (1%)
1	B	215/248 (86%)	-0.42	3 (1%) 73 77	13, 23, 40, 59	2 (0%)
All	All	430/496 (86%)	-0.43	4 (0%) 81 84	13, 23, 40, 59	5 (1%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	219	GLY	3.1
1	B	219	GLY	2.6
1	B	121	TYR	2.5
1	B	166	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	5SQ	A	64	25/26	0.97	0.05	12,16,18,20	0
1	5SQ	B	64	25/26	0.97	0.05	13,16,20,22	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DTT	B	301	8/8	0.95	0.13	23,33,45,50	0

6.5 Other polymers

There are no such residues in this entry.