



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 06:10 AM UTC

PDB ID : 1J20 / pdb_00001j20
Title : Crystal Structure of Thermus thermophilus HB8 Argininosuccinate Synthetase in complex with product
Authors : Goto, M.; Hirotsu, K.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2002-12-24
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	400	 2% 73% 21% . .
1	B	400	 2% 73% 21% . .
1	C	400	 6% 70% 23% . .
1	D	400	 3% 70% 22% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	AS1	A	520	X	-	X	-
4	AS1	B	2520	X	X	-	-
4	AS1	C	3520	X	-	X	-
4	AS1	D	4520	X	X	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 13318 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Argininosuccinate Synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	0	0
			3066	1963	533	561	9			
1	B	386	Total	C	N	O	S	0	0	0
			3066	1963	533	561	9			
1	C	386	Total	C	N	O	S	0	0	0
			3066	1963	533	561	9			
1	D	386	Total	C	N	O	S	0	0	0
			3066	1963	533	561	9			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is ADENOSINE MONOPHOSPHATE (CCD ID: AMP) (formula: $C_{10}H_{14}N_5O_7P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
3	D	1	Total	C	N	O	P	0	0
			23	10	5	7	1		


- Molecule 4 is ARGININOSUCCINATE (CCD ID: AS1) (formula: $C_{10}H_{18}N_4O_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			20	10	4	6		
4	B	1	Total	C	N	O	0	0
			20	10	4	6		
4	C	1	Total	C	N	O	0	0
			20	10	4	6		
4	D	1	Total	C	N	O	0	0
			20	10	4	6		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	230	Total	O	0	0
			230	230		
5	B	238	Total	O	0	0
			238	238		
5	C	194	Total	O	0	0
			194	194		
5	D	200	Total	O	0	0
			200	200		



● Molecule 1: Argininosuccinate Synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	228.78Å 228.78Å 159.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 2.00 10.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-2.00) 98.8 (10.00-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 2.01Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.212 , 0.239 0.213 , 0.240	Depositor DCC
R_{free} test set	20800 reflections (9.95%)	wwPDB-VP
Wilson B-factor (Å ²)	28.7	Xtriage
Anisotropy	0.003	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.47 , 61.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.003 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13318	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, AS1, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/3135	0.95	10/4237 (0.2%)
1	B	0.46	0/3135	0.96	8/4237 (0.2%)
1	C	0.48	0/3135	0.96	9/4237 (0.2%)
1	D	0.47	0/3135	0.96	10/4237 (0.2%)
All	All	0.47	0/12540	0.96	37/16948 (0.2%)

There are no bond length outliers.

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	263	GLY	N-CA-C	9.23	126.00	114.37
1	C	263	GLY	N-CA-C	8.85	125.52	114.37
1	B	263	GLY	N-CA-C	8.83	125.50	114.37
1	D	263	GLY	N-CA-C	8.65	123.91	113.79
1	B	394	GLU	N-CA-C	-7.20	104.37	113.01

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3066	0	3052	81	0
1	B	3066	0	3052	100	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3066	0	3052	105	0
1	D	3066	0	3052	101	0
2	A	5	0	0	0	0
2	B	5	0	0	1	0
2	C	5	0	0	1	0
2	D	5	0	0	0	0
3	A	23	0	12	2	0
3	B	23	0	12	3	0
3	C	23	0	12	1	0
3	D	23	0	12	2	0
4	A	20	0	13	8	0
4	B	20	0	13	6	0
4	C	20	0	13	9	0
4	D	20	0	11	8	0
5	A	230	0	0	12	0
5	B	238	0	0	8	0
5	C	194	0	0	7	0
5	D	200	0	0	5	0
All	All	13318	0	12306	367	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 367 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:520:AS1:CB	4:A:520:AS1:CA	1.80	1.59
4:C:3520:AS1:CB	4:C:3520:AS1:CA	1.80	1.56
4:B:2520:AS1:CB	4:B:2520:AS1:CA	1.80	1.55
4:D:4520:AS1:CB	4:D:4520:AS1:CA	1.81	1.54
1:A:151:LYS:H	1:A:151:LYS:HD2	0.99	1.11

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/400 (95%)	373 (98%)	6 (2%)	1 (0%)	36	35
1	B	380/400 (95%)	369 (97%)	11 (3%)	0	100	100
1	C	380/400 (95%)	372 (98%)	8 (2%)	0	100	100
1	D	380/400 (95%)	372 (98%)	7 (2%)	1 (0%)	36	35
All	All	1520/1600 (95%)	1486 (98%)	32 (2%)	2 (0%)	48	46

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	343	LYS
1	D	343	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/319 (97%)	297 (96%)	13 (4%)	26	25
1	B	310/319 (97%)	294 (95%)	16 (5%)	21	18
1	C	310/319 (97%)	296 (96%)	14 (4%)	24	23
1	D	310/319 (97%)	292 (94%)	18 (6%)	18	15
All	All	1240/1276 (97%)	1179 (95%)	61 (5%)	22	20

5 of 61 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	372	ASP
1	D	262	VAL
1	C	180	HIS
1	D	239	LEU
1	D	394	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	360	GLN
1	D	180	HIS
1	C	98	HIS
1	D	323	GLN
1	D	37	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	AMP	C	3510	-	25,25,25	2.69	16 (64%)	37,38,38	1.99	15 (40%)
2	SO4	D	4540	-	4,4,4	1.69	1 (25%)	6,6,6	0.86	0
4	AS1	A	520	-	18,19,19	3.88	5 (27%)	19,24,24	7.17	11 (57%)
4	AS1	D	4520	-	18,19,19	3.91	5 (27%)	19,24,24	7.78	13 (68%)
4	AS1	C	3520	-	18,19,19	3.80	5 (27%)	19,24,24	7.11	11 (57%)
4	AS1	B	2520	-	18,19,19	3.88	6 (33%)	19,24,24	7.09	13 (68%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	2540	-	4,4,4	1.79	1 (25%)	6,6,6	0.80	0
3	AMP	D	4510	-	25,25,25	2.78	13 (52%)	37,38,38	1.95	15 (40%)
2	SO4	A	540	-	4,4,4	1.52	1 (25%)	6,6,6	0.91	0
3	AMP	A	510	-	25,25,25	2.86	15 (60%)	37,38,38	1.94	13 (35%)
3	AMP	B	2510	-	25,25,25	2.93	18 (72%)	37,38,38	2.04	14 (37%)
2	SO4	C	3540	-	4,4,4	1.73	1 (25%)	6,6,6	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AMP	C	3510	-	-	1/10/26/26	0/3/3/3
4	AS1	D	4520	-	1/1/6/7	9/21/23/23	-
4	AS1	C	3520	-	1/1/6/7	10/21/23/23	-
4	AS1	B	2520	-	1/1/6/7	10/21/23/23	-
3	AMP	D	4510	-	-	0/10/26/26	0/3/3/3
4	AS1	A	520	-	1/1/6/7	10/21/23/23	-
3	AMP	A	510	-	-	0/10/26/26	0/3/3/3
3	AMP	B	2510	-	-	1/10/26/26	0/3/3/3

The worst 5 of 87 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	4520	AS1	CB-CA	13.28	1.81	1.53
4	A	520	AS1	CB-CA	12.89	1.80	1.53
4	B	2520	AS1	CB-CA	12.89	1.80	1.53
4	C	3520	AS1	CB-CA	12.77	1.80	1.53
4	A	520	AS1	CA-N1	-6.33	1.32	1.45

The worst 5 of 105 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	4520	AS1	C2-C1-N2	30.76	165.14	110.67
4	A	520	AS1	C2-C1-N2	27.64	159.61	110.67
4	B	2520	AS1	C2-C1-N2	27.39	159.17	110.67
4	C	3520	AS1	C2-C1-N2	27.36	159.12	110.67
4	A	520	AS1	C1-N2-C	7.18	138.34	121.86

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	520	AS1	C4
4	B	2520	AS1	C4
4	C	3520	AS1	C4
4	D	4520	AS1	C4

5 of 41 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	520	AS1	N2-C1-C2-C3
4	A	520	AS1	C2-C1-N2-C
4	A	520	AS1	N2-C-N1-CA
4	A	520	AS1	N3-C-N1-CA
4	A	520	AS1	CB-CA-N1-C


There are no ring outliers.

10 monomers are involved in 40 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	3510	AMP	1	0
4	A	520	AS1	8	0
4	D	4520	AS1	8	0
4	C	3520	AS1	9	0
4	B	2520	AS1	6	0
2	B	2540	SO4	1	0
3	D	4510	AMP	2	0
3	A	510	AMP	2	0
3	B	2510	AMP	3	0
2	C	3540	SO4	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand AMP C 3510




Ligand AS1 A 520









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/400 (96%)	-0.26	6 (1%) 70 70	20, 27, 49, 78	0
1	B	386/400 (96%)	-0.31	7 (1%) 67 67	20, 27, 47, 65	0
1	C	386/400 (96%)	0.02	23 (5%) 27 26	21, 29, 55, 72	0
1	D	386/400 (96%)	-0.13	12 (3%) 51 50	21, 30, 52, 72	0
All	All	1544/1600 (96%)	-0.17	48 (3%) 51 50	20, 29, 51, 78	0

The worst 5 of 48 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	395	ARG	5.0
1	C	395	ARG	5.0
1	D	395	ARG	4.6
1	B	395	ARG	4.2
1	C	165	PRO	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	AMP	A	510	23/23	0.70	0.20	36,45,68,68	23
3	AMP	C	3510	23/23	0.70	0.18	41,47,69,70	23
3	AMP	B	2510	23/23	0.71	0.18	31,41,60,63	23
3	AMP	D	4510	23/23	0.71	0.20	40,47,69,70	23
4	AS1	C	3520	20/20	0.72	0.22	23,35,54,54	0
4	AS1	A	520	20/20	0.73	0.21	24,36,53,54	0
4	AS1	D	4520	20/20	0.74	0.22	23,39,56,58	0
4	AS1	B	2520	20/20	0.75	0.22	22,36,59,61	0
2	SO4	B	2540	5/5	0.92	0.10	37,40,41,42	0
2	SO4	A	540	5/5	0.95	0.08	41,43,44,44	0
2	SO4	C	3540	5/5	0.95	0.09	43,44,45,45	0
2	SO4	D	4540	5/5	0.96	0.07	44,45,46,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around AMP C 3510:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around AMP B 2510:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AMP D 4510:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AS1 C 3520:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AS1 A 520:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around AS1 D 4520:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around AS1 B 2520:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.