



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 05:05 PM UTC

PDB ID : 6J2I / pdb_00006j2i
Title : Crystal structure of bat (Pteropus Alecto) MHC class I Ptal-N*01:01 in complex with H17N10 influenza-like virus-derivrd peptide H17N10-NP
Authors : Lu, D.; Liu, K.F.; Yue, C.; Lu, Q.; Cheng, H.; Chai, Y.; Qi, J.X.; Gao, G.F.; Liu, W.J.
Deposited on : 2019-01-01
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	2% 89% 11%
1	D	280	11% 84% 15% .
2	B	99	2% 91% 9%
2	E	99	3% 91% 8% .
3	C	9	44% 67% 33%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	9	56% 56% 33% 11%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6573 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ptal-N*01:01.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	0	0
			2288	1426	419	436	7			
1	D	280	Total	C	N	O	S	0	0	0
			2288	1426	419	436	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP A0A125R585
A	-1	SER	-	expression tag	UNP A0A125R585
A	0	HIS	-	expression tag	UNP A0A125R585
D	-2	GLY	-	expression tag	UNP A0A125R585
D	-1	SER	-	expression tag	UNP A0A125R585
D	0	HIS	-	expression tag	UNP A0A125R585

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	0	0
			827	527	140	157	3			
2	E	99	Total	C	N	O	S	0	0	0
			827	527	140	157	3			

- Molecule 3 is a protein called H17N10-NP.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			77	49	10	18			
3	F	9	Total	C	N	O	0	0	0
			76	49	10	17			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	74	Total 74	O 74	0	0
4	B	28	Total 28	O 28	0	0
4	D	55	Total 55	O 55	0	0
4	E	33	Total 33	O 33	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ptal-N*01:01




- Molecule 1: Ptal-N*01:01




- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin



- Molecule 3: H17N10-NP



● Molecule 3: H17N10-NP



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	100.94Å 104.64Å 180.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.32 – 2.30 36.32 – 2.31	Depositor EDS
% Data completeness (in resolution range)	97.0 (36.32-2.30) 97.0 (36.32-2.31)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 2.31Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.213 , 0.253 0.216 , 0.256	Depositor DCC
R_{free} test set	2106 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 27.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.063 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6573	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	4/2355 (0.2%)	0.43	0/3206
1	D	0.52	7/2355 (0.3%)	0.47	0/3206
2	B	0.16	0/850	0.36	0/1150
2	E	0.29	0/850	0.44	0/1150
3	C	0.17	0/78	0.49	0/101
3	F	0.15	0/77	0.48	0/101
All	All	0.40	11/6565 (0.2%)	0.44	0/8914

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	28	VAL	C-O	-7.74	1.15	1.24
1	D	29	ASP	C-O	-7.40	1.14	1.24
1	A	30	ASP	C-O	-6.85	1.14	1.24
1	D	274	THR	C-O	-6.41	1.16	1.24
1	A	28	VAL	C-O	-6.24	1.17	1.24
1	D	27	TYR	C-O	-6.11	1.16	1.23
1	A	27	TYR	C-O	-6.01	1.16	1.23
1	A	29	ASP	C-O	-5.72	1.16	1.24
1	D	275	LEU	C-O	-5.34	1.17	1.23
1	D	30	ASP	C-O	-5.22	1.16	1.24
1	D	27	TYR	N-CA	-5.10	1.39	1.46

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2288	0	2113	19	0
1	D	2288	0	2113	31	0
2	B	827	0	790	6	0
2	E	827	0	790	10	0
3	C	77	0	68	2	0
3	F	76	0	68	6	0
4	A	74	0	0	3	0
4	B	28	0	0	1	0
4	D	55	0	0	1	0
4	E	33	0	0	1	0
All	All	6573	0	5942	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:10:TYR:HA	2:E:99:MET:HE2	1.69	0.72
1:D:158:ARG:HB3	3:F:5:GLU:HG2	1.74	0.69
1:A:153:ALA:O	1:A:158:ARG:NH1	2.30	0.65
1:A:28:VAL:HG11	1:A:182:LEU:HD13	1.80	0.63
1:D:107:GLY:HA3	1:D:109:HIS:CE1	2.35	0.61
1:D:194:HIS:NE2	1:D:202:VAL:HG21	2.20	0.56
1:D:223:ASP:CG	1:D:259:ARG:HH11	2.12	0.56
1:D:19:GLU:HG3	1:D:20:PRO:HD2	1.87	0.55
1:D:0:HIS:HB3	1:D:108:PRO:HG3	1.89	0.55
1:A:85:ARG:HH21	1:A:92:GLU:HG3	1.70	0.54
1:D:85:ARG:HG3	1:D:90:GLN:HB2	1.88	0.54
1:A:0:HIS:HB3	1:A:108:PRO:HG3	1.89	0.53
2:B:9:VAL:O	2:B:99:MET:HE2	2.08	0.52
1:A:74:ALA:O	1:A:78:ARG:HD3	2.10	0.52
1:D:28:VAL:HG11	1:D:182:LEU:HD13	1.92	0.51
1:D:223:ASP:OD2	1:D:259:ARG:NH1	2.43	0.51
1:D:158:ARG:HG2	3:F:5:GLU:CD	2.36	0.51
1:D:130:ASN:OD1	1:D:137:THR:OG1	2.26	0.51
1:A:119:LEU:HD22	3:C:9:LEU:HD21	1.91	0.50
1:D:273:LEU:HD13	1:D:275:LEU:HD11	1.93	0.50
1:A:85:ARG:NH2	1:A:92:GLU:HG3	2.27	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:TYR:OH	1:A:100:ARG:HD2	2.12	0.50
1:A:20:PRO:HB2	1:A:78:ARG:HG3	1.93	0.49
2:B:49:VAL:HG21	4:B:105:HOH:O	2.12	0.49
3:F:9:LEU:H	3:F:9:LEU:HD12	1.76	0.49
1:D:227:GLN:HG3	1:D:230:GLU:HB2	1.95	0.48
1:D:233:LEU:HD11	1:D:246:LYS:HE3	1.94	0.48
2:B:22:PHE:CE2	2:B:69:GLU:HG2	2.50	0.47
2:E:99:MET:HE3	4:E:107:HOH:O	2.14	0.47
1:D:77:TYR:OH	1:D:100:ARG:HD2	2.15	0.46
1:A:130:ASN:ND2	1:A:135:SER:OG	2.45	0.46
1:D:256:GLU:O	1:D:259:ARG:HB2	2.16	0.46
1:A:51:TRP:O	1:A:54:LEU:HG	2.15	0.46
1:A:214:GLU:OE2	4:A:301:HOH:O	2.20	0.46
1:D:135:SER:HB2	4:D:317:HOH:O	2.16	0.46
2:E:24:ASN:HB3	2:E:65:LEU:HD11	1.98	0.45
1:D:40:ASN:HB2	1:D:43:PRO:HB3	1.98	0.45
1:D:146:THR:HG23	3:F:9:LEU:HA	1.99	0.45
1:D:237:ARG:HH22	2:E:99:MET:HG2	1.81	0.45
1:D:158:ARG:CB	3:F:5:GLU:HG2	2.44	0.44
2:E:10:TYR:CA	2:E:99:MET:HE2	2.43	0.44
1:A:84:VAL:HA	1:A:87:TYR:CD2	2.52	0.44
3:C:3:GLU:CD	3:C:5:GLU:HB2	2.43	0.44
1:D:223:ASP:CG	1:D:259:ARG:NH1	2.74	0.44
1:A:-2:GLY:N	4:A:309:HOH:O	2.49	0.43
2:B:25:CYS:HB2	2:B:39:LEU:HD21	2.00	0.43
1:D:149:LYS:HE3	3:F:9:LEU:C	2.43	0.43
2:B:6:LYS:HD2	2:B:6:LYS:HA	1.78	0.43
1:A:233:LEU:HD11	1:A:246:LYS:HE3	2.01	0.43
1:A:90:GLN:HG2	1:D:277:TRP:CE2	2.53	0.43
1:D:120:ALA:HB2	2:E:60:TRP:CE2	2.54	0.43
1:A:120:ALA:HB2	2:B:60:TRP:CE2	2.54	0.42
1:D:207:TRP:HH2	2:E:99:MET:SD	2.42	0.42
1:A:212:TYR:OH	4:A:302:HOH:O	2.21	0.41
1:D:235:GLU:HG2	2:E:8:GLN:NE2	2.35	0.41
1:D:124:ALA:HB2	2:E:1:ILE:HD12	2.03	0.41
1:D:179:ARG:HG3	1:D:183:LEU:HD12	2.03	0.41
1:D:56:GLU:HG3	1:D:60:PRO:HA	2.02	0.40
1:A:9:TYR:HB2	1:A:100:ARG:HB3	2.04	0.40
1:D:108:PRO:HD2	1:D:109:HIS:CE1	2.56	0.40
2:E:6:LYS:O	2:E:27:VAL:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/280 (99%)	274 (99%)	4 (1%)	0	100	100
1	D	278/280 (99%)	275 (99%)	3 (1%)	0	100	100
2	B	97/99 (98%)	95 (98%)	2 (2%)	0	100	100
2	E	97/99 (98%)	94 (97%)	3 (3%)	0	100	100
3	C	7/9 (78%)	5 (71%)	2 (29%)	0	100	100
3	F	7/9 (78%)	7 (100%)	0	0	100	100
All	All	764/776 (98%)	750 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/234 (100%)	234 (100%)	0	100	100
1	D	234/234 (100%)	234 (100%)	0	100	100
2	B	94/94 (100%)	94 (100%)	0	100	100
2	E	94/94 (100%)	93 (99%)	1 (1%)	65	81
3	C	8/8 (100%)	8 (100%)	0	100	100
3	F	8/8 (100%)	5 (62%)	3 (38%)	0	0
All	All	672/672 (100%)	668 (99%)	4 (1%)	78	89

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	99	MET
3	F	7	TYR
3	F	8	SER
3	F	9	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	GLN
1	A	191	HIS
1	A	227	GLN
1	D	191	HIS
1	D	227	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	280/280 (100%)	0.24	7 (2%) 58 60	17, 35, 55, 67	0
1	D	280/280 (100%)	0.58	30 (10%) 11 12	16, 38, 66, 83	0
2	B	99/99 (100%)	0.23	2 (2%) 65 66	18, 35, 54, 63	0
2	E	99/99 (100%)	0.25	3 (3%) 52 54	17, 34, 55, 68	0
3	C	9/9 (100%)	1.62	4 (44%) 0 0	33, 49, 65, 66	0
3	F	9/9 (100%)	2.61	5 (55%) 0 0	32, 41, 67, 73	0
All	All	776/776 (100%)	0.40	51 (6%) 24 26	16, 36, 62, 83	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	7	TYR	4.9
3	F	5	GLU	4.7
2	B	1	ILE	4.5
1	D	2	PHE	4.2
3	C	7	TYR	4.1
1	D	224	GLY	4.1
3	F	8	SER	3.7
2	E	1	ILE	3.7
3	F	9	LEU	3.6
3	F	6	GLY	3.5
2	E	99	MET	3.5
1	D	54	LEU	3.5
1	D	50	PRO	3.2
1	D	254	SER	3.1
1	D	113	LEU	3.1
3	C	5	GLU	3.1
1	D	109	HIS	3.0
1	D	-2	GLY	3.0
1	D	252	VAL	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	51	TRP	2.9
1	D	198	SER	2.8
1	A	-2	GLY	2.8
1	D	253	PRO	2.8
1	D	197	ILE	2.7
1	D	57	GLN	2.7
1	A	254	SER	2.6
1	D	200	ARG	2.6
1	D	255	GLY	2.6
1	D	42	SER	2.6
1	A	16	GLY	2.5
1	A	252	VAL	2.4
1	D	41	ALA	2.4
1	D	110	GLY	2.3
1	D	158	ARG	2.3
2	B	99	MET	2.3
1	D	-1	SER	2.3
3	C	8	SER	2.3
1	D	251	VAL	2.3
1	D	199	ASP	2.2
1	D	141	LEU	2.2
1	D	60	PRO	2.2
1	D	61	GLN	2.2
1	A	82	ASP	2.1
1	D	55	VAL	2.1
1	A	89	ASN	2.1
1	D	225	GLU	2.1
3	C	6	GLY	2.1
1	D	78	ARG	2.1
1	D	235	GLU	2.0
1	A	54	LEU	2.0
2	E	15	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.