



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 06:06 AM UTC

PDB ID : 1J34 / pdb_00001j34
Title : Crystal Structure of Mg(II)-and Ca(II)-bound Gla Domain of Factor IX Complexed with Binding Protein
Authors : Shikamoto, Y.; Morita, T.; Fujimoto, Z.; Mizuno, H.
Deposited on : 2003-01-20
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	129	2% 80% 19% .
2	B	123	3% 84% 15% .
3	C	46	2% 87% 11% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2934 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called coagulation factor IX-binding protein A chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	129	Total	C	N	O	S	0	0	0
			1030	648	170	205	7			

- Molecule 2 is a protein called coagulation factor IX-binding protein B chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	123	Total	C	N	O	S	0	0	0
			1017	648	173	185	11			

- Molecule 3 is a protein called Coagulation factor IX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	46	Total	C	N	O	S	0	0	0
			433	260	65	106	2			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	407	CGU	GLU	modified residue	UNP P00741
C	408	CGU	GLU	modified residue	UNP P00741
C	415	CGU	GLU	modified residue	UNP P00741
C	417	CGU	GLU	modified residue	UNP P00741
C	420	CGU	GLU	modified residue	UNP P00741
C	421	CGU	GLU	modified residue	UNP P00741
C	426	CGU	GLU	modified residue	UNP P00741
C	427	CGU	GLU	modified residue	UNP P00741
C	430	CGU	GLU	modified residue	UNP P00741
C	433	CGU	GLU	modified residue	UNP P00741
C	436	CGU	GLU	modified residue	UNP P00741
C	440	CGU	GLU	modified residue	UNP P00741

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Ca 1	0	0
4	B	1	Total 1	Ca 1	0	0
4	C	5	Total 5	Ca 5	0	0

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	3	Total 3	Mg 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	174	Total 174	O 174	0	0
6	B	151	Total 151	O 151	0	0
6	C	119	Total 119	O 119	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: coagulation factor IX-binding protein A chain



- Molecule 2: coagulation factor IX-binding protein B chain



- Molecule 3: Coagulation factor IX



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	128.55Å 37.20Å 62.63Å 90.00° 103.56° 90.00°	Depositor
Resolution (Å)	19.63 – 1.55 19.63 – 1.55	Depositor EDS
% Data completeness (in resolution range)	90.1 (19.63-1.55) 90.2 (19.63-1.55)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 1.55Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.183 , 0.212 0.183 , 0.212	Depositor DCC
R_{free} test set	3811 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å ²)	14.0	Xtriage
Anisotropy	0.627	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2934	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CGU, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1057	0.93	4/1428 (0.3%)
2	B	0.31	0/1054	0.81	3/1427 (0.2%)
3	C	0.29	0/287	0.66	0/366
All	All	0.32	0/2398	0.85	7/3221 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	95	ILE	N-CA-C	-11.62	91.84	108.58
1	A	39	LEU	N-CA-C	-6.37	101.69	110.35
1	A	77	LYS	N-CA-C	-6.15	105.60	113.23
2	B	316	GLN	N-CA-C	-6.03	100.51	109.86
2	B	239	LEU	N-CA-C	-5.71	102.59	110.35
2	B	271	VAL	N-CA-C	5.55	119.81	112.04
1	A	107	LYS	N-CA-C	5.07	117.19	111.11

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1030	0	950	19	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1017	0	902	12	0
3	C	433	0	340	4	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	5	0	0	0	0
5	C	3	0	0	0	0
6	A	174	0	0	2	0
6	B	151	0	0	1	0
6	C	119	0	0	1	0
All	All	2934	0	2192	31	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:ASN:ND2	3:C:446:VAL:HG13	1.95	0.82
2:B:270:ASN:HD22	2:B:294:SER:HB3	1.57	0.70
1:A:61:ASN:HD21	1:A:63:LYS:HB2	1.58	0.68
1:A:57:GLU:HG3	1:A:58:ASN:ND2	2.13	0.64
6:A:633:HOH:O	3:C:446:VAL:HG21	1.99	0.62
3:C:417:CGU:O	3:C:423:CYS:HB3	2.03	0.59
1:A:65:TYR:O	1:A:123:ASN:HB3	2.04	0.58
1:A:52:ALA:O	1:A:56:THR:HG23	2.05	0.57
2:B:218:SER:HB3	2:B:259:PHE:CE2	2.43	0.53
2:B:261:HIS:HE1	6:C:654:HOH:O	1.92	0.53
2:B:253:LYS:HB2	2:B:253:LYS:NZ	2.24	0.52
1:A:73:GLN:NE2	1:A:73:GLN:HA	2.25	0.51
2:B:286:ARG:CZ	2:B:286:ARG:HA	2.41	0.50
1:A:95:ILE:HG13	2:B:305:LYS:HB3	1.94	0.49
2:B:310:ALA:HB3	2:B:313:MET:HG3	1.93	0.49
1:A:113:LYS:HE3	6:A:1016:HOH:O	2.13	0.48
1:A:94:TRP:HA	2:B:306:TRP:HB2	1.95	0.47
1:A:110:GLY:O	1:A:112:ARG:HG3	2.15	0.47
1:A:61:ASN:ND2	1:A:63:LYS:HB2	2.28	0.45
1:A:73:GLN:HA	1:A:73:GLN:HE21	1.82	0.44
1:A:19:LEU:O	1:A:124:PRO:HA	2.18	0.44
2:B:203:PRO:HB2	2:B:206:TRP:CD1	2.53	0.43
1:A:18:LYS:O	1:A:18:LYS:HG3	2.19	0.43
1:A:97:ALA:HA	1:A:118:TYR:CD1	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:90:SER:HB3	6:B:798:HOH:O	2.18	0.42
1:A:18:LYS:NZ	1:A:18:LYS:HB2	2.35	0.42
2:B:222:ASN:HA	2:B:313:MET:O	2.19	0.42
1:A:67:TRP:HB2	1:A:125:PHE:HB3	2.03	0.41
1:A:70:LEU:HB2	2:B:279:TRP:CZ3	2.56	0.41
2:B:205:ASP:OD1	2:B:205:ASP:N	2.51	0.41
3:C:407:CGU:O	3:C:410:VAL:HG22	2.21	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:GLU:OE1	1:A:57:GLU:OE1[2_555]	1.64	0.56

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/129 (98%)	120 (94%)	7 (6%)	0	100	100
2	B	121/123 (98%)	116 (96%)	4 (3%)	1 (1%)	16	4
3	C	32/46 (70%)	31 (97%)	1 (3%)	0	100	100
All	All	280/298 (94%)	267 (95%)	12 (4%)	1 (0%)	30	13

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	301	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	111/111 (100%)	110 (99%)	1 (1%)	70	51
2	B	106/106 (100%)	104 (98%)	2 (2%)	50	22
3	C	31/31 (100%)	31 (100%)	0	100	100
All	All	248/248 (100%)	245 (99%)	3 (1%)	63	40

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	LYS
2	B	253	LYS
2	B	286	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	HIS
1	A	61	ASN
1	A	73	GLN
2	B	257	GLN
2	B	270	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CGU	C	408	4,3	9,11,12	1.05	0	10,14,16	0.84	0
3	CGU	C	407	4,3	9,11,12	1.06	0	10,14,16	0.93	1 (10%)
3	CGU	C	420	5,3	9,11,12	1.03	0	10,14,16	0.78	0
3	CGU	C	430	5,4,3	9,11,12	1.04	0	10,14,16	0.69	0
3	CGU	C	426	5,3	9,11,12	1.03	0	10,14,16	0.79	0
3	CGU	C	436	5,3	9,11,12	1.08	0	10,14,16	0.99	0
3	CGU	C	427	4,3	9,11,12	1.05	0	10,14,16	0.94	1 (10%)
3	CGU	C	440	5,3	9,11,12	1.01	0	10,14,16	0.88	0
3	CGU	C	417	4,3	9,11,12	1.04	0	10,14,16	1.03	0
3	CGU	C	415	5,3	9,11,12	1.07	0	10,14,16	0.80	0
3	CGU	C	421	4,3	9,11,12	1.00	0	10,14,16	1.02	0
3	CGU	C	433	3	9,11,12	1.12	0	10,14,16	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CGU	C	408	4,3	-	2/13/14/16	-
3	CGU	C	407	4,3	-	2/13/14/16	-
3	CGU	C	420	5,3	-	3/13/14/16	-
3	CGU	C	430	5,4,3	-	3/13/14/16	-
3	CGU	C	426	5,3	-	3/13/14/16	-
3	CGU	C	436	5,3	-	0/13/14/16	-
3	CGU	C	427	4,3	-	2/13/14/16	-
3	CGU	C	440	5,3	-	0/13/14/16	-
3	CGU	C	417	4,3	-	2/13/14/16	-
3	CGU	C	415	5,3	-	2/13/14/16	-
3	CGU	C	421	4,3	-	0/13/14/16	-
3	CGU	C	433	3	-	2/13/14/16	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	427	CGU	CB-CG-CD2	-2.04	108.97	113.11
3	C	407	CGU	CB-CG-CD2	-2.01	109.03	113.11

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	407	CGU	OE22-CD2-CG-CB
3	C	433	CGU	OE21-CD2-CG-CD1
3	C	433	CGU	OE22-CD2-CG-CD1
3	C	407	CGU	OE21-CD2-CG-CB
3	C	408	CGU	OE21-CD2-CG-CB
3	C	408	CGU	OE22-CD2-CG-CB
3	C	417	CGU	OE21-CD2-CG-CB
3	C	427	CGU	OE21-CD2-CG-CB
3	C	427	CGU	OE22-CD2-CG-CB
3	C	415	CGU	OE11-CD1-CG-CB
3	C	415	CGU	OE12-CD1-CG-CB
3	C	417	CGU	OE22-CD2-CG-CB
3	C	420	CGU	OE11-CD1-CG-CB
3	C	420	CGU	OE12-CD1-CG-CB
3	C	426	CGU	OE11-CD1-CG-CB
3	C	426	CGU	OE12-CD1-CG-CB
3	C	430	CGU	OE21-CD2-CG-CB
3	C	430	CGU	OE22-CD2-CG-CB
3	C	420	CGU	OE11-CD1-CG-CD2
3	C	426	CGU	OE12-CD1-CG-CD2
3	C	430	CGU	OE21-CD2-CG-CD1

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	407	CGU	1	0
3	C	417	CGU	1	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	129/129 (100%)	0.13	3 (2%) 61 69	13, 18, 33, 47	0
2	B	123/123 (100%)	0.24	4 (3%) 49 57	13, 20, 31, 39	0
3	C	34/46 (73%)	0.24	1 (2%) 53 62	13, 19, 32, 38	0
All	All	286/298 (95%)	0.19	8 (2%) 55 63	13, 20, 32, 47	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	60	GLN	3.3
2	B	323	ALA	3.0
1	A	1	ASP	3.0
3	C	446	VAL	2.7
2	B	201	ASP	2.4
2	B	286	ARG	2.2
1	A	61	ASN	2.1
2	B	285	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CGU	C	433	12/13	0.92	0.07	15,17,26,26	0
3	CGU	C	421	12/13	0.93	0.07	20,23,25,25	0
3	CGU	C	415	12/13	0.93	0.07	18,20,21,22	0
3	CGU	C	407	12/13	0.94	0.07	18,21,23,23	0
3	CGU	C	420	12/13	0.94	0.06	21,22,23,23	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CGU	C	440	12/13	0.94	0.06	15,16,17,18	0
3	CGU	C	417	12/13	0.95	0.06	16,17,18,19	0
3	CGU	C	430	12/13	0.96	0.05	12,14,16,16	0
3	CGU	C	426	12/13	0.96	0.05	12,12,14,15	0
3	CGU	C	427	12/13	0.96	0.05	12,14,16,18	0
3	CGU	C	436	12/13	0.97	0.04	13,14,17,17	0
3	CGU	C	408	12/13	0.97	0.05	14,17,18,19	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MG	C	507	1/1	0.95	0.06	23,23,23,23	0
4	CA	C	506	1/1	0.96	0.05	28,28,28,28	0
4	CA	B	512	1/1	0.97	0.04	20,20,20,20	0
4	CA	C	503	1/1	0.98	0.03	15,15,15,15	0
5	MG	C	501	1/1	0.98	0.06	11,11,11,11	0
4	CA	C	505	1/1	0.98	0.04	19,19,19,19	0
4	CA	C	502	1/1	0.99	0.03	14,14,14,14	0
4	CA	C	504	1/1	0.99	0.03	15,15,15,15	0
5	MG	C	508	1/1	0.99	0.03	15,15,15,15	0
4	CA	A	511	1/1	1.00	0.02	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.