



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:37 PM UTC

PDB ID : 8J3S / pdb_00008j3s
Title : Complex structure of human cytomegalovirus protease and a macrocyclic peptide ligand
Authors : Yoshida, S.; Sako, Y.; Nikaido, E.; Ueda, T.; Kozono, I.; Ichihashi, Y.; Nakahashi, A.; Onishi, M.; Yamatsu, Y.; Kato, T.; Nishikawa, J.; Tachibana, Y.
Deposited on : 2023-04-18
Resolution : 3.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	
1	B	263	
1	C	263	
1	D	263	
2	E	13	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	13	 100%
2	G	13	 100%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6452 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Assemblin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	220	Total	C	N	O	S	0	0	0
			1675	1055	300	315	5			
1	B	215	Total	C	N	O	S	0	0	0
			1604	1010	283	307	4			
1	C	208	Total	C	N	O	S	0	0	0
			1539	977	270	288	4			
1	D	190	Total	C	N	O	S	0	0	0
			1281	807	225	246	3			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	initiating methionine	UNP A0A0G2TMR2
A	-5	HIS	-	expression tag	UNP A0A0G2TMR2
A	-4	HIS	-	expression tag	UNP A0A0G2TMR2
A	-3	HIS	-	expression tag	UNP A0A0G2TMR2
A	-2	HIS	-	expression tag	UNP A0A0G2TMR2
A	-1	HIS	-	expression tag	UNP A0A0G2TMR2
A	0	HIS	-	expression tag	UNP A0A0G2TMR2
A	143	GLN	ALA	engineered mutation	UNP A0A0G2TMR2
B	-6	MET	-	initiating methionine	UNP A0A0G2TMR2
B	-5	HIS	-	expression tag	UNP A0A0G2TMR2
B	-4	HIS	-	expression tag	UNP A0A0G2TMR2
B	-3	HIS	-	expression tag	UNP A0A0G2TMR2
B	-2	HIS	-	expression tag	UNP A0A0G2TMR2
B	-1	HIS	-	expression tag	UNP A0A0G2TMR2
B	0	HIS	-	expression tag	UNP A0A0G2TMR2
B	143	GLN	ALA	engineered mutation	UNP A0A0G2TMR2
C	-6	MET	-	initiating methionine	UNP A0A0G2TMR2
C	-5	HIS	-	expression tag	UNP A0A0G2TMR2
C	-4	HIS	-	expression tag	UNP A0A0G2TMR2
C	-3	HIS	-	expression tag	UNP A0A0G2TMR2
C	-2	HIS	-	expression tag	UNP A0A0G2TMR2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	HIS	-	expression tag	UNP A0A0G2TMR2
C	0	HIS	-	expression tag	UNP A0A0G2TMR2
C	143	GLN	ALA	engineered mutation	UNP A0A0G2TMR2
D	-6	MET	-	initiating methionine	UNP A0A0G2TMR2
D	-5	HIS	-	expression tag	UNP A0A0G2TMR2
D	-4	HIS	-	expression tag	UNP A0A0G2TMR2
D	-3	HIS	-	expression tag	UNP A0A0G2TMR2
D	-2	HIS	-	expression tag	UNP A0A0G2TMR2
D	-1	HIS	-	expression tag	UNP A0A0G2TMR2
D	0	HIS	-	expression tag	UNP A0A0G2TMR2
D	143	GLN	ALA	engineered mutation	UNP A0A0G2TMR2




- Molecule 2 is a protein called PHE-ILE-THR-GLY-HIS-TYR-TRP-VAL-ARG-PHE-LEU-P
RO-CYS-GLY.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	13	Total	C	N	O	S	0	0	0
			110	74	19	16	1			
2	F	13	Total	C	N	O	S	0	0	0
			110	74	19	16	1			
2	G	13	Total	C	N	O	S	0	0	0
			110	74	19	16	1			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	14	Total	O	0	0
			14	14		
3	B	3	Total	O	0	0
			3	3		
3	C	5	Total	O	0	0
			5	5		
3	G	1	Total	O	0	0
			1	1		

- Molecule 1: Assemblin





● Molecule 2: PHE-ILE-THR-GLY-HIS-TYR-TRP-VAL-ARG-PHE-LEU-PRO-CYS-GLY



● Molecule 2: PHE-ILE-THR-GLY-HIS-TYR-TRP-VAL-ARG-PHE-LEU-PRO-CYS-GLY



There are no outlier residues recorded for this chain.

● Molecule 2: PHE-ILE-THR-GLY-HIS-TYR-TRP-VAL-ARG-PHE-LEU-PRO-CYS-GLY



There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	79.18Å 81.65Å 198.09Å 90.00° 91.13° 90.00°	Depositor
Resolution (Å)	50.00 – 3.09 50.00 – 3.09	Depositor EDS
% Data completeness (in resolution range)	97.3 (50.00-3.09) 97.3 (50.00-3.09)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 3.06Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.223 , 0.289 0.222 , 0.284	Depositor DCC
R_{free} test set	1143 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å ²)	62.2	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 52.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.029 for k,h,-l 0.027 for -k,-h,-l 0.029 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6452	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CCS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/1709	1.04	2/2322 (0.1%)
1	B	0.62	0/1634	1.03	0/2220
1	C	0.60	0/1571	1.05	3/2137 (0.1%)
1	D	0.60	0/1302	1.02	1/1776 (0.1%)
2	E	0.54	0/106	0.71	0/145
2	F	0.50	0/106	0.55	0/145
2	G	0.62	0/106	0.82	0/145
All	All	0.60	0/6534	1.02	6/8890 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	110	GLY	CA-C-N	8.41	128.84	120.52
1	C	110	GLY	C-N-CA	8.41	128.84	120.52
1	C	252	SER	N-CA-C	6.93	119.61	109.07
1	A	226	VAL	CB-CA-C	-6.04	104.24	111.97
1	A	206	ALA	N-CA-C	5.85	117.35	110.97
1	D	81	GLY	N-CA-C	5.40	117.01	111.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1675	0	1622	13	0
1	B	1604	0	1527	13	0
1	C	1539	0	1432	11	0
1	D	1281	0	1087	4	1
2	E	110	0	94	3	0
2	F	110	0	93	0	0
2	G	110	0	94	0	0
3	A	14	0	0	0	0
3	B	3	0	0	0	0
3	C	5	0	0	0	0
3	G	1	0	0	0	0
All	All	6452	0	5949	42	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:163:VAL:CG2	1:B:234:ARG:HG3	2.21	0.71
1:B:163:VAL:HG21	1:B:234:ARG:HG3	1.78	0.64
1:C:163:VAL:HG12	1:C:238:LEU:HD13	1.79	0.64
1:A:163:VAL:HG13	1:A:238:LEU:HD13	1.80	0.63
1:A:109:ARG:HH11	1:A:109:ARG:HG3	1.66	0.61
1:B:226:VAL:HA	1:B:229:LEU:HD12	1.87	0.55
1:B:163:VAL:HG22	1:B:234:ARG:HG3	1.89	0.54
1:B:170:LEU:HD22	1:B:242:LYS:HA	1.90	0.54
1:B:182:GLN:HA	1:B:192:ARG:HH21	1.73	0.53
1:B:243:GLN:HA	1:B:248:THR:HG23	1.90	0.53
1:B:221:LEU:HA	1:B:224:ASN:HD22	1.73	0.52
1:C:20:LEU:HD21	1:C:84:CYS:SG	2.50	0.51
1:B:245:VAL:HG23	1:B:247:VAL:HG23	1.93	0.51
1:D:14:VAL:HG22	1:D:175:ARG:HE	1.76	0.50
1:A:251:GLU:OE1	1:B:183:ARG:NH2	2.44	0.50
1:C:226:VAL:HA	1:C:229:LEU:HD12	1.95	0.49
1:A:131:LEU:HD13	1:A:171:ALA:HB2	1.94	0.49
1:C:184:PHE:HB2	1:C:187:LEU:HD12	1.94	0.49
1:A:193:ASP:OD1	1:A:196:ARG:NH1	2.43	0.48
2:E:1:PHE:HB2	2:E:11:LEU:HB3	1.96	0.48
1:C:97:VAL:HG13	1:C:123:PHE:CD2	2.49	0.48
1:A:20:LEU:HD21	1:A:84:CYS:SG	2.54	0.47
1:A:96:ILE:HG21	1:A:221:LEU:HD12	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:185:PRO:HD2	1:D:246:GLY:HA3	1.97	0.47
1:A:109:ARG:HH11	1:A:109:ARG:CG	2.28	0.46
1:B:243:GLN:HA	1:B:248:THR:CG2	2.46	0.46
2:E:3:THR:OG1	2:E:8:VAL:O	2.34	0.45
1:C:12:THR:HA	1:C:13:PRO:HD3	1.85	0.45
1:B:62:ASN:HD22	1:B:224:ASN:CG	2.25	0.44
1:C:20:LEU:HD13	1:C:133:LEU:HD13	1.99	0.44
1:A:35:PRO:HD2	1:A:38:VAL:HB	1.99	0.44
1:A:187:LEU:O	1:A:192:ARG:NH1	2.51	0.44
1:C:113:SER:HA	1:C:114:PRO:HA	1.67	0.43
1:C:61:ILE:HD13	1:C:160:LEU:HB2	2.00	0.42
1:C:97:VAL:HG13	1:C:123:PHE:HD2	1.84	0.42
1:B:221:LEU:O	1:B:224:ASN:HB2	2.20	0.41
1:A:227:ASP:HB3	2:E:7:TRP:CD1	2.55	0.41
1:A:249:GLU:O	1:A:251:GLU:N	2.54	0.41
1:C:95:GLU:OE2	1:C:98:ARG:NH2	2.52	0.40
1:D:113:SER:HA	1:D:114:PRO:HA	1.73	0.40
1:A:113:SER:HA	1:A:114:PRO:HA	1.85	0.40
1:D:184:PHE:HA	1:D:185:PRO:HD3	1.85	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:218:SER:OG	1:D:218:SER:OG[2_656]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	212/263 (81%)	198 (93%)	13 (6%)	1 (0%)	24 57

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	207/263 (79%)	194 (94%)	13 (6%)	0	100	100
1	C	200/263 (76%)	187 (94%)	12 (6%)	1 (0%)	24	57
1	D	176/263 (67%)	157 (89%)	17 (10%)	2 (1%)	11	39
2	E	11/13 (85%)	11 (100%)	0	0	100	100
2	F	11/13 (85%)	9 (82%)	2 (18%)	0	100	100
2	G	11/13 (85%)	10 (91%)	1 (9%)	0	100	100
All	All	828/1091 (76%)	766 (92%)	58 (7%)	4 (0%)	24	57

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	114	PRO
1	D	111	PRO
1	A	247	VAL
1	C	114	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/223 (78%)	153 (87%)	22 (13%)	4	19
1	B	163/223 (73%)	150 (92%)	13 (8%)	11	37
1	C	148/223 (66%)	140 (95%)	8 (5%)	20	50
1	D	107/223 (48%)	99 (92%)	8 (8%)	12	39
2	E	9/11 (82%)	9 (100%)	0	100	100
2	F	9/11 (82%)	9 (100%)	0	100	100
2	G	9/11 (82%)	9 (100%)	0	100	100
All	All	620/925 (67%)	569 (92%)	51 (8%)	10	36

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	LEU
1	A	37	ASP
1	A	59	LEU
1	A	79	ARG
1	A	85	LEU
1	A	108	SER
1	A	109	ARG
1	A	112	VAL
1	A	131	LEU
1	A	163	VAL
1	A	165	ARG
1	A	180	VAL
1	A	181	THR
1	A	183	ARG
1	A	186	ASP
1	A	207	VAL
1	A	222	LEU
1	A	226	VAL
1	A	233	GLU
1	A	238	LEU
1	A	249	GLU
1	A	252	SER
1	B	37	ASP
1	B	59	LEU
1	B	80	ASP
1	B	85	LEU
1	B	131	LEU
1	B	163	VAL
1	B	180	VAL
1	B	183	ARG
1	B	192	ARG
1	B	207	VAL
1	B	222	LEU
1	B	226	VAL
1	B	238	LEU
1	C	37	ASP
1	C	59	LEU
1	C	84	CYS
1	C	102	GLU
1	C	103	LYS
1	C	131	LEU
1	C	163	VAL
1	C	180	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	59	LEU
1	D	98	ARG
1	D	102	GLU
1	D	131	LEU
1	D	181	THR
1	D	199	TRP
1	D	222	LEU
1	D	229	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	B	25	GLN
1	B	62	ASN
1	B	198	GLN
1	B	224	ASN
1	C	71	HIS
1	C	76	GLN
1	D	62	ASN
1	D	63	HIS
1	D	224	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CCS	E	13	2	7,8,10	0.80	0	4,8,12	0.62	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CCS	G	13	2	7,8,10	0.85	0	4,8,12	0.69	0
2	CCS	F	13	2	7,8,10	0.83	0	4,8,12	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CCS	E	13	2	-	0/4/7/10	-
2	CCS	G	13	2	-	0/4/7/10	-
2	CCS	F	13	2	-	1/4/7/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	13	CCS	CA-CB-SG-CD

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	220/263 (83%)	-0.44	0 100 100	22, 34, 58, 87	0
1	B	215/263 (81%)	-0.01	3 (1%) 73 53	32, 51, 89, 111	0
1	C	208/263 (79%)	0.25	5 (2%) 59 38	37, 62, 90, 103	0
1	D	190/263 (72%)	1.07	24 (12%) 8 4	71, 105, 145, 162	0
2	E	12/13 (92%)	-0.58	0 100 100	26, 35, 44, 54	0
2	F	12/13 (92%)	-0.15	0 100 100	54, 64, 89, 96	0
2	G	12/13 (92%)	-0.06	0 100 100	37, 55, 75, 82	0
All	All	869/1091 (79%)	0.17	32 (3%) 45 25	22, 57, 123, 162	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	246	GLY	3.5
1	D	251	GLU	3.5
1	D	58	PRO	3.2
1	D	12	THR	3.2
1	B	204	SER	2.9
1	D	59	LEU	2.9
1	D	132	SER	2.9
1	D	253	TYR	2.8
1	C	12	THR	2.8
1	D	134	SER	2.7
1	D	62	ASN	2.7
1	C	154	PRO	2.6
1	D	245	VAL	2.6
1	D	192	ARG	2.6
1	C	186	ASP	2.5
1	D	80	ASP	2.4
1	B	205	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	87	CYS	2.4
1	D	73	ALA	2.4
1	D	61	ILE	2.3
1	C	249	GLU	2.3
1	D	67	ALA	2.3
1	D	235	LEU	2.3
1	D	17	GLY	2.2
1	D	129	ALA	2.2
1	D	57	LEU	2.2
1	D	154	PRO	2.2
1	C	135	SER	2.2
1	B	28	ASP	2.2
1	D	60	ASN	2.1
1	D	180	VAL	2.1
1	D	13	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CCS	G	13	9/11	0.78	0.13	78,83,93,96	0
2	CCS	F	13	9/11	0.86	0.09	94,98,102,104	0
2	CCS	E	13	9/11	0.87	0.09	57,65,69,71	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.