



wwPDB EM Validation Summary Report ⓘ

Mar 25, 2026 – 09:14 AM UTC

PDB ID : 9J35 / pdb_00009j35
EMDB ID : EMD-61106
Title : Cryo-EM structure of Arabidopsis CNGC5 in nanodisc
Authors : Wang, J.P.; Zhang, X.; Zhang, P.
Deposited on : 2024-08-07
Resolution : 2.71 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY


The reported resolution of this entry is 2.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	10297 (2.21 - 3.21)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	726	
1	B	726	
1	C	726	
1	D	726	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 16960 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable cyclic nucleotide-gated ion channel 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	523	Total	C	N	O	S	0	0
			4240	2752	710	753	25		
1	B	523	Total	C	N	O	S	0	0
			4240	2752	710	753	25		
1	C	523	Total	C	N	O	S	0	0
			4240	2752	710	753	25		
1	D	523	Total	C	N	O	S	0	0
			4240	2752	710	753	25		

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	MET	-	initiating methionine	UNP Q8RWS9
A	-7	ASP	-	expression tag	UNP Q8RWS9
A	-6	TYR	-	expression tag	UNP Q8RWS9
A	-5	LYS	-	expression tag	UNP Q8RWS9
A	-4	ASP	-	expression tag	UNP Q8RWS9
A	-3	ASP	-	expression tag	UNP Q8RWS9
A	-2	ASP	-	expression tag	UNP Q8RWS9
A	-1	ASP	-	expression tag	UNP Q8RWS9
A	0	LYS	-	expression tag	UNP Q8RWS9
B	-8	MET	-	initiating methionine	UNP Q8RWS9
B	-7	ASP	-	expression tag	UNP Q8RWS9
B	-6	TYR	-	expression tag	UNP Q8RWS9
B	-5	LYS	-	expression tag	UNP Q8RWS9
B	-4	ASP	-	expression tag	UNP Q8RWS9
B	-3	ASP	-	expression tag	UNP Q8RWS9
B	-2	ASP	-	expression tag	UNP Q8RWS9
B	-1	ASP	-	expression tag	UNP Q8RWS9
B	0	LYS	-	expression tag	UNP Q8RWS9
C	-8	MET	-	initiating methionine	UNP Q8RWS9
C	-7	ASP	-	expression tag	UNP Q8RWS9
C	-6	TYR	-	expression tag	UNP Q8RWS9
C	-5	LYS	-	expression tag	UNP Q8RWS9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-4	ASP	-	expression tag	UNP Q8RWS9
C	-3	ASP	-	expression tag	UNP Q8RWS9
C	-2	ASP	-	expression tag	UNP Q8RWS9
C	-1	ASP	-	expression tag	UNP Q8RWS9
C	0	LYS	-	expression tag	UNP Q8RWS9
D	-8	MET	-	initiating methionine	UNP Q8RWS9
D	-7	ASP	-	expression tag	UNP Q8RWS9
D	-6	TYR	-	expression tag	UNP Q8RWS9
D	-5	LYS	-	expression tag	UNP Q8RWS9
D	-4	ASP	-	expression tag	UNP Q8RWS9
D	-3	ASP	-	expression tag	UNP Q8RWS9
D	-2	ASP	-	expression tag	UNP Q8RWS9
D	-1	ASP	-	expression tag	UNP Q8RWS9
D	0	LYS	-	expression tag	UNP Q8RWS9

L572	I573	W574	A575	L576	D577	P578	K579	S580	G581	V582	N583	A611	F614	ARG	ARG	LEU	HIS	SER	ARG	GLN	VAL	GLN	HIS	THR	PHE	ARG	PHE	TYR	HIS	HIS	GLN	TRP	ARG	THR	TRP	ALA	ALA	CYS	PHE	ILE	GLN	ALA	ALA	TRP	ARG	ARG	CYS	LYS	ARG	LYS	LYS	MET	GLU	ALA	GLU
ALA	GLU	ALA	ALA	ALA	VAL	SER	SER	SER	THR	ALA	GLY	PRO	SER	SER	ILE	GLY	ALA	ALA	PHE	LEU	ALA	ALA	ASN	ALA	LEU	ALA	ARG	THR	ILE	HIS	ARG	ASN	ARG	ASN	THR	LYS	ILE	ARG	ASP	LEU	VAL	VAL	LYS	LEU	GLN	LYS	PRO	PRO	GLU	PRO	ASP	PHE	THR	ALA	ASP

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	195793	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	5.587	Depositor
Minimum map value	-3.682	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.102	Depositor
Recommended contour level	0.53	Depositor
Map size (Å)	335.52, 335.52, 335.52	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.932, 0.932, 0.932	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	10/4337 (0.2%)	0.83	19/5877 (0.3%)
1	B	0.78	15/4337 (0.3%)	0.93	24/5877 (0.4%)
1	C	0.58	9/4337 (0.2%)	0.83	17/5877 (0.3%)
1	D	0.60	7/4337 (0.2%)	0.89	17/5877 (0.3%)
All	All	0.66	41/17348 (0.2%)	0.87	77/23508 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	411	LEU	C-N	21.09	1.59	1.33
1	B	236	PRO	N-CD	-17.55	1.23	1.47
1	B	411	LEU	C-N	17.14	1.54	1.33
1	B	378	LEU	C-N	15.60	1.57	1.33
1	D	512	MET	C-N	15.48	1.57	1.33

The worst 5 of 77 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	378	LEU	O-C-N	-18.70	97.37	122.43
1	D	378	LEU	CA-C-N	15.64	149.18	121.66
1	D	378	LEU	C-N-CA	15.64	149.18	121.66
1	B	378	LEU	O-C-N	-15.08	103.72	122.27
1	B	378	LEU	CA-C-N	11.83	141.84	121.14

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	414	ASN	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4240	0	4285	43	0
1	B	4240	0	4283	47	0
1	C	4240	0	4285	42	0
1	D	4240	0	4285	54	0
All	All	16960	0	17138	156	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 156 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:297:CYS:SG	1:D:303:CYS:SG	1.46	1.36
1:C:297:CYS:SG	1:C:303:CYS:SG	1.34	1.34
1:A:297:CYS:SG	1:A:303:CYS:SG	1.34	1.33
1:A:297:CYS:SG	1:A:303:CYS:CB	2.44	1.06
1:B:235:ILE:N	1:B:236:PRO:HD2	1.68	1.06

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	519/726 (72%)	480 (92%)	38 (7%)	1 (0%)	43	66
1	B	519/726 (72%)	477 (92%)	42 (8%)	0	100	100
1	C	519/726 (72%)	476 (92%)	43 (8%)	0	100	100
1	D	519/726 (72%)	488 (94%)	31 (6%)	0	100	100
All	All	2076/2904 (72%)	1921 (92%)	154 (7%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	236	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	463/631 (73%)	460 (99%)	3 (1%)	78	90
1	B	463/631 (73%)	459 (99%)	4 (1%)	70	86
1	C	463/631 (73%)	458 (99%)	5 (1%)	65	84
1	D	463/631 (73%)	461 (100%)	2 (0%)	84	92
All	All	1852/2524 (73%)	1838 (99%)	14 (1%)	70	87

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	145	THR
1	C	197	LEU
1	D	295	GLU
1	C	445	LEU
1	D	234	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	416	GLN
1	D	215	ASN
1	C	487	HIS
1	D	312	ASN
1	B	204	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-61106. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 189




Y Index: 189




Z Index: 211


6.3.2 Raw map



X Index: 162



Y Index: 198




Z Index: 210


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.53. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 97 nm³; this corresponds to an approximate mass of 88 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of $0.369 \text{ \AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.369 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.71	-	-
Author-provided FSC curve	2.71	2.99	2.75
Unmasked-calculated*	3.10	3.58	3.16

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.10 differs from the reported value 2.71 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-61106 and PDB model 9J35. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.53 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.53).

9.4 Atom inclusion [i](#)



At the recommended contour level, 87% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.53) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8160	0.5940
A	0.8180	0.5930
B	0.8150	0.5940
C	0.8140	0.5930
D	0.8180	0.5970

