



wwPDB EM Validation Summary Report ⓘ

Mar 10, 2026 – 03:17 AM UTC

PDB ID : 9J36 / pdb_00009j36
EMDB ID : EMD-61107
Title : Cryo-EM structure of Arabidopsis CNGC5
Authors : Wang, J.P.; Zhang, P.; Zhang, X.
Deposited on : 2024-08-07
Resolution : 2.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	7115 (2.00 - 3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	726	 12% 57% 14% • 28%
1	B	726	 12% 56% 15% • 28%
1	C	726	 12% 58% 13% • 28%
1	D	726	 12% 57% 14% • 28%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 16960 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable cyclic nucleotide-gated ion channel 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	523	Total	C	N	O	S	0	0
			4240	2752	710	753	25		
1	C	523	Total	C	N	O	S	0	0
			4240	2752	710	753	25		
1	D	523	Total	C	N	O	S	0	0
			4240	2752	710	753	25		
1	B	523	Total	C	N	O	S	0	0
			4240	2752	710	753	25		


There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	MET	-	initiating methionine	UNP Q8RWS9
A	-7	ASP	-	expression tag	UNP Q8RWS9
A	-6	TYR	-	expression tag	UNP Q8RWS9
A	-5	LYS	-	expression tag	UNP Q8RWS9
A	-4	ASP	-	expression tag	UNP Q8RWS9
A	-3	ASP	-	expression tag	UNP Q8RWS9
A	-2	ASP	-	expression tag	UNP Q8RWS9
A	-1	ASP	-	expression tag	UNP Q8RWS9
A	0	LYS	-	expression tag	UNP Q8RWS9
C	-8	MET	-	initiating methionine	UNP Q8RWS9
C	-7	ASP	-	expression tag	UNP Q8RWS9
C	-6	TYR	-	expression tag	UNP Q8RWS9
C	-5	LYS	-	expression tag	UNP Q8RWS9
C	-4	ASP	-	expression tag	UNP Q8RWS9
C	-3	ASP	-	expression tag	UNP Q8RWS9
C	-2	ASP	-	expression tag	UNP Q8RWS9
C	-1	ASP	-	expression tag	UNP Q8RWS9
C	0	LYS	-	expression tag	UNP Q8RWS9
D	-8	MET	-	initiating methionine	UNP Q8RWS9
D	-7	ASP	-	expression tag	UNP Q8RWS9
D	-6	TYR	-	expression tag	UNP Q8RWS9
D	-5	LYS	-	expression tag	UNP Q8RWS9


Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-4	ASP	-	expression tag	UNP Q8RWS9
D	-3	ASP	-	expression tag	UNP Q8RWS9
D	-2	ASP	-	expression tag	UNP Q8RWS9
D	-1	ASP	-	expression tag	UNP Q8RWS9
D	0	LYS	-	expression tag	UNP Q8RWS9
B	-8	MET	-	initiating methionine	UNP Q8RWS9
B	-7	ASP	-	expression tag	UNP Q8RWS9
B	-6	TYR	-	expression tag	UNP Q8RWS9
B	-5	LYS	-	expression tag	UNP Q8RWS9
B	-4	ASP	-	expression tag	UNP Q8RWS9
B	-3	ASP	-	expression tag	UNP Q8RWS9
B	-2	ASP	-	expression tag	UNP Q8RWS9
B	-1	ASP	-	expression tag	UNP Q8RWS9
B	0	LYS	-	expression tag	UNP Q8RWS9



• Molecule 1: Probable cyclic nucleotide-gated ion channel 5




ASN
ALA
LEU
ARG
THR
ILE
HIS
ARG
ASN
ARG
ASN
ARG
THR
LYS
ILE
ARG
ASP
LEU
VAL
LYS
LEU
GLN
LYS
PRO
PRO
GLU
PRO
PHE
THR
ALA
ASP

- Molecule 1: Probable cyclic nucleotide-gated ion channel 5

Chain B: 

MET
ASP
TYR
LYS
ASP
GLY
ASP
ASP
LYS
LYS
MET
ALA
GLY
LYS
ARG
GLU
ASN
PHE
VAL
ARG
VAL
ASP
LYS
LEU
ASP
SER
SER
SER
VAL
ALA
PHE
GLN
ASN
TYR
ALA
SER
ASN
PHE
SER
GLY
GLN
LEU
HIS
PRO
ILE
HIS
ALA
SER
ASN
GLU
THR
SER
ARG
PHE
LYS



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	204715	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	5.919	Depositor
Minimum map value	-3.980	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.099	Depositor
Recommended contour level	0.47	Depositor
Map size (Å)	335.52, 335.52, 335.52	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.932, 0.932, 0.932	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	10/4337 (0.2%)	0.83	15/5877 (0.3%)
1	B	0.53	9/4337 (0.2%)	0.83	14/5877 (0.2%)
1	C	0.52	10/4337 (0.2%)	0.87	12/5877 (0.2%)
1	D	0.53	9/4337 (0.2%)	0.83	14/5877 (0.2%)
All	All	0.53	38/17348 (0.2%)	0.84	55/23508 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
1	C	0	1
1	D	0	1
All	All	0	5

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	384	GLY	C-N	-8.74	1.21	1.33
1	A	384	GLY	C-N	-8.61	1.21	1.33
1	A	217	SER	C-N	-8.48	1.22	1.33
1	B	384	GLY	C-N	-8.08	1.22	1.33
1	A	383	GLN	CA-C	-7.49	1.42	1.52

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	384	GLY	CA-C-N	17.69	147.20	121.02
1	C	384	GLY	C-N-CA	17.69	147.20	121.02
1	C	384	GLY	O-C-N	-14.52	103.83	122.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	158	GLN	N-CA-C	-13.09	96.94	113.16
1	B	158	GLN	N-CA-C	-13.06	96.97	113.16

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	217	SER	Mainchain
1	A	256	ALA	Peptide
1	B	256	ALA	Peptide
1	C	256	ALA	Peptide
1	D	256	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4240	0	4282	99	0
1	B	4240	0	4282	104	0
1	C	4240	0	4283	98	0
1	D	4240	0	4282	103	0
All	All	16960	0	17129	370	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 370 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:547:THR:CG2	1:A:589:ARG:HG2	1.63	1.28
1:B:547:THR:CG2	1:B:589:ARG:HG2	1.63	1.28
1:C:547:THR:CG2	1:C:589:ARG:HG2	1.63	1.27
1:D:547:THR:CG2	1:D:589:ARG:HG2	1.63	1.27
1:C:155:MET:O	1:C:159:LEU:HG	1.34	1.27

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	519/726 (72%)	484 (93%)	35 (7%)	0	100	100
1	B	519/726 (72%)	485 (93%)	34 (7%)	0	100	100
1	C	519/726 (72%)	485 (93%)	34 (7%)	0	100	100
1	D	519/726 (72%)	485 (93%)	34 (7%)	0	100	100
All	All	2076/2904 (72%)	1939 (93%)	137 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	463/631 (73%)	459 (99%)	4 (1%)	70	87
1	B	463/631 (73%)	459 (99%)	4 (1%)	70	87
1	C	463/631 (73%)	459 (99%)	4 (1%)	70	87
1	D	463/631 (73%)	459 (99%)	4 (1%)	70	87
All	All	1852/2524 (73%)	1836 (99%)	16 (1%)	68	87

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	381	LEU
1	B	224	GLN
1	D	109	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	109	LEU
1	C	501	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	438	GLN
1	B	154	HIS
1	B	613	GLN
1	B	340	ASN
1	D	613	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-61107. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 172




Y Index: 172




Z Index: 139


6.3.2 Raw map



X Index: 188



Y Index: 188




Z Index: 143


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) ⓘ


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.47. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 94 nm^3 ; this corresponds to an approximate mass of 85 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.400 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.400 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.50	-	-
Author-provided FSC curve	2.50	2.92	2.61
Unmasked-calculated*	3.29	3.81	3.33

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.29 differs from the reported value 2.5 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-61107 and PDB model 9J36. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay [i](#)


This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.47).

9.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.47) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7390	0.5310
A	0.7400	0.5340
B	0.7410	0.5320
C	0.7340	0.5240
D	0.7420	0.5340

