



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 02:13 PM UTC

PDB ID : 1J4T / pdb_00001j4t
Title : Structure of Artocarpin: a Lectin with Mannose Specificity (Form 2)
Authors : Pratap, J.V.; Jeyaprakash, A.A.; Rani, P.G.; Sekar, K.; Surolia, A.; Vijayan, M.
Deposited on : 2001-10-30
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	5% 51% 34% 10%
1	B	149	7% 46% 36% 10%
1	C	149	12% 42% 36% 10%
1	D	149	5% 49% 36% 11%
1	E	149	9% 53% 31% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	149	
1	G	149	
1	H	149	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	AYA	A	1	X	X	-	-
1	AYA	B	1	X	X	X	-
1	AYA	C	1	X	X	-	-
1	AYA	E	1	X	-	-	-
1	AYA	F	1	-	X	-	-
1	AYA	G	1	X	-	-	-
1	AYA	H	1	X	X	-	-

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9922 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Artocarpin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	149	Total	C	N	O	S	0	0	0
			1132	727	182	222	1			
1	B	149	Total	C	N	O	S	0	0	0
			1132	727	182	222	1			
1	C	149	Total	C	N	O	S	0	0	0
			1132	727	182	222	1			
1	D	149	Total	C	N	O	S	0	0	0
			1132	727	182	222	1			
1	E	149	Total	C	N	O	S	0	0	0
			1128	724	181	222	1			
1	F	149	Total	C	N	O	S	0	0	0
			1128	724	181	222	1			
1	G	149	Total	C	N	O	S	0	0	0
			1132	727	182	222	1			
1	H	149	Total	C	N	O	S	0	0	0
			1132	727	182	222	1			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	AYA	ALA	modified residue	UNP Q7M1T4
A	9	SER	PRO	conflict	UNP Q7M1T4
A	20	GLU	ASP	conflict	UNP Q7M1T4
A	49	ASP	GLU	conflict	UNP Q7M1T4
A	70	LYS	ARG	conflict	UNP Q7M1T4
A	84	GLY	ALA	conflict	UNP Q7M1T4
A	145	ILE	VAL	conflict	UNP Q7M1T4
A	148	SER	ALA	conflict	UNP Q7M1T4
B	1	AYA	ALA	modified residue	UNP Q7M1T4
B	9	SER	PRO	conflict	UNP Q7M1T4
B	20	GLU	ASP	conflict	UNP Q7M1T4
B	49	ASP	GLU	conflict	UNP Q7M1T4
B	70	LYS	ARG	conflict	UNP Q7M1T4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	84	GLY	ALA	conflict	UNP Q7M1T4
B	145	ILE	VAL	conflict	UNP Q7M1T4
B	148	SER	ALA	conflict	UNP Q7M1T4
C	1	AYA	ALA	modified residue	UNP Q7M1T4
C	9	SER	PRO	conflict	UNP Q7M1T4
C	20	GLU	ASP	conflict	UNP Q7M1T4
C	49	ASP	GLU	conflict	UNP Q7M1T4
C	70	LYS	ARG	conflict	UNP Q7M1T4
C	84	GLY	ALA	conflict	UNP Q7M1T4
C	145	ILE	VAL	conflict	UNP Q7M1T4
C	148	SER	ALA	conflict	UNP Q7M1T4
D	1	AYA	ALA	modified residue	UNP Q7M1T4
D	9	SER	PRO	conflict	UNP Q7M1T4
D	20	GLU	ASP	conflict	UNP Q7M1T4
D	49	ASP	GLU	conflict	UNP Q7M1T4
D	70	LYS	ARG	conflict	UNP Q7M1T4
D	84	GLY	ALA	conflict	UNP Q7M1T4
D	145	ILE	VAL	conflict	UNP Q7M1T4
D	148	SER	ALA	conflict	UNP Q7M1T4
E	1	AYA	ALA	modified residue	UNP Q7M1T4
E	9	SER	PRO	conflict	UNP Q7M1T4
E	20	GLU	ASP	conflict	UNP Q7M1T4
E	49	ASP	GLU	conflict	UNP Q7M1T4
E	70	LYS	ARG	conflict	UNP Q7M1T4
E	84	GLY	ALA	conflict	UNP Q7M1T4
E	145	ILE	VAL	conflict	UNP Q7M1T4
E	148	SER	ALA	conflict	UNP Q7M1T4
F	1	AYA	ALA	modified residue	UNP Q7M1T4
F	9	SER	PRO	conflict	UNP Q7M1T4
F	20	GLU	ASP	conflict	UNP Q7M1T4
F	49	ASP	GLU	conflict	UNP Q7M1T4
F	70	LYS	ARG	conflict	UNP Q7M1T4
F	84	GLY	ALA	conflict	UNP Q7M1T4
F	145	ILE	VAL	conflict	UNP Q7M1T4
F	148	SER	ALA	conflict	UNP Q7M1T4
G	1	AYA	ALA	modified residue	UNP Q7M1T4
G	9	SER	PRO	conflict	UNP Q7M1T4
G	20	GLU	ASP	conflict	UNP Q7M1T4
G	49	ASP	GLU	conflict	UNP Q7M1T4
G	70	LYS	ARG	conflict	UNP Q7M1T4
G	84	GLY	ALA	conflict	UNP Q7M1T4
G	145	ILE	VAL	conflict	UNP Q7M1T4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	148	SER	ALA	conflict	UNP Q7M1T4
H	1	AYA	ALA	modified residue	UNP Q7M1T4
H	9	SER	PRO	conflict	UNP Q7M1T4
H	20	GLU	ASP	conflict	UNP Q7M1T4
H	49	ASP	GLU	conflict	UNP Q7M1T4
H	70	LYS	ARG	conflict	UNP Q7M1T4
H	84	GLY	ALA	conflict	UNP Q7M1T4
H	145	ILE	VAL	conflict	UNP Q7M1T4
H	148	SER	ALA	conflict	UNP Q7M1T4


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	118	Total O 118 118	0	0
2	B	123	Total O 123 123	0	0
2	C	121	Total O 121 121	0	0
2	D	121	Total O 121 121	0	0
2	E	93	Total O 93 93	0	0
2	F	115	Total O 115 115	0	0
2	G	102	Total O 102 102	0	0
2	H	81	Total O 81 81	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Artocarpin




• Molecule 1: Artocarpin



• Molecule 1: Artocarpin



• Molecule 1: Artocarpin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.69Å 72.19Å 92.63Å 90.00° 101.15° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40 20.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.40) 82.3 (20.00-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.91 (at 2.41Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.191 , 0.258 0.203 , 0.260	Depositor DCC
R_{free} test set	1536 reflections (4.18%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.22 , 72.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9922	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 64.61 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.0588e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AYA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.33	3/1155 (0.3%)	2.79	154/1569 (9.8%)
1	B	1.34	6/1155 (0.5%)	2.86	151/1569 (9.6%)
1	C	1.35	5/1155 (0.4%)	2.92	158/1569 (10.1%)
1	D	1.24	7/1155 (0.6%)	3.14	156/1569 (9.9%)
1	E	1.20	5/1151 (0.4%)	3.09	148/1565 (9.5%)
1	F	1.63	9/1151 (0.8%)	2.90	151/1565 (9.6%)
1	G	1.59	7/1155 (0.6%)	2.72	138/1569 (8.8%)
1	H	1.24	7/1155 (0.6%)	3.00	144/1569 (9.2%)
All	All	1.37	49/9232 (0.5%)	2.93	1200/12544 (9.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	8
1	B	1	4
1	C	1	8
1	D	0	5
1	E	1	5
1	F	0	7
1	G	1	5
1	H	1	4
All	All	6	46

The worst 5 of 49 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	70	LYS	CD-CE	32.69	2.50	1.52
1	F	70	LYS	CD-CE	30.66	2.44	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	33	TYR	C-N	18.56	1.59	1.33
1	A	34	LYS	C-N	16.98	1.58	1.33
1	C	33	TYR	C-N	15.29	1.55	1.33

The worst 5 of 1200 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	34	LYS	O-C-N	-43.70	64.47	122.59
1	D	34	LYS	O-C-N	-33.93	77.46	122.59
1	H	9	SER	O-C-N	-33.53	77.99	122.59
1	H	89	LEU	O-C-N	-31.42	91.07	122.18
1	B	8	GLY	O-C-N	-27.23	100.28	122.88

5 of 6 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	1	AYA	CA
1	B	1	AYA	CA
1	C	1	AYA	CA
1	E	1	AYA	CA
1	G	1	AYA	CA

5 of 46 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	AYA	Mainchain
1	A	34	LYS	Mainchain
1	A	42	VAL	Mainchain
1	A	8	GLY	Peptide,Mainchain
1	A	9	SER	Mainchain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1132	0	1084	72	0
1	B	1132	0	1084	60	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1132	0	1084	59	0
1	D	1132	0	1081	86	0
1	E	1128	0	1071	50	0
1	F	1128	0	1068	70	0
1	G	1132	0	1082	57	0
1	H	1132	0	1083	75	0
2	A	118	0	0	7	0
2	B	123	0	0	9	0
2	C	121	0	0	4	0
2	D	121	0	0	7	0
2	E	93	0	0	4	0
2	F	115	0	0	6	0
2	G	102	0	0	6	0
2	H	81	0	0	6	0
All	All	9922	0	8637	493	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 493 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:8:GLY:HA2	1:D:123:PRO:HG2	1.22	1.18
1:D:18:TRP:HZ3	1:D:136:THR:HG23	0.98	1.10
1:D:18:TRP:CZ3	1:D:136:THR:HG23	1.89	1.07
1:A:61:PRO:HD2	2:A:181:HOH:O	1.57	1.04
1:E:60:LEU:HD11	1:E:89:LEU:HD11	1.47	0.97

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	147/149 (99%)	133 (90%)	12 (8%)	2 (1%)	9	13
1	B	147/149 (99%)	136 (92%)	8 (5%)	3 (2%)	6	7
1	C	147/149 (99%)	131 (89%)	11 (8%)	5 (3%)	3	2
1	D	147/149 (99%)	132 (90%)	12 (8%)	3 (2%)	6	7
1	E	147/149 (99%)	131 (89%)	12 (8%)	4 (3%)	4	4
1	F	147/149 (99%)	135 (92%)	11 (8%)	1 (1%)	18	28
1	G	147/149 (99%)	139 (95%)	6 (4%)	2 (1%)	9	13
1	H	147/149 (99%)	128 (87%)	17 (12%)	2 (1%)	9	13
All	All	1176/1192 (99%)	1065 (91%)	89 (8%)	22 (2%)	6	8

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	SER
1	A	110	GLY
1	B	9	SER
1	B	110	GLY
1	C	9	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	120/122 (98%)	97 (81%)	23 (19%)	1	2
1	B	120/122 (98%)	93 (78%)	27 (22%)	1	1
1	C	120/122 (98%)	90 (75%)	30 (25%)	0	1
1	D	120/122 (98%)	93 (78%)	27 (22%)	1	1
1	E	119/122 (98%)	91 (76%)	28 (24%)	1	1
1	F	119/122 (98%)	96 (81%)	23 (19%)	1	2
1	G	120/122 (98%)	96 (80%)	24 (20%)	1	2
1	H	120/122 (98%)	85 (71%)	35 (29%)	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	958/976 (98%)	741 (77%)	217 (23%)	1 1

5 of 217 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	69	LEU
1	F	70	LYS
1	H	87	SER
1	E	95	VAL
1	E	141	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	104	ASN
1	H	104	ASN
1	E	56	HIS
1	G	47	ASN
1	E	47	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	AYA	H	1	1	6,7,8	1.44	1 (16%)	6,8,10	10.82	6 (100%)
1	AYA	F	1	1	6,7,8	1.65	1 (16%)	6,8,10	10.99	6 (100%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	AYA	C	1	1	6,7,8	1.26	1 (16%)	6,8,10	11.40	5 (83%)
1	AYA	A	1	1	6,7,8	1.08	0	6,8,10	11.15	5 (83%)
1	AYA	G	1	1	6,7,8	2.44	1 (16%)	6,8,10	10.67	3 (50%)
1	AYA	E	1	1	6,7,8	0.81	0	6,8,10	10.73	3 (50%)
1	AYA	D	1	1	6,7,8	1.81	1 (16%)	6,8,10	11.61	3 (50%)
1	AYA	B	1	1	6,7,8	1.68	2 (33%)	6,8,10	10.46	6 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	AYA	H	1	1	1/1/2/4	5/5/6/8	-
1	AYA	F	1	1	-	5/5/6/8	-
1	AYA	C	1	1	1/1/2/4	5/5/6/8	-
1	AYA	A	1	1	1/1/2/4	5/5/6/8	-
1	AYA	G	1	1	1/1/2/4	4/5/6/8	-
1	AYA	E	1	1	1/1/2/4	4/5/6/8	-
1	AYA	D	1	1	-	4/5/6/8	-
1	AYA	B	1	1	1/1/2/4	3/5/6/8	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	1	AYA	CA-N	-5.46	1.40	1.46
1	D	1	AYA	CA-N	3.62	1.50	1.46
1	F	1	AYA	CA-N	3.22	1.50	1.46
1	H	1	AYA	OT-CT	2.90	1.29	1.23
1	B	1	AYA	CA-N	-2.82	1.43	1.46

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	1	AYA	CB-CA-N	23.84	136.60	109.68
1	C	1	AYA	CB-CA-N	23.71	136.45	109.68
1	G	1	AYA	CB-CA-N	23.22	135.89	109.68
1	A	1	AYA	CB-CA-N	23.01	135.66	109.68
1	E	1	AYA	CB-CA-N	21.68	134.16	109.68

5 of 6 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	1	AYA	CA
1	B	1	AYA	CA
1	C	1	AYA	CA
1	E	1	AYA	CA
1	G	1	AYA	CA

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1	AYA	CB-CA-N-CT
1	A	1	AYA	OT-CT-N-CA
1	A	1	AYA	CM-CT-N-CA
1	A	1	AYA	O-C-CA-CB
1	B	1	AYA	CB-CA-N-CT

There are no ring outliers.

7 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	H	1	AYA	2	0
1	F	1	AYA	2	0
1	C	1	AYA	1	0
1	A	1	AYA	1	0
1	E	1	AYA	1	0
1	D	1	AYA	1	0
1	B	1	AYA	4	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	F	2
1	C	1
1	D	1
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	22:SER	C	23:TYR	N	1.20
1	D	8:GLY	C	9:SER	N	1.20
1	F	9:SER	C	10:TRP	N	1.20
1	B	89:LEU	C	90:ALA	N	1.18
1	F	90:ALA	C	91:THR	N	1.14

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	148/149 (99%)	-0.63	2 (1%) 73 69	7, 20, 37, 60	0
1	B	148/149 (99%)	-0.54	0 100 100	11, 24, 37, 46	0
1	C	148/149 (99%)	-0.62	1 (0%) 84 81	7, 22, 34, 57	0
1	D	148/149 (99%)	-0.38	2 (1%) 73 69	11, 26, 44, 66	0
1	E	148/149 (99%)	-0.54	2 (1%) 73 69	8, 23, 47, 61	0
1	F	148/149 (99%)	-0.41	4 (2%) 56 52	9, 26, 44, 63	0
1	G	148/149 (99%)	-0.69	1 (0%) 84 81	10, 21, 33, 52	0
1	H	148/149 (99%)	-0.48	2 (1%) 73 69	11, 26, 40, 60	0
All	All	1184/1192 (99%)	-0.54	14 (1%) 76 73	7, 24, 43, 66	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	91	THR	4.8
1	C	23	TYR	4.1
1	F	91	THR	3.6
1	A	91	THR	3.1
1	D	89	LEU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	AYA	A	1	8/9	0.91	0.07	24,26,31,35	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	AYA	E	1	8/9	0.91	0.06	12,20,22,22	0
1	AYA	B	1	8/9	0.92	0.08	30,33,37,38	0
1	AYA	F	1	8/9	0.92	0.06	17,23,32,32	0
1	AYA	D	1	8/9	0.93	0.07	22,28,32,34	0
1	AYA	H	1	8/9	0.93	0.06	23,25,30,32	0
1	AYA	G	1	8/9	0.94	0.05	22,26,31,31	0
1	AYA	C	1	8/9	0.95	0.06	18,23,29,34	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.