



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 02:22 PM UTC

PDB ID : 6J4N / pdb_00006j4n
Title : Structure of papua new guinea MBL-1(PNGM-1) native
Authors : Hong, M.K.; Park, K.S.; Jeon, J.H.; Lee, J.H.; Park, Y.S.; Lee, S.H.; Kang, L.W.
Deposited on : 2019-01-10
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	372	4% 86% 13% .
1	B	372	4% 86% 13% .
1	C	372	8% 85% 13% .
1	D	372	5% 89% 9% .
1	E	372	5% 83% 15% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	372	 8% 84% 15% •
1	G	372	 8% 85% 14% •
1	H	372	 12% 84% 14% •

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 24786 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metallo-beta-lactamases PNGM-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	372	Total	C	N	O	S	0	0	0
			2933	1853	506	554	20			
1	B	372	Total	C	N	O	S	0	0	0
			2933	1853	506	554	20			
1	C	372	Total	C	N	O	S	0	0	0
			2933	1853	506	554	20			
1	D	372	Total	C	N	O	S	0	0	0
			2933	1853	506	554	20			
1	E	372	Total	C	N	O	S	0	0	0
			2933	1853	506	554	20			
1	F	372	Total	C	N	O	S	0	0	0
			2933	1853	506	554	20			
1	G	372	Total	C	N	O	S	0	0	0
			2933	1853	506	554	20			
1	H	372	Total	C	N	O	S	0	0	0
			2933	1853	506	554	20			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		
2	D	2	Total	Zn	0	0
			2	2		
2	E	2	Total	Zn	0	0
			2	2		
2	F	2	Total	Zn	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	2	Total 2	Zn 2	0	0
2	H	2	Total 2	Zn 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	186	Total 186	O 186	0	0
3	B	205	Total 205	O 205	0	0
3	C	165	Total 165	O 165	0	0
3	D	181	Total 181	O 181	0	0
3	E	173	Total 173	O 173	0	0
3	F	132	Total 132	O 132	0	0
3	G	137	Total 137	O 137	0	0
3	H	127	Total 127	O 127	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Metallo-beta-lactamases PNGM-1




• Molecule 1: Metallo-beta-lactamases PNGM-1




• Molecule 1: Metallo-beta-lactamases PNGM-1





• Molecule 1: Metallo-beta-lactamases PNGM-1









• Molecule 1: Metallo-beta-lactamases PNGM-1




• Molecule 1: Metallo-beta-lactamases PNGM-1




• Molecule 1: Metallo-beta-lactamases PNGM-1



• Molecule 1: Metallo-beta-lactamases PNGM-1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	122.27Å 83.02Å 163.47Å 90.00° 110.55° 90.00°	Depositor
Resolution (Å)	45.39 – 2.10 45.39 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.1 (45.39-2.10) 98.1 (45.39-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	12.82 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.190 , 0.244 0.198 , 0.248	Depositor DCC
R_{free} test set	8773 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.012 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	24786	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 84.26 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7453e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	0/3019	1.21	2/4115 (0.0%)
1	B	1.10	2/3019 (0.1%)	1.19	1/4115 (0.0%)
1	C	1.06	1/3019 (0.0%)	1.21	2/4115 (0.0%)
1	D	1.10	2/3019 (0.1%)	1.25	5/4115 (0.1%)
1	E	1.08	1/3019 (0.0%)	1.25	7/4115 (0.2%)
1	F	1.08	0/3019	1.27	9/4115 (0.2%)
1	G	1.10	1/3019 (0.0%)	1.22	0/4115
1	H	1.10	0/3019	1.27	1/4115 (0.0%)
All	All	1.09	7/24152 (0.0%)	1.23	27/32920 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1
1	F	0	1
1	H	0	1
All	All	0	3

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	271	GLN	N-CA	6.69	1.51	1.46
1	C	271	GLN	N-CA	5.63	1.50	1.46
1	D	40	MET	N-CA	5.39	1.49	1.46
1	E	4	GLY	C-O	5.25	1.29	1.23
1	B	56	LEU	C-O	5.16	1.29	1.23

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	66	ILE	CA-C-N	7.39	127.36	120.34
1	D	66	ILE	C-N-CA	7.39	127.36	120.34
1	C	66	ILE	CA-C-N	7.04	127.03	120.34
1	C	66	ILE	C-N-CA	7.04	127.03	120.34
1	F	344	ASN	CA-C-O	-6.47	115.65	121.67

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	343	ARG	Peptide
1	F	152	ILE	Peptide
1	H	337	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2933	0	2754	36	1
1	B	2933	0	2754	33	1
1	C	2933	0	2754	34	0
1	D	2933	0	2754	24	0
1	E	2933	0	2754	43	1
1	F	2933	0	2754	43	1
1	G	2933	0	2754	33	0
1	H	2933	0	2754	42	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
2	G	2	0	0	0	0
2	H	2	0	0	0	0
3	A	186	0	0	2	0
3	B	205	0	0	4	0
3	C	165	0	0	1	0
3	D	181	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	173	0	0	3	0
3	F	132	0	0	3	0
3	G	137	0	0	1	0
3	H	127	0	0	2	0
All	All	24786	0	22032	241	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 241 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:ARG:HD2	1:F:329:ALA:O	1.53	1.05
1:C:44:ARG:HD2	1:H:329:ALA:O	1.58	1.03
1:C:329:ALA:O	1:H:44:ARG:HD2	1.61	0.99
1:B:329:ALA:O	1:E:44:ARG:HD2	1.65	0.96
1:A:329:ALA:O	1:F:44:ARG:HD2	1.67	0.95

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:347:SER:O	1:F:344:ASN:ND2[2_455]	2.02	0.18
1:B:347:SER:O	1:E:344:ASN:ND2[2_555]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/372 (100%)	354 (96%)	15 (4%)	1 (0%)	36	36
1	B	370/372 (100%)	356 (96%)	12 (3%)	2 (0%)	24	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	370/372 (100%)	348 (94%)	17 (5%)	5 (1%)	9	5
1	D	370/372 (100%)	349 (94%)	17 (5%)	4 (1%)	11	8
1	E	370/372 (100%)	349 (94%)	16 (4%)	5 (1%)	9	5
1	F	370/372 (100%)	353 (95%)	13 (4%)	4 (1%)	11	8
1	G	370/372 (100%)	357 (96%)	11 (3%)	2 (0%)	24	22
1	H	370/372 (100%)	352 (95%)	14 (4%)	4 (1%)	11	8
All	All	2960/2976 (100%)	2818 (95%)	115 (4%)	27 (1%)	14	10

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	336	SER
1	D	336	SER
1	E	338	ASP
1	E	341	PRO
1	E	343	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/302 (100%)	292 (97%)	10 (3%)	33	37
1	B	302/302 (100%)	289 (96%)	13 (4%)	26	27
1	C	302/302 (100%)	287 (95%)	15 (5%)	22	22
1	D	302/302 (100%)	289 (96%)	13 (4%)	26	27
1	E	302/302 (100%)	286 (95%)	16 (5%)	20	19
1	F	302/302 (100%)	290 (96%)	12 (4%)	28	29
1	G	302/302 (100%)	285 (94%)	17 (6%)	19	18
1	H	302/302 (100%)	289 (96%)	13 (4%)	26	27
All	All	2416/2416 (100%)	2307 (96%)	109 (4%)	24	25

5 of 109 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	254	LEU
1	F	222	LYS
1	H	183	SER
1	E	310	VAL
1	E	362	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	369	GLN
1	G	243	ASN
1	G	175	GLN
1	G	257	HIS
1	E	82	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 16 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	372/372 (100%)	0.07	15 (4%)	42	44	9, 18, 49, 87	0
1	B	372/372 (100%)	0.02	14 (3%)	44	46	7, 15, 48, 81	0
1	C	372/372 (100%)	0.28	28 (7%)	20	21	10, 22, 57, 101	0
1	D	372/372 (100%)	0.15	17 (4%)	37	39	10, 19, 50, 92	0
1	E	372/372 (100%)	0.33	20 (5%)	31	33	10, 22, 45, 100	0
1	F	372/372 (100%)	0.60	28 (7%)	20	21	12, 27, 50, 92	0
1	G	372/372 (100%)	0.66	29 (7%)	19	20	12, 27, 47, 88	0
1	H	372/372 (100%)	0.86	46 (12%)	8	8	13, 28, 54, 90	0
All	All	2976/2976 (100%)	0.37	197 (6%)	24	26	7, 23, 50, 101	0

The worst 5 of 197 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	339	LEU	7.3
1	C	341	PRO	6.4
1	D	373	GLY	6.3
1	C	373	GLY	6.2
1	D	341	PRO	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	H	402	1/1	0.98	0.03	23,23,23,23	0
2	ZN	D	401	1/1	0.99	0.07	29,29,29,29	0
2	ZN	E	401	1/1	0.99	0.04	27,27,27,27	0
2	ZN	E	402	1/1	0.99	0.02	18,18,18,18	0
2	ZN	F	401	1/1	0.99	0.03	31,31,31,31	0
2	ZN	G	401	1/1	0.99	0.07	28,28,28,28	0
2	ZN	G	402	1/1	0.99	0.02	19,19,19,19	0
2	ZN	H	401	1/1	0.99	0.05	31,31,31,31	0
2	ZN	C	402	1/1	0.99	0.02	24,24,24,24	0
2	ZN	B	402	1/1	1.00	0.01	16,16,16,16	0
2	ZN	C	401	1/1	1.00	0.05	30,30,30,30	0
2	ZN	F	402	1/1	1.00	0.03	21,21,21,21	0
2	ZN	A	401	1/1	1.00	0.05	25,25,25,25	0
2	ZN	A	402	1/1	1.00	0.01	20,20,20,20	0
2	ZN	D	402	1/1	1.00	0.01	19,19,19,19	0
2	ZN	B	401	1/1	1.00	0.06	24,24,24,24	0

6.5 Other polymers

There are no such residues in this entry.