



wwPDB EM Validation Summary Report ⓘ

Mar 25, 2026 – 05:15 AM UTC

PDB ID : 8J4F / pdb_00008j4f
EMDB ID : EMD-35975
Title : Structure of human Nav1.7 in complex with Hardwickii acid
Authors : Wu, Q.R.; Yan, N.
Deposited on : 2023-04-19
Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14081 (2.50 - 3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2028	 9% 53% 10% 37%
2	B	218	 53% 24% 21%
3	C	2	 50% 50%
3	D	2	 50% 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	2	 100%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 12419 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sodium channel protein type 9 subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1273	Total	C	N	O	S	0	0
			10259	6799	1613	1770	77		

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-39	TRP	-	expression tag	UNP Q15858
A	-38	SER	-	expression tag	UNP Q15858
A	-37	HIS	-	expression tag	UNP Q15858
A	-36	PRO	-	expression tag	UNP Q15858
A	-35	GLN	-	expression tag	UNP Q15858
A	-34	PHE	-	expression tag	UNP Q15858
A	-33	GLU	-	expression tag	UNP Q15858
A	-32	LYS	-	expression tag	UNP Q15858
A	-31	GLY	-	expression tag	UNP Q15858
A	-30	GLY	-	expression tag	UNP Q15858
A	-29	GLY	-	expression tag	UNP Q15858
A	-28	ALA	-	expression tag	UNP Q15858
A	-27	ARG	-	expression tag	UNP Q15858
A	-26	GLY	-	expression tag	UNP Q15858
A	-25	GLY	-	expression tag	UNP Q15858
A	-24	SER	-	expression tag	UNP Q15858
A	-23	GLY	-	expression tag	UNP Q15858
A	-22	GLY	-	expression tag	UNP Q15858
A	-21	GLY	-	expression tag	UNP Q15858
A	-20	SER	-	expression tag	UNP Q15858
A	-19	TRP	-	expression tag	UNP Q15858
A	-18	SER	-	expression tag	UNP Q15858
A	-17	HIS	-	expression tag	UNP Q15858
A	-16	PRO	-	expression tag	UNP Q15858
A	-15	GLN	-	expression tag	UNP Q15858
A	-14	PHE	-	expression tag	UNP Q15858
A	-13	GLU	-	expression tag	UNP Q15858
A	-12	LYS	-	expression tag	UNP Q15858

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	GLY	-	expression tag	UNP Q15858
A	-10	PHE	-	expression tag	UNP Q15858
A	-9	ASP	-	expression tag	UNP Q15858
A	-8	TYR	-	expression tag	UNP Q15858
A	-7	LYS	-	expression tag	UNP Q15858
A	-6	ASP	-	expression tag	UNP Q15858
A	-5	ASP	-	expression tag	UNP Q15858
A	-4	ASP	-	expression tag	UNP Q15858
A	-3	ASP	-	expression tag	UNP Q15858
A	-2	LYS	-	expression tag	UNP Q15858
A	-1	GLY	-	expression tag	UNP Q15858
A	0	THR	-	expression tag	UNP Q15858

- Molecule 2 is a protein called Sodium channel subunit beta-1.


Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	173	Total	C	N	O	S	0	0
			1416	902	232	272	10		

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	2	Total	C	N	O	0	0
			28	16	2	10		
3	D	2	Total	C	N	O	0	0
			28	16	2	10		
3	E	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				AltConf
4	A	1	Total	C	N	O	0
			14	8	1	5	
4	A	1	Total	C	N	O	0
			14	8	1	5	
4	B	1	Total	C	N	O	0
			14	8	1	5	
4	B	1	Total	C	N	O	0
			14	8	1	5	
4	B	1	Total	C	N	O	0
			14	8	1	5	

- Molecule 5 is (4 {a} {R},5 {S},6 {R},8 {a} {R})-5-[2-(furan-3-yl)ethyl]-5,6,8 {a}-trimethyl-3,4,4 {a},6,7,8-hexahydronaphthalene-1-carboxylic acid (CCD ID: UK0) (formula: C₂₀H₂₈O₃).




Mol	Chain	Residues	Atoms			AltCon
5	A	1	Total	C	O	0
			23	20	3	

- Molecule 6 is CHOLESTEROL HEMISUCCINATE (CCD ID: Y01) (formula: $C_{31}H_{50}O_4$).




Mol	Chain	Residues	Atoms			AltConf
6	A	1	Total 35	C 31	O 4	0
6	A	1	Total 35	C 31	O 4	0
6	A	1	Total 35	C 31	O 4	0

- Molecule 7 is (3beta,14beta,17beta,25R)-3-[4-methoxy-3-(methoxymethyl)butoxy]spirost-5-en (CCD ID: 9Z9) (formula: C₃₄H₅₆O₅).



Mol	Chain	Residues	Atoms			AltConf
			Total	C	O	
7	A	1	39	34	5	0

- Molecule 8 is 1-O-OCTADECYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: LPE) (formula: C₂₆H₅₇NO₆P).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
8	A	1	25	17	1	6	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
8	A	1	Total	C	N	O	P	0
			18	10	1	6	1	
8	A	1	Total	C	N	O	P	0
			28	20	1	6	1	
8	A	1	Total	C	N	O	P	0
			25	17	1	6	1	
8	A	1	Total	C	N	O	P	0
			25	17	1	6	1	
8	A	1	Total	C	N	O	P	0
			25	17	1	6	1	
8	A	1	Total	C	N	O	P	0
			25	17	1	6	1	
8	A	1	Total	C	N	O	P	0
			25	17	1	6	1	
8	B	1	Total	C	N	O	P	0
			17	9	1	6	1	

- Molecule 9 is 1,2-DIOLEOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PCW) (formula: $C_{44}H_{85}NO_8P$).




Mol	Chain	Residues	Atoms					AltConf
9	A	1	Total	C	N	O	P	0
			50	40	1	8	1	
9	A	1	Total	C	N	O	P	0
			47	37	1	8	1	


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
9	A	1	Total	C	N	O	P	0
			44	34	1	8	1	
9	A	1	Total	C	N	O	P	0
			44	34	1	8	1	



● Molecule 2: Sodium channel subunit beta-1



GLU
TYR
LEU
ALA
ILE
THR
SER
GLU
SER
LYS
GLU
ASN
CYS
THR
GLY
VAL
GLN
VAL
ALA
GLU

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




MAG1
MAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	220477	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1500	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	3.677	Depositor
Minimum map value	-2.185	Depositor
Average map value	0.009	Depositor
Map value standard deviation	0.094	Depositor
Recommended contour level	0.55	Depositor
Map size (Å)	281.0624, 281.0624, 281.0624	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0979, 1.0979, 1.0979	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, PCW, UK0, LPE, Y01, 9Z9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/10508	0.44	0/14235
2	B	0.95	3/1442 (0.2%)	1.29	7/1949 (0.4%)
All	All	0.42	3/11950 (0.0%)	0.61	7/16184 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	95	SER	CA-CB	-5.99	1.43	1.53
2	B	105	SER	C-O	-5.50	1.16	1.23
2	B	106	ILE	C-O	-5.26	1.18	1.23

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	129	PHE	CA-C-O	-6.50	114.04	121.58
2	B	93	ASN	N-CA-C	-6.13	98.53	108.52
2	B	132	TYR	CA-CB-CG	5.76	124.26	113.90
2	B	129	PHE	CA-CB-CG	5.54	119.34	113.80
2	B	107	PHE	CA-C-O	-5.21	114.73	120.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10259	0	10490	125	0
2	B	1416	0	1380	33	0
3	C	28	0	25	1	0
3	D	28	0	25	0	0
3	E	28	0	25	0	0
4	A	28	0	26	0	0
4	B	42	0	39	2	0
5	A	23	0	0	2	0
6	A	105	0	147	1	0
7	A	39	0	0	2	0
8	A	221	0	304	7	0
8	B	17	0	19	2	0
9	A	185	0	257	7	0
All	All	12419	0	12737	163	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 163 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:51:ALA:HB2	2:B:127:LEU:HD13	1.42	1.01
1:A:1324:VAL:HG21	1:A:1455:VAL:HG21	1.75	0.68
1:A:86:LYS:HB2	1:A:102:ALA:HB3	1.76	0.67
1:A:1502:ASN:HB3	1:A:1505:GLN:HB2	1.79	0.65
1:A:202:THR:HB	1:A:213:LEU:HD13	1.79	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1261/2028 (62%)	1235 (98%)	26 (2%)	0	100	100
2	B	171/218 (78%)	168 (98%)	3 (2%)	0	100	100
All	All	1432/2246 (64%)	1403 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1135/1807 (63%)	1126 (99%)	9 (1%)	73	86
2	B	157/190 (83%)	148 (94%)	9 (6%)	18	52
All	All	1292/1997 (65%)	1274 (99%)	18 (1%)	57	80

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	136	THR
2	B	189	THR
2	B	180	TYR
1	A	1568	SER
2	B	127	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1551	ASN
2	B	61	GLN
2	B	143	HIS
1	A	1256	ASN
1	A	900	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	C	1	3,1	14,14,15	0.44	0	17,19,21	1.34	2 (11%)
3	NAG	C	2	3	14,14,15	0.42	0	17,19,21	0.97	1 (5%)
3	NAG	D	1	3,1	14,14,15	0.33	0	17,19,21	0.75	0
3	NAG	D	2	3	14,14,15	0.32	0	17,19,21	0.61	0
3	NAG	E	1	3,2	14,14,15	0.68	0	17,19,21	1.17	1 (5%)
3	NAG	E	2	3	14,14,15	0.78	0	17,19,21	1.22	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	1/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	NAG	D	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	1/6/23/26	0/1/1/1
3	NAG	E	1	3,2	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-O5-C5	3.44	116.80	112.19
3	E	1	NAG	O5-C1-C2	-2.91	106.78	111.29
3	C	1	NAG	O4-C4-C3	2.73	116.81	110.38
3	E	2	NAG	C2-N2-C7	-2.39	119.70	122.90
3	C	2	NAG	C1-O5-C5	2.27	115.23	112.19

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	2	NAG	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6
3	D	2	NAG	C1-C2-N2-C7
3	E	2	NAG	C4-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	B	301	2	14,14,15	0.43	0	17,19,21	0.98	1 (5%)
6	Y01	A	2005	-	38,38,38	0.44	0	57,57,57	0.50	0
4	NAG	B	302	2	14,14,15	0.51	0	17,19,21	0.74	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	LPE	A	2015	-	24,24,33	0.28	0	28,30,39	0.36	0
9	PCW	A	2011	-	46,46,53	0.49	0	52,54,61	0.36	0
4	NAG	A	2001	1	14,14,15	0.39	0	17,19,21	0.49	0
8	LPE	A	2014	-	24,24,33	0.29	0	28,30,39	0.32	0
5	UK0	A	2003	-	25,25,25	1.92	6 (24%)	36,38,38	1.67	7 (19%)
4	NAG	B	303	2	14,14,15	0.89	0	17,19,21	0.86	1 (5%)
8	LPE	A	2018	-	24,24,33	0.28	0	28,30,39	0.41	0
8	LPE	A	2016	-	24,24,33	0.28	0	28,30,39	0.37	0
9	PCW	A	2012	-	43,43,53	0.31	0	49,51,61	0.35	0
9	PCW	A	2008	-	49,49,53	0.32	0	55,57,61	0.33	0
6	Y01	A	2004	-	38,38,38	0.47	0	57,57,57	0.61	0
8	LPE	A	2017	-	24,24,33	0.28	0	28,30,39	0.35	0
6	Y01	A	2020	-	38,38,38	0.47	0	57,57,57	0.59	0
9	PCW	A	2019	-	43,43,53	0.39	0	49,51,61	0.35	0
8	LPE	A	2013	-	24,24,33	0.28	0	28,30,39	0.39	0
4	NAG	A	2002	1	14,14,15	0.46	0	17,19,21	1.22	1 (5%)
8	LPE	A	2010	-	27,27,33	0.27	0	31,33,39	0.33	0
8	LPE	A	2009	-	17,17,33	0.41	0	21,23,39	0.46	0
7	9Z9	A	2006	-	44,44,44	0.26	0	64,68,68	0.56	0
8	LPE	B	304	-	16,16,33	0.41	0	20,22,39	0.49	0
8	LPE	A	2007	-	24,24,33	0.41	0	25,27,39	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	B	301	2	-	0/6/23/26	0/1/1/1
6	Y01	A	2005	-	-	6/19/77/77	0/4/4/4
4	NAG	B	302	2	-	3/6/23/26	0/1/1/1
8	LPE	A	2015	-	-	3/25/25/34	-
9	PCW	A	2011	-	-	14/50/50/57	-
4	NAG	A	2001	1	-	2/6/23/26	0/1/1/1
8	LPE	A	2014	-	-	3/25/25/34	-
5	UK0	A	2003	-	-	6/10/46/46	0/3/3/3
4	NAG	B	303	2	-	2/6/23/26	0/1/1/1
8	LPE	A	2018	-	-	6/25/25/34	-
8	LPE	A	2016	-	-	5/25/25/34	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	PCW	A	2012	-	-	7/47/47/57	-
9	PCW	A	2008	-	-	10/53/53/57	-
6	Y01	A	2004	-	-	9/19/77/77	0/4/4/4
8	LPE	A	2017	-	-	2/25/25/34	-
6	Y01	A	2020	-	-	4/19/77/77	0/4/4/4
9	PCW	A	2019	-	-	16/47/47/57	-
8	LPE	A	2013	-	-	4/25/25/34	-
4	NAG	A	2002	1	-	0/6/23/26	0/1/1/1
8	LPE	A	2010	-	-	3/28/28/34	-
8	LPE	A	2009	-	-	8/18/18/34	-
7	9Z9	A	2006	-	-	4/12/100/100	0/6/6/6
8	LPE	B	304	-	-	9/17/17/34	-
8	LPE	A	2007	-	-	15/25/25/34	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	2003	UK0	C6-C7	5.95	1.43	1.33
5	A	2003	UK0	C04-C2	-3.96	1.50	1.56
5	A	2003	UK0	C5-C6	-2.57	1.43	1.50
5	A	2003	UK0	C21-C23	2.25	1.39	1.34
5	A	2003	UK0	C08-C06	-2.18	1.50	1.54

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	2003	UK0	C1-C07-C04	-4.60	109.21	114.38
4	A	2002	NAG	C1-O5-C5	4.36	118.03	112.19
5	A	2003	UK0	C18-C11-C04	-3.92	111.00	117.05
5	A	2003	UK0	C08-C06-C2	3.52	112.25	108.36
5	A	2003	UK0	C06-C7-C6	-2.61	118.95	123.26

There are no chirality outliers.

5 of 141 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	302	NAG	C8-C7-N2-C2
4	B	302	NAG	O7-C7-N2-C2
5	A	2003	UK0	C13-C04-C11-C18

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	2003	UK0	C07-C04-C11-C18
5	A	2003	UK0	C2-C04-C11-C18


There are no ring outliers.


14 monomers are involved in 21 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	302	NAG	1	0
8	A	2015	LPE	2	0
9	A	2011	PCW	1	0
8	A	2014	LPE	1	0
5	A	2003	UK0	2	0
4	B	303	NAG	1	0
8	A	2016	LPE	1	0
9	A	2012	PCW	2	0
9	A	2008	PCW	4	0
6	A	2020	Y01	1	0
8	A	2013	LPE	3	0
8	A	2009	LPE	2	0
7	A	2006	9Z9	2	0
8	B	304	LPE	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.












































5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-35975. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 120




Y Index: 132




Z Index: 133


6.3.2 Raw map



X Index: 124



Y Index: 132




Z Index: 133


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z

6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.55. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 66 nm³; this corresponds to an approximate mass of 60 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.333 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.333 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.00	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.45	4.01	3.52

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.45 differs from the reported value 3.0 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-35975 and PDB model 8J4F. Per-residue inclusion information can be found in section 3 on page 11.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.55 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.55).

9.4 Atom inclusion [i](#)



At the recommended contour level, 78% of all backbone atoms, 72% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.55) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7160	0.5010
A	0.7050	0.4940
B	0.8030	0.5590
C	0.6430	0.3270
D	0.5000	0.3440
E	0.9290	0.5840

