



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:38 AM UTC

PDB ID : 2J5V / pdb_00002j5v
Title : GLUTAMATE 5-KINASE FROM ESCHERICHIA COLI COMPLEXED
WITH GLUTAMYL-5-PHOSPHATE AND PYROGLUTAMIC ACID
Authors : Marco-Marin, C.; Gil-Ortiz, F.; Perez-Arellano, I.; Cervera, J.; Fita, I.; Rubio,
V.
Deposited on : 2006-09-19
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	367	
1	B	367	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PCA	A	1369	-	-	X	-
3	PCA	B	1368	-	-	X	-
4	RGP	A	1370	X	-	-	-
4	RGP	B	1369	X	-	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5087 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLUTAMATE 5-KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	323	Total	C	N	O	S	0	0	0
			2429	1512	450	459	8			
1	B	325	Total	C	N	O	S	0	0	0
			2437	1516	452	461	8			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	129	VAL	ILE	engineered mutation	UNP P0A7B5
B	129	VAL	ILE	engineered mutation	UNP P0A7B5

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PYROGLUTAMIC ACID (CCD ID: PCA) (formula: C₅H₇NO₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			9	5	1	3		
3	B	1	Total	C	N	O	0	0
			9	5	1	3		

- Molecule 4 is GAMMA-GLUTAMYL PHOSPHATE (CCD ID: RGP) (formula: $C_5H_{10}NO_7P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			14	5	1	7	1		
4	B	1	Total	C	N	O	P	0	0
			14	5	1	7	1		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	89	Total	O	0	0
			89	89		
6	B	75	Total	O	0	0
			75	75		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GLUTAMATE 5-KINASE



• Molecule 1: GLUTAMATE 5-KINASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	101.45Å 101.45Å 178.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.39 – 2.50 88.23 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.5 (88.39-2.50) 99.5 (88.23-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.06 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.192 , 0.244 (Not available) , 0.244	Depositor DCC
R_{free} test set	1667 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	37.7	Xtriage
Anisotropy	0.145	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 29.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5087	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PCA, SO4, RGP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/2460	0.89	1/3332 (0.0%)
1	B	0.71	0/2468	0.92	0/3342
All	All	0.70	0/4928	0.90	1/6674 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	251	VAL	N-CA-C	-5.63	100.91	108.35

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	366	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2429	0	2473	35	0
1	B	2437	0	2479	29	0
2	A	1	0	0	0	0
3	A	9	0	5	10	0
3	B	9	0	5	5	0
4	A	14	0	7	2	0
4	B	14	0	7	0	0
5	A	5	0	0	1	0
5	B	5	0	0	0	0
6	A	89	0	0	3	0
6	B	75	0	0	0	0
All	All	5087	0	4976	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:53:ILE:H	3:B:1368:PCA:HB3	1.14	1.08
1:A:53:ILE:H	3:A:1369:PCA:HB3	1.24	1.02
1:B:53:ILE:N	3:B:1368:PCA:HB3	1.84	0.92
1:A:53:ILE:N	3:A:1369:PCA:HB3	1.92	0.84
1:A:53:ILE:H	3:A:1369:PCA:CB	1.91	0.84
5:A:1371:SO4:O4	6:A:2088:HOH:O	2.01	0.77
1:A:44:ARG:NH2	1:A:126:ASP:OD2	2.18	0.76
1:B:278:ILE:HD12	1:B:313:ILE:CD1	2.16	0.74
1:B:278:ILE:HD12	1:B:313:ILE:HD13	1.67	0.74
1:B:59:HIS:CG	1:B:82:ARG:HD2	2.24	0.72
1:B:44:ARG:NH2	1:B:126:ASP:OD2	2.24	0.71
1:B:53:ILE:H	3:B:1368:PCA:CB	2.01	0.69
1:A:52:ALA:N	3:A:1369:PCA:HB2	2.08	0.68
1:A:278:ILE:HG12	1:A:313:ILE:HD12	1.75	0.67
1:A:53:ILE:HB	3:A:1369:PCA:HB3	1.77	0.66
1:B:280:VAL:HG13	1:B:284:ALA:HB3	1.80	0.63
1:A:238:LYS:HD3	1:A:251:VAL:HG13	1.80	0.61
1:B:52:ALA:O	1:B:79:GLY:HA3	2.00	0.60
1:A:121:LEU:HD13	1:A:131:PRO:HG2	1.84	0.60
1:B:278:ILE:CD1	1:B:313:ILE:HD13	2.33	0.59
1:A:69:ILE:O	1:A:73:GLN:HG3	2.03	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:269:ILE:HB	1:B:365:ILE:HD12	1.84	0.57
1:A:59:HIS:CG	1:A:82:ARG:HD2	2.40	0.57
1:A:103:LEU:HD11	1:A:152:LEU:HD11	1.87	0.56
1:A:22:ARG:NH1	6:A:2003:HOH:O	2.41	0.54
1:B:30:GLU:OE2	1:B:33:ARG:NH1	2.41	0.54
1:A:241:VAL:HG23	1:A:251:VAL:HG11	1.90	0.53
1:B:121:LEU:HD13	1:B:131:PRO:HG2	1.90	0.53
1:B:103:LEU:HD11	1:B:152:LEU:HD11	1.91	0.52
1:A:53:ILE:CB	3:A:1369:PCA:HB3	2.39	0.52
1:A:52:ALA:O	1:A:79:GLY:HA3	2.11	0.51
1:A:238:LYS:HD3	1:A:251:VAL:CG1	2.41	0.51
1:A:10:LYS:NZ	4:A:1370:RGP:OA2	2.29	0.50
1:B:52:ALA:N	3:B:1368:PCA:HB2	2.28	0.49
1:A:30:GLU:OE2	1:A:33:ARG:NH1	2.47	0.48
1:B:103:LEU:CD1	1:B:152:LEU:HD11	2.44	0.48
1:A:102:LEU:HD12	1:B:102:LEU:HD12	1.96	0.48
1:B:118:ARG:O	1:B:122:ARG:HG2	2.14	0.48
1:A:262:LEU:HD12	6:A:2039:HOH:O	2.14	0.47
1:A:10:LYS:C	1:A:10:LYS:HD3	2.41	0.46
1:B:238:LYS:HD3	1:B:251:VAL:CG2	2.46	0.45
1:A:85:GLN:HG3	1:B:85:GLN:CD	2.42	0.45
1:A:280:VAL:HG13	1:A:284:ALA:HB3	1.99	0.45
1:A:366:THR:O	1:A:367:ARG:HB2	2.16	0.45
1:A:149:ASN:HA	1:A:152:LEU:HB3	1.99	0.44
1:B:53:ILE:HB	3:B:1368:PCA:HB3	1.98	0.44
1:B:317:ASN:OD1	1:B:319:GLU:HB3	2.18	0.44
1:B:118:ARG:HG3	1:B:159:LEU:O	2.18	0.44
1:A:53:ILE:CA	3:A:1369:PCA:HB3	2.48	0.43
1:A:85:GLN:HG3	1:B:85:GLN:CG	2.48	0.43
1:B:169:THR:HG22	1:B:217:LYS:HE2	2.00	0.43
1:A:319:GLU:O	1:A:319:GLU:HG3	2.19	0.43
1:A:51:GLY:C	3:A:1369:PCA:HB2	2.43	0.43
1:A:281:ASP:HB3	1:A:284:ALA:H	1.83	0.42
1:B:111:ARG:HD3	1:B:268:TRP:CD1	2.54	0.42
1:B:111:ARG:HG3	1:B:115:LEU:HD22	2.02	0.41
1:B:38:LEU:HD11	1:B:242:ILE:HG22	2.03	0.41
1:A:139:VAL:HG11	1:B:103:LEU:HB3	2.02	0.41
3:A:1369:PCA:O	4:A:1370:RGP:HG1	2.21	0.41
1:A:53:ILE:O	1:A:57:ARG:HG3	2.21	0.41
1:A:53:ILE:N	3:A:1369:PCA:CB	2.65	0.40
1:A:344:GLN:CD	1:A:344:GLN:H	2.29	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/367 (87%)	311 (98%)	7 (2%)	1 (0%)	36	55
1	B	321/367 (88%)	312 (97%)	9 (3%)	0	100	100
All	All	640/734 (87%)	623 (97%)	16 (2%)	1 (0%)	43	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	281	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/285 (89%)	241 (95%)	12 (5%)	23	47
1	B	253/285 (89%)	235 (93%)	18 (7%)	13	29
All	All	506/570 (89%)	476 (94%)	30 (6%)	18	37

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	GLU
1	A	74	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	83	LEU
1	A	110	ASP
1	A	121	LEU
1	A	135	GLU
1	A	260	THR
1	A	277	GLU
1	A	280	VAL
1	A	282	GLU
1	A	298	LYS
1	A	353	GLU
1	B	25	ARG
1	B	69	ILE
1	B	74	LEU
1	B	88	GLU
1	B	92	SER
1	B	103	LEU
1	B	111	ARG
1	B	115	LEU
1	B	121	LEU
1	B	135	GLU
1	B	247	GLU
1	B	260	THR
1	B	280	VAL
1	B	308	SER
1	B	309	ARG
1	B	313	ILE
1	B	344	GLN
1	B	347	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	59	HIS
1	B	151	ASN
1	B	258	GLN
1	B	264	ASN
1	B	306	ASN
1	B	344	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	RGP	A	1370	-	11,13,13	0.83	0	13,18,18	1.39	1 (7%)
5	SO4	B	1370	-	4,4,4	0.33	0	6,6,6	0.35	0
3	PCA	A	1369	-	9,9,9	0.93	0	12,12,12	2.43	8 (66%)
3	PCA	B	1368	-	9,9,9	1.36	2 (22%)	12,12,12	2.09	5 (41%)
5	SO4	A	1371	2	4,4,4	0.28	0	6,6,6	0.35	0
4	RGP	B	1369	-	11,13,13	0.80	0	13,18,18	1.10	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	RGP	A	1370	-	1/1/4/4	5/12/14/14	-
3	PCA	B	1368	-	-	4/4/13/13	0/1/1/1
3	PCA	A	1369	-	-	0/4/13/13	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	RGP	B	1369	-	1/1/4/4	5/12/14/14	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1368	PCA	CA-C	2.73	1.59	1.52
3	B	1368	PCA	O-C	2.04	1.28	1.22

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1369	PCA	CA-N-CD	-4.90	110.33	114.48
3	B	1368	PCA	CG-CB-CA	4.62	109.89	104.49
3	B	1368	PCA	CA-N-CD	-3.70	111.35	114.48
3	A	1369	PCA	CG-CB-CA	3.18	108.21	104.49
3	A	1369	PCA	OE-CD-CG	-2.73	121.84	126.72
3	A	1369	PCA	O-C-CA	-2.47	114.29	122.26
3	A	1369	PCA	CB-CA-N	2.41	104.62	102.70
4	B	1369	RGP	OA3-PA-OA2	2.28	116.37	107.80
4	A	1370	RGP	OE2-CD-CG	2.28	121.88	111.87
3	B	1368	PCA	CG-CD-N	2.23	113.85	108.39
3	B	1368	PCA	OE-CD-CG	-2.20	122.79	126.72
4	B	1369	RGP	OE2-CD-CG	2.15	121.30	111.87
3	A	1369	PCA	C-CA-N	-2.14	109.31	111.94
3	A	1369	PCA	OXT-C-CA	2.08	120.54	113.51
3	B	1368	PCA	CB-CG-CD	-2.07	101.21	104.41
3	A	1369	PCA	CG-CD-N	2.02	113.33	108.39

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	1370	RGP	CA
4	B	1369	RGP	CA

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1368	PCA	O-C-CA-N
3	B	1368	PCA	OXT-C-CA-N
3	B	1368	PCA	OXT-C-CA-CB
4	A	1370	RGP	N-CA-CB-CG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	1369	RGP	OXT-C-CA-N
4	A	1370	RGP	O-C-CA-N
4	B	1369	RGP	O-C-CA-N
3	B	1368	PCA	O-C-CA-CB
4	B	1369	RGP	CA-CB-CG-CD
4	A	1370	RGP	OXT-C-CA-N
4	A	1370	RGP	C-CA-CB-CG
4	B	1369	RGP	OXT-C-CA-CB
4	B	1369	RGP	O-C-CA-CB
4	A	1370	RGP	OE2-CD-CG-CB

There are no ring outliers.

4 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1370	RGP	2	0
3	A	1369	PCA	10	0
3	B	1368	PCA	5	0
5	A	1371	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/367 (88%)	0.28	12 (3%) 45 40	31, 36, 44, 55	0
1	B	325/367 (88%)	0.41	18 (5%) 30 27	28, 36, 45, 55	0
All	All	648/734 (88%)	0.35	30 (4%) 37 33	28, 36, 44, 55	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	214	MET	4.5
1	B	278	ILE	3.6
1	A	260	THR	3.4
1	B	319	GLU	3.4
1	B	342	HIS	2.9
1	B	281	ASP	2.9
1	B	171	GLN	2.7
1	B	212	GLY	2.7
1	A	290	GLU	2.7
1	A	259	ALA	2.6
1	A	342	HIS	2.6
1	B	21	ARG	2.5
1	B	64	GLU	2.4
1	B	291	ARG	2.4
1	B	343	SER	2.3
1	B	344	GLN	2.3
1	B	346	ILE	2.3
1	A	361	ARG	2.3
1	B	340	GLY	2.2
1	A	344	GLN	2.2
1	B	213	GLY	2.2
1	A	347	ASP	2.2
1	A	281	ASP	2.2
1	A	64	GLU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	282	GLU	2.2
1	A	110	ASP	2.1
1	B	298	LYS	2.1
1	A	351	GLY	2.1
1	B	321	ARG	2.0
1	B	347	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PCA	B	1368	9/9	0.62	0.24	35,39,41,43	9
3	PCA	A	1369	9/9	0.70	0.23	32,36,39,41	9
4	RGP	B	1369	14/14	0.72	0.30	25,31,38,38	14
5	SO4	B	1370	5/5	0.77	0.21	57,57,58,59	5
5	SO4	A	1371	5/5	0.84	0.16	52,53,54,54	5
4	RGP	A	1370	14/14	0.85	0.16	45,49,54,54	4
2	MG	A	1368	1/1	0.93	0.06	19,19,19,19	1

6.5 Other polymers [i](#)

There are no such residues in this entry.