



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 02:48 PM EDT

PDB ID : 4J5P / pdb_00004j5p
Title : Crystal Structure of a Covalently Bound alpha-Ketoheterocycle Inhibitor (Ph
enhexyl/Oxadiazole/Pyridine) to a Humanized Variant of Fatty Acid Amide
Hydrolase
Authors : Otrubova, K.; Brown, M.; McCormick, M.S.; Han, G.W.; O'Neal, S.T.; Cra-
vatt, B.F.; Stevens, R.C.; Lichtman, A.H.; Boger, D.L.
Deposited on : 2013-02-08
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	573	3% 86% 9% 5%
1	B	573	3% 87% 7% 5%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8787 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fatty-acid amide hydrolase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	544	Total	C	N	O	S	0	0	0
			4198	2679	717	772	30			
1	B	545	Total	C	N	O	S	0	0	0
			4207	2684	719	774	30			

There are 58 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	7	MET	-	expression tag	UNP P97612
A	8	GLY	-	expression tag	UNP P97612
A	9	SER	-	expression tag	UNP P97612
A	10	SER	-	expression tag	UNP P97612
A	11	HIS	-	expression tag	UNP P97612
A	12	HIS	-	expression tag	UNP P97612
A	13	HIS	-	expression tag	UNP P97612
A	14	HIS	-	expression tag	UNP P97612
A	15	HIS	-	expression tag	UNP P97612
A	16	HIS	-	expression tag	UNP P97612
A	17	SER	-	expression tag	UNP P97612
A	18	SER	-	expression tag	UNP P97612
A	19	GLY	-	expression tag	UNP P97612
A	20	LEU	-	expression tag	UNP P97612
A	21	VAL	-	expression tag	UNP P97612
A	22	PRO	-	expression tag	UNP P97612
A	23	ARG	-	expression tag	UNP P97612
A	24	GLY	-	expression tag	UNP P97612
A	25	SER	-	expression tag	UNP P97612
A	26	HIS	-	expression tag	UNP P97612
A	27	MET	-	expression tag	UNP P97612
A	28	ALA	-	expression tag	UNP P97612
A	29	SER	-	expression tag	UNP P97612
A	192	PHE	LEU	engineered mutation	UNP P97612
A	194	TYR	PHE	engineered mutation	UNP P97612

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	377	THR	ALA	engineered mutation	UNP P97612
A	435	ASN	SER	engineered mutation	UNP P97612
A	491	VAL	ILE	engineered mutation	UNP P97612
A	495	MET	VAL	engineered mutation	UNP P97612
B	7	MET	-	expression tag	UNP P97612
B	8	GLY	-	expression tag	UNP P97612
B	9	SER	-	expression tag	UNP P97612
B	10	SER	-	expression tag	UNP P97612
B	11	HIS	-	expression tag	UNP P97612
B	12	HIS	-	expression tag	UNP P97612
B	13	HIS	-	expression tag	UNP P97612
B	14	HIS	-	expression tag	UNP P97612
B	15	HIS	-	expression tag	UNP P97612
B	16	HIS	-	expression tag	UNP P97612
B	17	SER	-	expression tag	UNP P97612
B	18	SER	-	expression tag	UNP P97612
B	19	GLY	-	expression tag	UNP P97612
B	20	LEU	-	expression tag	UNP P97612
B	21	VAL	-	expression tag	UNP P97612
B	22	PRO	-	expression tag	UNP P97612
B	23	ARG	-	expression tag	UNP P97612
B	24	GLY	-	expression tag	UNP P97612
B	25	SER	-	expression tag	UNP P97612
B	26	HIS	-	expression tag	UNP P97612
B	27	MET	-	expression tag	UNP P97612
B	28	ALA	-	expression tag	UNP P97612
B	29	SER	-	expression tag	UNP P97612
B	192	PHE	LEU	engineered mutation	UNP P97612
B	194	TYR	PHE	engineered mutation	UNP P97612
B	377	THR	ALA	engineered mutation	UNP P97612
B	435	ASN	SER	engineered mutation	UNP P97612
B	491	VAL	ILE	engineered mutation	UNP P97612
B	495	MET	VAL	engineered mutation	UNP P97612

- Molecule 2 is (1S)-1-{5-[5-(bromomethyl)pyridin-2-yl]-1,3-oxazol-2-yl}-7-phenylheptan-1-ol (CCD ID: BR1) (formula: C₂₂H₂₅BrN₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			26	22	2	2		
2	B	1	Total	C	N	O	0	0
			26	22	2	2		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	124	Total	O	0	0
			124	124		
5	B	170	Total	O	0	0
			170	170		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Fatty-acid amide hydrolase 1



• Molecule 1: Fatty-acid amide hydrolase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.07 Å 104.07 Å 261.03 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.38 – 2.30 33.38 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.2 (33.38-2.30) 96.7 (33.38-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	48.72 (at 2.29 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.220 , 0.265 0.221 , 0.259	Depositor DCC
R_{free} test set	3602 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	55.3	Xtriage
Anisotropy	0.295	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 42.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8787	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CL, BR1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.05	4/4292 (0.1%)	1.09	8/5822 (0.1%)
1	B	1.10	2/4301 (0.0%)	1.12	9/5834 (0.2%)
All	All	1.08	6/8593 (0.1%)	1.11	17/11656 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	368	ILE	CA-CB	-7.81	1.48	1.53
1	B	167	ASP	C-O	-7.38	1.14	1.23
1	B	543	VAL	C-O	-5.78	1.18	1.24
1	A	139	VAL	C-O	-5.19	1.19	1.24
1	A	34	ARG	N-CA	5.13	1.51	1.46
1	A	376	SER	N-CA	5.06	1.52	1.46

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	257	THR	N-CA-C	-6.36	101.53	110.50
1	B	575	GLN	N-CA-C	6.35	118.20	111.28
1	B	257	THR	N-CA-C	-6.07	101.42	110.23
1	B	124	GLN	N-CA-C	5.88	117.69	111.28
1	A	275	ALA	N-CA-C	5.83	118.11	111.11
1	B	258	GLY	N-CA-C	5.73	119.82	112.83
1	A	258	GLY	N-CA-C	5.63	119.70	112.83
1	A	432	PHE	N-CA-C	-5.58	105.20	111.28
1	B	121	CYS	N-CA-C	5.53	119.86	113.18
1	B	49	ARG	CB-CA-C	-5.52	101.98	110.81
1	A	121	CYS	N-CA-C	5.34	119.64	113.18
1	B	49	ARG	CG-CD-NE	-5.31	100.31	112.00
1	A	281	GLY	CA-C-N	5.17	125.38	120.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	281	GLY	C-N-CA	5.17	125.38	120.31
1	A	315	ARG	NE-CZ-NH2	5.07	123.77	119.20
1	B	255	LYS	CB-CA-C	-5.05	105.90	110.44
1	B	432	PHE	N-CA-C	-5.02	105.81	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4198	0	4252	13	0
1	B	4207	0	4260	20	0
2	A	26	0	22	0	0
2	B	26	0	22	2	0
3	A	14	0	20	0	0
3	B	21	0	30	0	0
4	B	1	0	0	0	0
5	A	124	0	0	1	0
5	B	170	0	0	8	0
All	All	8787	0	8606	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:304:THR:O	1:B:463:LYS:HE3	1.77	0.84
1:B:34:ARG:HA	5:B:704:HOH:O	1.92	0.70
1:A:80:LEU:HD13	1:A:288:GLU:HG3	1.84	0.59
1:A:237:ASP:HA	1:A:241:SER:HB2	1.85	0.58
1:B:80:LEU:HD13	1:B:288:GLU:HG3	1.86	0.56
1:B:352:ARG:HD2	5:B:710:HOH:O	2.05	0.56
1:B:237:ASP:HA	1:B:241:SER:HB2	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:121:CYS:HA	1:B:124:GLN:HB2	1.89	0.54
1:B:168:CYS:SG	1:B:171:VAL:HG23	2.48	0.54
1:B:105:TRP:NE1	5:B:715:HOH:O	2.40	0.54
1:B:45:ARG:NH2	5:B:763:HOH:O	2.41	0.52
1:A:535:LEU:HG	1:A:539:MET:HE2	1.91	0.52
1:B:575:GLN:H	1:B:575:GLN:CD	2.16	0.51
1:B:61:ARG:N	5:B:719:HOH:O	2.42	0.51
1:B:535:LEU:HG	1:B:539:MET:HE2	1.93	0.51
1:A:168:CYS:SG	1:A:171:VAL:HG23	2.53	0.48
1:B:191:MET:HE3	2:B:601:BR1:C22	2.43	0.48
1:B:209:LYS:NZ	5:B:822:HOH:O	2.45	0.47
1:B:508:PRO:HB3	5:B:742:HOH:O	2.16	0.46
1:B:87:GLN:NE2	5:B:714:HOH:O	2.50	0.44
1:B:269:CYS:N	2:B:601:BR1:C21	2.81	0.44
1:A:448:GLN:HG2	5:A:701:HOH:O	2.18	0.43
1:B:97:PHE:HB2	1:B:125:LEU:HD21	2.00	0.43
1:A:185:THR:HB	1:A:220:GLY:HA3	2.01	0.43
1:B:190:SER:O	1:B:191:MET:HB2	2.20	0.42
1:A:97:PHE:HB2	1:A:125:LEU:HD21	2.01	0.41
1:A:346:LEU:C	1:A:346:LEU:HD13	2.46	0.41
1:A:416:ARG:HG3	1:A:437:ARG:CZ	2.50	0.41
1:A:190:SER:O	1:A:191:MET:HB2	2.21	0.41
1:A:142:LYS:HE2	1:A:191:MET:SD	2.61	0.41
1:A:551:CYS:HB3	1:A:564:PHE:CD1	2.56	0.41
1:B:153:THR:O	1:B:154:LEU:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	542/573 (95%)	523 (96%)	18 (3%)	1 (0%)	43 55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	543/573 (95%)	524 (96%)	18 (3%)	1 (0%)	43	55
All	All	1085/1146 (95%)	1047 (96%)	36 (3%)	2 (0%)	43	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	193	SER
1	A	193	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	462/487 (95%)	440 (95%)	22 (5%)	23	35
1	B	463/487 (95%)	444 (96%)	19 (4%)	27	41
All	All	925/974 (95%)	884 (96%)	41 (4%)	25	38

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	LYS
1	A	57	LYS
1	A	60	GLN
1	A	61	ARG
1	A	80	LEU
1	A	84	GLN
1	A	119	THR
1	A	133	LEU
1	A	280	LEU
1	A	294	LEU
1	A	295	LYS
1	A	302	LEU
1	A	316	GLU
1	A	317	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	352	ARG
1	A	386	ARG
1	A	416	ARG
1	A	418	LEU
1	A	420	LEU
1	A	446	LYS
1	A	537	LYS
1	A	576	LYS
1	B	46	GLN
1	B	61	ARG
1	B	65	GLN
1	B	80	LEU
1	B	84	GLN
1	B	119	THR
1	B	120	ASP
1	B	133	LEU
1	B	280	LEU
1	B	294	LEU
1	B	302	LEU
1	B	315	ARG
1	B	352	ARG
1	B	416	ARG
1	B	418	LEU
1	B	463	LYS
1	B	465	MET
1	B	575	GLN
1	B	576	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	GLN
1	A	124	GLN
1	A	203	GLN
1	A	334	ASN
1	A	351	GLN
1	A	366	ASN
1	A	367	ASN
1	A	434	ASN
1	A	448	GLN
1	A	449	HIS
1	A	557	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	570	GLN
1	B	65	GLN
1	B	124	GLN
1	B	334	ASN
1	B	351	GLN
1	B	366	ASN
1	B	367	ASN
1	B	390	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BR1	B	601	1	27,28,29	1.61	3 (11%)	30,36,37	2.13	7 (23%)
3	PEG	B	603	-	6,6,6	0.63	0	5,5,5	0.29	0
3	PEG	B	604	-	6,6,6	0.78	0	5,5,5	0.50	0
3	PEG	B	602	-	6,6,6	0.57	0	5,5,5	0.40	0
3	PEG	A	603	-	6,6,6	0.73	0	5,5,5	0.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BR1	A	601	1	27,28,29	1.43	5 (18%)	30,36,37	2.10	10 (33%)
3	PEG	A	602	-	6,6,6	0.44	0	5,5,5	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BR1	B	601	1	-	7/17/17/19	0/3/3/3
3	PEG	B	603	-	-	2/4/4/4	-
3	PEG	B	604	-	-	2/4/4/4	-
3	PEG	B	602	-	-	2/4/4/4	-
3	PEG	A	603	-	-	3/4/4/4	-
2	BR1	A	601	1	-	7/17/17/19	0/3/3/3
3	PEG	A	602	-	-	2/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	BR1	C14-N1	5.23	1.37	1.29
2	B	601	BR1	C7-C8	-3.18	1.42	1.51
2	A	601	BR1	C14-N1	2.91	1.34	1.29
2	B	601	BR1	C15-C16	2.90	1.39	1.34
2	A	601	BR1	C7-C8	-2.77	1.43	1.51
2	A	601	BR1	C2-C1	2.72	1.57	1.53
2	A	601	BR1	O2-C16	-2.35	1.34	1.38
2	A	601	BR1	C15-C16	2.12	1.38	1.34

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	BR1	O2-C16-C15	5.27	112.59	106.93
2	B	601	BR1	C18-C19-C20	-4.78	115.86	121.37
2	A	601	BR1	C16-C15-N1	-4.27	105.84	110.07
2	A	601	BR1	O2-C16-C15	4.24	111.49	106.93
2	B	601	BR1	C16-C15-N1	-4.23	105.87	110.07
2	A	601	BR1	C18-C19-C20	-4.15	116.59	121.37
2	B	601	BR1	O2-C16-C17	3.72	124.11	116.98
2	A	601	BR1	O2-C16-C17	3.43	123.55	116.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	BR1	C1-C14-N1	-3.39	124.71	128.78
2	A	601	BR1	C15-N1-C14	3.31	107.00	104.33
2	B	601	BR1	C5-C6-C7	-3.19	100.40	113.74
2	A	601	BR1	C21-C20-C22	-3.01	115.33	121.09
2	A	601	BR1	O2-C14-N1	-2.51	111.52	114.10
2	B	601	BR1	C15-N1-C14	2.31	106.20	104.33
2	A	601	BR1	C1-C14-N1	-2.29	126.03	128.78
2	A	601	BR1	C5-C6-C7	-2.29	104.19	113.74
2	A	601	BR1	C19-C20-C22	2.10	121.24	116.60

There are no chirality outliers.

All (25) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	BR1	C2-C1-C14-O2
2	A	601	BR1	C2-C1-C14-N1
2	B	601	BR1	C2-C1-C14-O2
2	B	601	BR1	O1-C1-C14-N1
2	B	601	BR1	C2-C1-C14-N1
3	B	604	PEG	O2-C3-C4-O4
3	A	603	PEG	O2-C3-C4-O4
3	B	602	PEG	O2-C3-C4-O4
3	A	602	PEG	O2-C3-C4-O4
3	A	603	PEG	C4-C3-O2-C2
3	B	603	PEG	C1-C2-O2-C3
3	A	602	PEG	O1-C1-C2-O2
3	B	602	PEG	O1-C1-C2-O2
2	A	601	BR1	O1-C1-C14-O2
2	B	601	BR1	O1-C1-C14-O2
2	A	601	BR1	C2-C3-C4-C5
3	B	604	PEG	C4-C3-O2-C2
2	A	601	BR1	O1-C1-C14-N1
3	A	603	PEG	O1-C1-C2-O2
2	B	601	BR1	C4-C5-C6-C7
2	B	601	BR1	C6-C7-C8-C9
2	B	601	BR1	C2-C3-C4-C5
2	A	601	BR1	C6-C7-C8-C13
2	A	601	BR1	C6-C7-C8-C9
3	B	603	PEG	C4-C3-O2-C2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	BR1	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	544/573 (94%)	0.20	15 (2%) 55 57	37, 57, 88, 136	0
1	B	545/573 (95%)	0.26	20 (3%) 45 48	39, 57, 98, 117	0
All	All	1089/1146 (95%)	0.23	35 (3%) 50 52	37, 57, 95, 136	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	33	GLY	4.0
1	A	576	LYS	3.4
1	B	69	LEU	3.3
1	B	33	GLY	3.1
1	B	123	THR	3.1
1	B	76	THR	3.0
1	B	125	LEU	2.9
1	B	575	GLN	2.8
1	B	72	GLU	2.7
1	B	73	ALA	2.7
1	A	570	GLN	2.7
1	B	404	LEU	2.6
1	B	208	TRP	2.6
1	A	574	PRO	2.6
1	B	36	LYS	2.5
1	B	432	PHE	2.5
1	A	404	LEU	2.5
1	A	226	GLY	2.5
1	B	67	PRO	2.5
1	B	119	THR	2.3
1	A	64	LEU	2.3
1	A	357	GLY	2.3
1	A	575	GLN	2.2
1	B	524	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	124	GLN	2.2
1	A	323	ARG	2.2
1	B	576	LYS	2.1
1	A	69	LEU	2.1
1	B	64	LEU	2.1
1	A	354	GLU	2.1
1	B	417	LEU	2.1
1	A	568	VAL	2.1
1	A	119	THR	2.0
1	A	389	LEU	2.0
1	B	121	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PEG	A	603	7/7	0.84	0.13	54,59,75,76	0
2	BR1	B	601	26/27	0.87	0.17	49,62,68,71	0
3	PEG	B	602	7/7	0.87	0.14	68,76,81,82	0
3	PEG	B	603	7/7	0.87	0.14	82,93,106,107	0
3	PEG	B	604	7/7	0.88	0.12	62,69,78,84	0
2	BR1	A	601	26/27	0.90	0.17	48,68,77,81	0
3	PEG	A	602	7/7	0.91	0.11	69,69,74,74	0
4	CL	B	605	1/1	0.97	0.06	48,48,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around BR1 B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BR1 A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.