



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:06 PM UTC

PDB ID : 8J5W / pdb_00008j5w
Title : The crystal structure of TrkA(F589L) kinase in complex with N-(3-cyclopropyl-5-((4-methylpiperazin-1-yl)methyl)phenyl)-4⁶-methyl-14-oxo-5-oxa-13-aza-1(3,6)-imidazo[1,2-b]pyridazina-4(1,3)-benzenacyclotetradecaphan-2-yne-4⁵-carboxamide
Authors : Zhang, Z.M.; Wang, Y.J.
Deposited on : 2023-04-24
Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	7% 80% 14% • 5%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2417 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called High affinity nerve growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	63	0	0
			2368	1513	427	412	16			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	589	LEU	PHE	engineered mutation	UNP P04629

- Molecule 2 is N-(3-cyclopropyl-5-((4-methylpiperazin-1-yl)methyl)phenyl)-4⁶-methyl-14-oxo-5-oxa-13-aza-1(3,6)-imidazo[1,2-b]pyridazina-4(1,3)-benzenacyclotetradecaphan-2-yne-4⁵-carboxamide (CCD ID: A4U) (formula: C₃₉H₄₅N₇O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			49	39	7	3		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: High affinity nerve growth factor receptor



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 1 2	Depositor
Cell constants a, b, c, α , β , γ	51.68Å 51.68Å 229.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.05 – 2.28 29.05 – 2.28	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.05-2.28) 99.9 (29.05-2.28)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.80 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.9_1692+SVN	Depositor
R, R_{free}	0.217 , 0.266 0.228 , 0.270	Depositor DCC
R_{free} test set	1650 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.243	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.070 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2417	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A4U

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/2424	0.77	4/3279 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	676	TYR	N-CA-C	-6.55	103.89	112.41
1	A	767	GLU	CA-C-N	5.31	124.76	119.24
1	A	767	GLU	C-N-CA	5.31	124.76	119.24
1	A	501	CYS	N-CA-C	-5.26	106.12	112.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2368	0	2353	24	0
2	A	49	0	0	1	0
All	All	2417	0	2353	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:669:PHE:HB3	1:A:671:MET:H	1.57	0.69
1:A:668:ASP:N	2:A:801:A4U:O27	2.21	0.68
1:A:488:GLY:O	1:A:489:HIS:ND1	2.29	0.66
1:A:606:PRO:HA	1:A:607:ASP:C	2.19	0.65
1:A:499:ASP:N	1:A:501:CYS:H	2.03	0.56
1:A:488:GLY:HA2	1:A:647:VAL:H	1.72	0.54
1:A:669:PHE:HB3	1:A:671:MET:N	2.24	0.53
1:A:748:ARG:NH1	1:A:752:CYS:O	2.46	0.46
1:A:669:PHE:HA	1:A:670:GLY:HA3	1.77	0.46
1:A:780:ARG:O	1:A:784:LEU:HG	2.16	0.45
1:A:669:PHE:HB3	1:A:672:SER:H	1.79	0.45
1:A:717:LEU:HG	1:A:760:MET:HE2	1.98	0.45
1:A:610:LEU:HD11	1:A:620:GLY:H	1.83	0.44
1:A:499:ASP:N	1:A:500:ALA:HA	2.32	0.43
1:A:513:LYS:HD2	1:A:528:GLU:HG3	2.01	0.43
1:A:692:ARG:NH1	1:A:728:TRP:O	2.52	0.42
1:A:755:GLU:HG3	1:A:795:LEU:HG	2.01	0.42
1:A:564:LEU:O	1:A:568:GLN:NE2	2.34	0.42
1:A:563:LEU:HD23	1:A:563:LEU:HA	1.88	0.41
1:A:553:ALA:O	1:A:557:PHE:N	2.45	0.41
1:A:669:PHE:HB2	1:A:672:SER:OG	2.21	0.41
1:A:617:VAL:HG22	1:A:662:LEU:HD21	2.03	0.40
1:A:753:PRO:HA	1:A:754:PRO:HD3	1.99	0.40
1:A:741:THR:C	1:A:766:ARG:HH12	2.29	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	290/313 (93%)	280 (97%)	9 (3%)	1 (0%)	36	44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	675	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/262 (94%)	237 (96%)	10 (4%)	28	40

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	486	LEU
1	A	535	GLU
1	A	548	GLU
1	A	554	ARG
1	A	607	ASP
1	A	671	MET
1	A	686	ARG
1	A	710	VAL
1	A	725	LYS
1	A	795	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	530	HIS
1	A	571	HIS
1	A	604	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	A4U	A	801	-	54,55,55	4.89	30 (55%)	70,77,77	2.36	14 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A4U	A	801	-	-	11/35/49/49	0/5/7/7

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	A4U	C13-N14	-10.69	1.27	1.47
2	A	801	A4U	C07-C02	10.22	1.53	1.40
2	A	801	A4U	C26-C22	9.92	1.53	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	A4U	C21-C22	9.69	1.53	1.39
2	A	801	A4U	C06-C07	8.92	1.53	1.39
2	A	801	A4U	C04-C03	8.71	1.53	1.39
2	A	801	A4U	C21-C12	8.60	1.53	1.39
2	A	801	A4U	C26-C10	8.54	1.53	1.39
2	A	801	A4U	C06-C05	8.47	1.53	1.39
2	A	801	A4U	C04-C05	8.28	1.52	1.39
2	A	801	A4U	C11-C12	8.01	1.52	1.39
2	A	801	A4U	C11-C10	7.59	1.51	1.39
2	A	801	A4U	C37-N36	7.49	1.46	1.33
2	A	801	A4U	C42-C41	6.13	1.53	1.41
2	A	801	A4U	C47-C46	5.03	1.50	1.42
2	A	801	A4U	N40-N39	4.69	1.50	1.38
2	A	801	A4U	C03-C48	4.42	1.52	1.43
2	A	801	A4U	C43-C42	4.38	1.45	1.35
2	A	801	A4U	C41-N44	4.18	1.39	1.33
2	A	801	A4U	C45-C46	3.35	1.40	1.37
2	A	801	A4U	C38-N39	3.21	1.43	1.32
2	A	801	A4U	C43-C38	3.12	1.47	1.42
2	A	801	A4U	C19-N17	-3.07	1.39	1.46
2	A	801	A4U	C38-C37	2.94	1.54	1.48
2	A	801	A4U	C08-N09	2.75	1.43	1.35
2	A	801	A4U	C16-N17	-2.75	1.39	1.46
2	A	801	A4U	O28-C05	2.74	1.43	1.37
2	A	801	A4U	C13-C12	2.63	1.56	1.51
2	A	801	A4U	C15-N14	-2.56	1.40	1.46
2	A	801	A4U	C20-N14	-2.35	1.40	1.46

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	A4U	C45-C46-N40	13.42	108.36	103.61
2	A	801	A4U	C45-N44-C41	5.42	108.73	105.11
2	A	801	A4U	C46-C45-N44	-5.31	106.66	112.35
2	A	801	A4U	C41-N40-N39	-5.13	121.67	126.75
2	A	801	A4U	C25-C23-C22	-4.51	115.12	121.28
2	A	801	A4U	C20-N14-C15	3.95	117.35	108.84
2	A	801	A4U	C24-C23-C22	-3.42	116.61	121.28
2	A	801	A4U	C41-N40-C46	3.30	109.38	106.06
2	A	801	A4U	C02-C03-C48	-2.92	118.41	121.14
2	A	801	A4U	C38-N39-N40	2.88	120.36	116.12
2	A	801	A4U	C16-C15-N14	2.54	115.77	110.65

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	A4U	C19-C20-N14	2.19	115.06	110.65
2	A	801	A4U	C29-O28-C05	-2.05	112.59	117.93
2	A	801	A4U	C42-C41-N44	2.05	134.30	130.59

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	A4U	C46-C47-C48-C03
2	A	801	A4U	C06-C05-O28-C29
2	A	801	A4U	C04-C05-O28-C29
2	A	801	A4U	C29-C30-C31-C32
2	A	801	A4U	C30-C29-O28-C05
2	A	801	A4U	C31-C32-C33-C34
2	A	801	A4U	C02-C03-C48-C47
2	A	801	A4U	C04-C03-C48-C47
2	A	801	A4U	C12-C13-N14-C15
2	A	801	A4U	C30-C31-C32-C33
2	A	801	A4U	O49-C37-C38-C43

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	A4U	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/313 (95%)	0.28	21 (7%) 22 23	20, 50, 94, 116	20 (6%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	610	LEU	4.4
1	A	675	ILE	4.2
1	A	557	PHE	3.9
1	A	672	SER	3.7
1	A	487	GLN	3.6
1	A	501	CYS	3.4
1	A	667	GLY	3.3
1	A	500	ALA	3.2
1	A	486	LEU	3.1
1	A	521	PHE	2.9
1	A	611	LEU	2.9
1	A	613	GLY	2.9
1	A	534	PRO	2.9
1	A	669	PHE	2.6
1	A	607	ASP	2.6
1	A	670	GLY	2.2
1	A	551	GLU	2.2
1	A	612	ALA	2.2
1	A	686	ARG	2.2
1	A	552	SER	2.1
1	A	606	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	A4U	A	801	49/49	0.71	0.23	48,90,107,153	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.