



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:14 AM UTC

PDB ID : 1J6Z / pdb_00001j6z
Title : UNCOMPLEXED ACTIN
Authors : Otterbein, L.R.; Graceffa, P.; Dominguez, R.
Deposited on : 2001-05-15
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1025 (1.54-1.54)
Ramachandran outliers	187476	1007 (1.54-1.54)
Sidechain outliers	187428	1007 (1.54-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	375	 83% 14% . .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3350 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACTIN ALPHA 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	369	Total	C	N	O	S	0	3	0
			2883	1824	486	552	21			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	73	HIC	HIS	modified residue	UNP P68135

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	6	Total	Ca	0	0
			6	6		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is TETRAMETHYLRHODAMINE-5-MALEIMIDE (CCD ID: RHO) (formula: $C_{28}H_{25}N_3O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			30	24	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	404	Total	O	0	0
			404	404		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: ACTIN ALPHA 1

Chain A:  83% 14% ..



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	112.81Å 37.50Å 85.26Å 90.00° 108.26° 90.00°	Depositor
Resolution (Å)	82.00 – 1.54	Depositor
% Data completeness (in resolution range)	97.0 (82.00-1.54)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.179 , 0.223	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3350	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, RHO, HIC, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/2947	1.51	30/3992 (0.8%)

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	51	ASP	CA-CB-CG	18.97	131.57	112.60
1	A	147	ARG	CG-CD-NE	8.96	131.71	112.00
1	A	22	ALA	CA-C-N	-7.53	111.49	123.31
1	A	22	ALA	C-N-CA	-7.53	111.49	123.31
1	A	35	VAL	CA-C-N	-7.25	117.04	122.33
1	A	35	VAL	C-N-CA	-7.25	117.04	122.33
1	A	201	VAL	O-C-N	-7.14	113.64	122.57
1	A	297	ASN	CA-CB-CG	6.76	119.36	112.60
1	A	356	TRP	CA-C-O	-6.47	114.69	121.55
1	A	28	ARG	NH1-CZ-NH2	6.31	127.51	119.30
1	A	147	ARG	CA-CB-CG	6.27	126.64	114.10
1	A	125	GLU	CB-CG-CD	6.16	123.08	112.60
1	A	90	PHE	CA-CB-CG	6.15	119.95	113.80
1	A	62	ARG	NE-CZ-NH2	-6.03	113.77	119.20
1	A	309	ILE	CA-C-O	-5.97	114.75	120.95
1	A	173	HIS	CA-CB-CG	-5.95	107.85	113.80
1	A	184	ASP	CA-CB-CG	-5.68	106.92	112.60
1	A	332	PRO	O-C-N	-5.66	118.71	121.31
1	A	183	ARG	NE-CZ-NH2	-5.52	114.23	119.20
1	A	30	VAL	CA-C-O	-5.50	114.92	120.31
1	A	194	THR	CA-C-N	5.42	128.08	120.28
1	A	194	THR	C-N-CA	5.42	128.08	120.28
1	A	335	ARG	N-CA-C	5.25	118.79	112.38
1	A	162	ASN	CA-CB-CG	5.24	117.84	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	150	GLY	CA-C-N	-5.19	116.37	123.12
1	A	150	GLY	C-N-CA	-5.19	116.37	123.12
1	A	274	ILE	CA-C-O	-5.12	115.09	120.57
1	A	50	LYS	CA-C-N	5.10	128.06	120.31
1	A	50	LYS	C-N-CA	5.10	128.06	120.31
1	A	354	GLN	CA-C-O	-5.03	113.20	119.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2883	0	2854	28	0
2	A	6	0	0	0	0
3	A	27	0	12	0	0
4	A	30	0	22	0	0
5	A	404	0	0	9	0
All	All	3350	0	2888	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:PRO:HG3	5:A:1495:HOH:O	1.90	0.71
1:A:153:LEU:HD23	1:A:299:MET:HE3	1.75	0.69
1:A:324:THR:HG23	5:A:1690:HOH:O	1.95	0.65
1:A:372:ARG:NH2	5:A:1632:HOH:O	2.31	0.63
1:A:299:MET:HE2	1:A:304:THR:HB	1.81	0.63
1:A:275:HIS:HA	1:A:313:MET:HE1	1.85	0.58
1:A:201:VAL:HG22	5:A:1664:HOH:O	2.04	0.57
1:A:237:GLU:HG2	5:A:1628:HOH:O	2.05	0.56
1:A:223:PHE:O	1:A:227:MET:HG2	2.08	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:ALA:O	1:A:172:PRO:HD3	2.09	0.53
1:A:227:MET:HE3	1:A:227:MET:HA	1.91	0.51
1:A:176:MET:HE2	1:A:284:LYS:HD2	1.93	0.50
1:A:41:GLN:OE1	1:A:44:MET:HE3	2.12	0.50
1:A:299:MET:HE1	1:A:309:ILE:HD13	1.94	0.50
1:A:334:GLU:HG3	5:A:1698:HOH:O	2.11	0.49
1:A:191:LYS:HE2	5:A:1737:HOH:O	2.14	0.47
1:A:299:MET:HE2	1:A:299:MET:HB3	1.95	0.44
1:A:372:ARG:CZ	5:A:1632:HOH:O	2.64	0.44
1:A:169:TYR:CE2	1:A:171:LEU:HD23	2.54	0.43
1:A:104:LEU:HD23	1:A:104:LEU:C	2.43	0.43
1:A:191:LYS:HG2	5:A:1737:HOH:O	2.18	0.43
1:A:105:LEU:HD11	1:A:123:MET:HG3	2.01	0.43
1:A:8:LEU:HG	1:A:101:HIS:HB3	2.02	0.42
1:A:37:ARG:HH22	1:A:84:LYS:HE3	1.85	0.41
1:A:176:MET:CE	1:A:284:LYS:HD2	2.50	0.41
1:A:189:LEU:HA	1:A:192:ILE:HG12	2.02	0.41
1:A:169:TYR:HD1	1:A:372:ARG:HH22	1.67	0.41
1:A:332:PRO:HA	1:A:333:PRO:HD3	1.97	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	369/375 (98%)	362 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	313/317 (99%)	307 (98%)	6 (2%)	50 20

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	68	LYS
1	A	71	ILE
1	A	147	ARG
1	A	227	MET
1	A	234	SER
1	A	372	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
1	A	121	GLN
1	A	297	ASN
1	A	353	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	HIC	A	73	1	10,11,12	0.67	0	9,14,16	3.26	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	HIC	A	73	1	-	1/5/6/8	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	73	HIC	NE2-CE1-ND1	-8.53	109.40	112.66
1	A	73	HIC	CD2-NE2-CE1	-2.69	105.65	106.71
1	A	73	HIC	CE1-ND1-CG	2.61	109.98	105.80
1	A	73	HIC	CD2-CG-ND1	-2.14	104.78	109.43

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	73	HIC	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ADP	A	1380	2	28,29,29	1.10	2 (7%)	43,45,45	1.25	6 (13%)
4	RHO	A	1381	-	33,33,40	3.73	18 (54%)	41,49,60	2.11	15 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	1380	2	-	0/16/32/32	0/3/3/3
4	RHO	A	1381	-	-	1/14/41/58	0/4/4/5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1381	RHO	C8-C14	13.37	1.47	1.35
4	A	1381	RHO	C22-C17	5.54	1.48	1.39
4	A	1381	RHO	C13-C12	-5.42	1.39	1.49
4	A	1381	RHO	OH2-C23	5.23	1.37	1.22
4	A	1381	RHO	C-N	4.55	1.48	1.37
4	A	1381	RHO	C18-C17	4.48	1.48	1.40
4	A	1381	RHO	C4-C3	4.40	1.46	1.39
4	A	1381	RHO	C19-C20	4.30	1.46	1.39
4	A	1381	RHO	C11-N1	3.78	1.47	1.36
4	A	1381	RHO	C19-C18	3.71	1.45	1.39
4	A	1381	RHO	C12-C11	3.54	1.45	1.35
3	A	1380	ADP	C5-N7	-3.22	1.33	1.39
4	A	1381	RHO	C17-C8	-3.17	1.45	1.49
4	A	1381	RHO	C3-C2	3.06	1.46	1.40
3	A	1380	ADP	C2-N1	2.98	1.39	1.33
4	A	1381	RHO	OH3-C23	-2.83	1.22	1.30
4	A	1381	RHO	C22-C21	2.78	1.43	1.38
4	A	1381	RHO	C13-C14	-2.63	1.45	1.50
4	A	1381	RHO	C20-N2	2.33	1.46	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1381	RHO	C9-C14	2.31	1.47	1.44

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1381	RHO	OH3-C23-C18	5.52	130.98	115.28
4	A	1381	RHO	OH3-C23-OH2	-5.07	112.46	123.35
4	A	1381	RHO	C1-C-N	-4.63	116.23	121.33
4	A	1381	RHO	OH1-C9-C14	3.10	122.36	119.17
4	A	1381	RHO	OH1-C2-C3	3.09	124.38	121.57
4	A	1381	RHO	C21-C20-C19	3.02	122.50	118.62
4	A	1381	RHO	C21-C20-N2	-2.94	115.49	120.90
4	A	1381	RHO	OH1-C9-C10	-2.60	112.86	116.73
4	A	1381	RHO	C7-N-C	-2.58	116.12	120.51
3	A	1380	ADP	C5-C4-N3	2.58	130.26	126.72
3	A	1380	ADP	C4-N9-C8	-2.56	103.05	105.74
3	A	1380	ADP	C4-C5-N7	2.46	113.39	110.58
4	A	1381	RHO	C-C1-C2	-2.30	116.86	120.39
4	A	1381	RHO	OH2-C23-C18	-2.27	116.55	121.97
3	A	1380	ADP	C2-N1-C6	2.21	122.36	118.73
4	A	1381	RHO	C5-C-C1	2.18	123.45	119.12
4	A	1381	RHO	C22-C17-C18	2.16	121.69	119.26
4	A	1381	RHO	C12-C13-C14	2.16	117.57	112.29
3	A	1380	ADP	O3B-PB-O2B	2.15	115.87	107.80
3	A	1380	ADP	N3-C2-N1	-2.11	125.38	128.58
4	A	1381	RHO	C5-C4-C3	-2.07	117.45	120.86

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1381	RHO	C19-C18-C23-OH3


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.