



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:53 PM UTC

PDB ID : 4J6O / pdb_00004j6o
Title : Crystal Structure of the Phosphatase Domain of C. thermocellum (Bacterial) PnkP
Authors : Wang, L.; Smith, P.; Shuman, S.
Deposited on : 2013-02-11
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	2% 88% 11% .
1	B	264	5% 87% 10% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	B	504	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4686 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Metallophosphoesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	261	Total	C	N	O	S	0	12	0
			2127	1362	363	394	8			
1	B	258	Total	C	N	O	S	0	6	0
			2068	1320	351	388	9			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	169	SER	-	expression tag	UNP A3DJ38
A	170	MET	-	expression tag	UNP A3DJ38
A	425	LEU	-	expression tag	UNP A3DJ38
A	426	GLU	-	expression tag	UNP A3DJ38
A	427	HIS	-	expression tag	UNP A3DJ38
A	428	HIS	-	expression tag	UNP A3DJ38
A	429	HIS	-	expression tag	UNP A3DJ38
A	430	HIS	-	expression tag	UNP A3DJ38
A	431	HIS	-	expression tag	UNP A3DJ38
A	432	HIS	-	expression tag	UNP A3DJ38
B	169	SER	-	expression tag	UNP A3DJ38
B	170	MET	-	expression tag	UNP A3DJ38
B	425	LEU	-	expression tag	UNP A3DJ38
B	426	GLU	-	expression tag	UNP A3DJ38
B	427	HIS	-	expression tag	UNP A3DJ38
B	428	HIS	-	expression tag	UNP A3DJ38
B	429	HIS	-	expression tag	UNP A3DJ38
B	430	HIS	-	expression tag	UNP A3DJ38
B	431	HIS	-	expression tag	UNP A3DJ38
B	432	HIS	-	expression tag	UNP A3DJ38

- Molecule 2 is CITRIC ACID (CCD ID: CIT) (formula: C₆H₈O₇).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		
2	A	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	8	5		
4	A	1	Total	C	O	0	0
			13	8	5		
4	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0

- Molecule 6 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Mn 1 1	0	0
6	B	1	Total Mn 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	205	Total O 205 205	0	0
7	B	148	Total O 148 148	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Metallophosphoesterase



- Molecule 1: Metallophosphoesterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.70Å 38.66Å 92.95Å 90.00° 91.80° 90.00°	Depositor
Resolution (Å)	38.69 – 1.60 38.69 – 1.60	Depositor EDS
% Data completeness (in resolution range)	96.1 (38.69-1.60) 99.0 (38.69-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.8.1 _1168	Depositor
R, R_{free}	0.164 , 0.206 0.168 , 0.210	Depositor DCC
R_{free} test set	3170 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.325	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.022 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4686	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, CIT, GOL, PG4, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/2198	0.94	1/2967 (0.0%)
1	B	0.72	0/2120	0.93	0/2861
All	All	0.76	0/4318	0.93	1/5828 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	252	VAL	N-CA-C	-5.05	105.62	110.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2127	0	2139	20	0
1	B	2068	0	2081	25	0
2	A	26	0	10	0	0
2	B	13	0	5	0	0
3	A	10	0	14	5	0
4	A	26	0	36	6	0
4	B	13	0	18	1	0
5	A	18	0	24	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	30	0	40	7	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	205	0	0	4	0
7	B	148	0	0	8	0
All	All	4686	0	4367	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:370:ALA:HB2	3:A:502:PGE:H52	1.55	0.88
1:B:377[B]:THR:HG23	7:B:629:HOH:O	1.85	0.76
1:A:329:GLU:HA	4:A:504:PG4:H61	1.71	0.72
1:B:340:GLU:OE2	7:B:639:HOH:O	2.08	0.71
1:B:346:GLU:OE2	5:B:506:GOL:O2	2.08	0.70
1:B:378:PRO:HG2	5:B:504:GOL:H11	1.75	0.67
1:A:288:GLU:OE2	7:A:661:HOH:O	2.14	0.65
1:B:348[A]:THR:HG23	1:B:350:GLU:H	1.63	0.64
1:B:273:ASN:ND2	5:B:508:GOL:O3	2.27	0.62
1:B:377[B]:THR:HB	5:B:504:GOL:H32	1.81	0.61
1:B:174:ASN:HB3	7:B:717:HOH:O	1.99	0.61
4:A:503:PG4:H32	4:A:503:PG4:H82	1.85	0.59
1:B:219[B]:ARG:NH1	7:B:695:HOH:O	2.31	0.59
1:A:263[A]:ASN:ND2	5:A:506:GOL:H2	2.18	0.58
1:A:172:TRP:HB3	1:A:309[B]:VAL:HG11	1.87	0.57
4:A:503:PG4:H52	7:A:737:HOH:O	2.05	0.56
1:B:376:HIS:O	7:B:602:HOH:O	2.18	0.55
1:A:263[A]:ASN:HD22	5:A:506:GOL:H2	1.72	0.55
1:A:173:ASN:ND2	7:A:714:HOH:O	2.31	0.55
1:A:263[B]:ASN:ND2	5:A:506:GOL:O1	2.40	0.55
1:B:267:LYS:HD2	1:B:283:LEU:HD22	1.89	0.54
1:A:245:LEU:HB3	1:A:249:MET:HE2	1.90	0.53
1:B:245:LEU:HB3	1:B:249:MET:HE2	1.91	0.53
1:B:339:ARG:NH1	7:B:684:HOH:O	2.41	0.53
1:B:378:PRO:HD2	5:B:504:GOL:H32	1.91	0.52
1:B:173[B]:ASN:OD1	1:B:332:GLY:HA2	2.10	0.52
1:B:377[A]:THR:HB	5:B:504:GOL:H32	1.92	0.51
1:A:384[B]:LYS:HG2	1:A:389:ILE:HD12	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:370:ALA:CB	3:A:502:PGE:H52	2.37	0.49
1:A:369:LYS:HD3	3:A:502:PGE:H5	1.95	0.48
1:B:347:THR:OG1	5:B:505:GOL:H32	2.14	0.48
1:B:270:ARG:NH2	7:B:684:HOH:O	2.46	0.47
1:A:379:GLN:O	1:A:393:THR:HA	2.14	0.47
1:A:201:LYS:O	4:A:503:PG4:H62	2.14	0.47
1:A:405:ARG:CD	4:A:503:PG4:H41	2.45	0.47
4:B:503:PG4:H11	4:B:503:PG4:H31	1.57	0.46
3:A:502:PGE:H4	3:A:502:PGE:H6	1.50	0.46
1:B:245:LEU:C	1:B:249:MET:HE2	2.41	0.45
1:A:266[B]:VAL:HG11	1:A:343:LEU:HD13	1.97	0.45
1:A:410:GLU:HB3	4:A:503:PG4:H11	1.98	0.45
3:A:502:PGE:H62	7:A:719:HOH:O	2.18	0.44
1:B:170[B]:MET:HE3	1:B:170[B]:MET:HB3	1.85	0.44
1:A:224:GLU:HG2	5:A:507:GOL:H32	2.00	0.43
1:B:425:LEU:N	1:B:426:GLU:HA	2.32	0.42
1:B:290:LEU:HD23	1:B:290:LEU:HA	1.86	0.42
1:B:379:GLN:O	1:B:393:THR:HA	2.20	0.42
1:A:184:ILE:O	1:A:403:ALA:HA	2.20	0.42
1:A:235:VAL:O	1:A:236:ASP:HB2	2.21	0.41
1:B:227:LYS:NZ	7:B:699:HOH:O	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/264 (103%)	262 (97%)	9 (3%)	0	100	100
1	B	262/264 (99%)	243 (93%)	19 (7%)	0	100	100
All	All	533/528 (101%)	505 (95%)	28 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/222 (103%)	226 (99%)	3 (1%)	61	40
1	B	222/222 (100%)	220 (99%)	2 (1%)	70	55
All	All	451/444 (102%)	446 (99%)	5 (1%)	63	46

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	170	MET
1	A	213	VAL
1	A	298	ILE
1	B	213	VAL
1	B	425	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	429	HIS
1	B	174	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 2 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CIT	A	509	6	12,12,12	1.14	1 (8%)	17,17,17	4.18	10 (58%)
4	PG4	A	503	-	12,12,12	0.68	0	11,11,11	1.07	1 (9%)
2	CIT	B	502	6	12,12,12	1.06	0	17,17,17	3.49	10 (58%)
5	GOL	A	507	-	5,5,5	0.36	0	5,5,5	0.74	0
5	GOL	B	506	-	5,5,5	0.47	0	5,5,5	0.56	0
5	GOL	B	507	-	5,5,5	0.52	0	5,5,5	0.56	0
4	PG4	A	504	-	12,12,12	0.67	0	11,11,11	0.80	0
5	GOL	B	505	-	5,5,5	0.42	0	5,5,5	0.37	0
3	PGE	A	502	-	9,9,9	0.65	0	8,8,8	0.79	0
4	PG4	B	503	-	12,12,12	0.62	0	11,11,11	0.94	1 (9%)
5	GOL	B	508	-	5,5,5	0.32	0	5,5,5	0.45	0
5	GOL	A	505	-	5,5,5	0.33	0	5,5,5	0.43	0
5	GOL	A	506	-	5,5,5	0.34	0	5,5,5	0.48	0
5	GOL	B	504	-	5,5,5	0.41	0	5,5,5	0.39	0
2	CIT	A	501	-	12,12,12	1.12	0	17,17,17	1.68	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	A	509	6	-	0/16/16/16	-
4	PG4	A	503	-	-	6/10/10/10	-
2	CIT	B	502	6	-	0/16/16/16	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	507	-	-	4/4/4/4	-
5	GOL	B	506	-	-	1/4/4/4	-
5	GOL	B	507	-	-	2/4/4/4	-
4	PG4	A	504	-	-	5/10/10/10	-
5	GOL	B	505	-	-	2/4/4/4	-
3	PGE	A	502	-	-	5/7/7/7	-
4	PG4	B	503	-	-	8/10/10/10	-
5	GOL	B	508	-	-	1/4/4/4	-
5	GOL	A	505	-	-	2/4/4/4	-
5	GOL	A	506	-	-	0/4/4/4	-
5	GOL	B	504	-	-	4/4/4/4	-
2	CIT	A	501	-	-	3/16/16/16	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	509	CIT	O1-C1	2.30	1.29	1.22

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	509	CIT	C3-C4-C5	-8.80	89.86	113.92
2	A	509	CIT	O3-C5-C4	8.59	147.29	122.95
2	B	502	CIT	O5-C6-C3	-8.34	105.93	122.09
2	A	509	CIT	O4-C5-O3	-6.81	105.83	123.33
2	B	502	CIT	O7-C3-C6	6.38	118.00	108.96
2	A	509	CIT	C4-C3-C2	-5.17	96.06	109.31
2	A	509	CIT	C4-C3-C6	5.13	121.39	110.03
2	B	502	CIT	C4-C3-C6	4.55	120.09	110.03
2	A	501	CIT	O6-C6-C3	4.09	120.98	113.14
2	B	502	CIT	C3-C4-C5	-3.54	104.25	113.92
2	B	502	CIT	C2-C3-C6	-3.53	102.23	110.03
2	B	502	CIT	O6-C6-O5	3.45	134.89	123.86
2	A	509	CIT	O2-C1-O1	-3.24	115.00	123.33
2	B	502	CIT	O6-C6-C3	3.13	119.15	113.14
2	A	509	CIT	O5-C6-C3	-3.02	116.25	122.09
2	B	502	CIT	O7-C3-C2	-2.90	102.76	109.38
2	A	509	CIT	O2-C1-C2	2.66	122.76	114.35
2	A	501	CIT	O4-C5-C4	2.54	122.39	114.35
2	A	509	CIT	O7-C3-C2	-2.49	103.70	109.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	PG4	C5-O3-C4	2.44	123.93	113.26
2	A	509	CIT	O4-C5-C4	-2.36	106.87	114.35
2	B	502	CIT	O4-C5-C4	-2.15	107.53	114.35
2	A	501	CIT	O4-C5-O3	-2.09	117.96	123.33
2	B	502	CIT	C3-C2-C1	2.04	119.50	113.92
4	B	503	PG4	C3-O2-C2	2.03	122.13	113.26

There are no chirality outliers.

All (43) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	507	GOL	O1-C1-C2-C3
4	B	503	PG4	C1-C2-O2-C3
4	A	504	PG4	O2-C3-C4-O3
4	B	503	PG4	O2-C3-C4-O3
4	B	503	PG4	O3-C5-C6-O4
3	A	502	PGE	C6-C5-O3-C4
4	A	503	PG4	O4-C7-C8-O5
5	A	505	GOL	C1-C2-C3-O3
5	A	507	GOL	C1-C2-C3-O3
5	B	504	GOL	O1-C1-C2-C3
5	B	504	GOL	C1-C2-C3-O3
5	B	505	GOL	C1-C2-C3-O3
5	B	507	GOL	C1-C2-C3-O3
5	A	507	GOL	O1-C1-C2-O2
5	A	507	GOL	O2-C2-C3-O3
5	B	504	GOL	O1-C1-C2-O2
5	B	504	GOL	O2-C2-C3-O3
3	A	502	PGE	O1-C1-C2-O2
3	A	502	PGE	O3-C5-C6-O4
4	B	503	PG4	O1-C1-C2-O2
5	B	507	GOL	O2-C2-C3-O3
4	A	503	PG4	O3-C5-C6-O4
4	B	503	PG4	C3-C4-O3-C5
4	B	503	PG4	O4-C7-C8-O5
5	B	505	GOL	O2-C2-C3-O3
4	A	503	PG4	C1-C2-O2-C3
2	A	501	CIT	C1-C2-C3-C4
4	A	504	PG4	C3-C4-O3-C5
5	B	506	GOL	C1-C2-C3-O3
3	A	502	PGE	C3-C4-O3-C5
3	A	502	PGE	C4-C3-O2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	503	PG4	C6-C5-O3-C4
4	A	504	PG4	C1-C2-O2-C3
2	A	501	CIT	C1-C2-C3-O7
2	A	501	CIT	C1-C2-C3-C6
4	A	503	PG4	C8-C7-O4-C6
5	A	505	GOL	O2-C2-C3-O3
4	B	503	PG4	C8-C7-O4-C6
4	A	503	PG4	O2-C3-C4-O3
4	A	503	PG4	C3-C4-O3-C5
4	A	504	PG4	O3-C5-C6-O4
5	B	508	GOL	O1-C1-C2-C3
4	A	504	PG4	C8-C7-O4-C6

There are no ring outliers.

10 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	503	PG4	5	0
5	A	507	GOL	1	0
5	B	506	GOL	1	0
4	A	504	PG4	1	0
5	B	505	GOL	1	0
3	A	502	PGE	5	0
4	B	503	PG4	1	0
5	B	508	GOL	1	0
5	A	506	GOL	3	0
5	B	504	GOL	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/264 (98%)	-0.04	4 (1%) 72 76	15, 33, 63, 138	12 (4%)
1	B	258/264 (97%)	0.14	12 (4%) 36 38	13, 38, 86, 120	6 (2%)
All	All	519/528 (98%)	0.05	16 (3%) 51 54	13, 35, 76, 138	18 (3%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	429	HIS	3.2
1	A	170	MET	3.0
1	B	291	ALA	2.8
1	B	170[A]	MET	2.7
1	B	424	ALA	2.6
1	A	172	TRP	2.6
1	A	169	SER	2.4
1	B	349	GLY	2.3
1	B	423	PRO	2.3
1	B	354	TYR	2.3
1	B	213	VAL	2.3
1	B	274	GLY	2.3
1	B	425	LEU	2.3
1	B	172	TRP	2.2
1	B	171	LEU	2.2
1	B	277	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	B	505	6/6	0.63	0.20	68,69,70,71	6
5	GOL	B	506	6/6	0.66	0.15	70,71,72,72	6
4	PG4	B	503	13/13	0.68	0.17	49,58,58,59	13
5	GOL	A	505	6/6	0.72	0.10	63,63,64,65	0
4	PG4	A	504	13/13	0.75	0.13	59,59,63,63	13
5	GOL	B	504	6/6	0.75	0.13	58,60,61,63	0
5	GOL	B	508	6/6	0.76	0.12	61,63,64,65	6
5	GOL	B	507	6/6	0.81	0.13	59,60,60,61	6
2	CIT	A	501	13/13	0.81	0.11	55,70,74,83	13
5	GOL	A	506	6/6	0.86	0.12	68,68,70,71	0
3	PGE	A	502	10/10	0.86	0.13	36,42,47,47	10
5	GOL	A	507	6/6	0.87	0.10	64,65,65,66	0
4	PG4	A	503	13/13	0.90	0.10	34,38,40,41	13
2	CIT	A	509	13/13	0.95	0.07	25,27,35,39	0
2	CIT	B	502	13/13	0.96	0.05	28,30,37,40	0
6	MN	A	508	1/1	1.00	0.02	18,18,18,18	0
6	MN	B	501	1/1	1.00	0.02	26,26,26,26	0

6.5 Other polymers ⓘ

There are no such residues in this entry.