



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:22 PM UTC

PDB ID : 5J6D / pdb_00005j6d
Title : Discovery of acyl guanidine tryptophan hydroxylase-1 inhibitors
Authors : Stein, A.J.; Goldberg, D.R.; De Lombaert, S.
Deposited on : 2016-04-04
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	4% 93% 5%
1	B	307	41% 71% 9% 19%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan 5-hydroxylase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	292	Total	C	N	O	S	0	1	0
			2317	1494	385	425	13			
1	B	248	Total	C	N	O	S	0	1	0
			1768	1136	293	326	13			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	403	HIS	-	expression tag	UNP P17752
A	404	HIS	-	expression tag	UNP P17752
A	405	HIS	-	expression tag	UNP P17752
A	406	HIS	-	expression tag	UNP P17752
A	407	HIS	-	expression tag	UNP P17752
A	408	HIS	-	expression tag	UNP P17752
B	403	HIS	-	expression tag	UNP P17752
B	404	HIS	-	expression tag	UNP P17752
B	405	HIS	-	expression tag	UNP P17752
B	406	HIS	-	expression tag	UNP P17752
B	407	HIS	-	expression tag	UNP P17752
B	408	HIS	-	expression tag	UNP P17752

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	B	1	Total	Fe	0	0
			1	1		

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		
3	B	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is 4-[(N-{[2-(3-methoxyphenoxy)-6-(piperidin-1-yl)phenyl]methyl}carbamimidoyl)carbamoyl]-L-phenylalanine (CCD ID: 6H5) (formula: C₃₀H₃₅N₅O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			40	30	5	5		
5	B	1	Total	C	N	O	0	0
			40	30	5	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	123	Total	O	0	0
			123	123		
6	B	17	Total	O	0	0
			17	17		



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tryptophan 5-hydroxylase 1



• Molecule 1: Tryptophan 5-hydroxylase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.73Å 63.56Å 156.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.90 50.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-1.90) 99.9 (50.00-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.210 , 0.248 0.218 , 0.255	Depositor DCC
R_{free} test set	2245 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	33.2	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4335	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, TRS, FE, 6H5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/2384	0.86	0/3230
1	B	0.73	0/1812	1.13	7/2472 (0.3%)
All	All	0.75	0/4196	0.99	7/5702 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	124[A]	MET	CA-C-O	-16.12	102.90	120.80
1	B	124[B]	MET	CA-C-O	-16.12	102.90	120.80
1	B	368	THR	N-CA-C	9.63	125.17	109.96
1	B	124[A]	MET	N-CA-C	9.25	124.94	109.76
1	B	124[B]	MET	N-CA-C	9.25	124.94	109.76
1	B	227	PHE	N-CA-C	-6.30	99.45	109.59
1	B	189	ALA	N-CA-C	5.20	118.41	110.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	367	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2317	0	2236	3	0
1	B	1768	0	1518	26	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	8	0	9	0	0
3	B	8	0	10	0	0
4	A	6	0	8	0	0
4	B	6	0	8	0	0
5	A	40	0	0	0	0
5	B	40	0	0	0	0
6	A	123	0	0	0	0
6	B	17	0	0	0	0
All	All	4335	0	3789	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:124[B]:MET:HE3	1:B:124[B]:MET:HA	1.36	1.07
1:B:221:LEU:HD11	1:B:278:VAL:HG13	1.54	0.89
1:B:182:ASN:O	1:B:186:PRO:HD3	1.84	0.77
1:B:124[B]:MET:HE3	1:B:124[B]:MET:CA	2.15	0.76
1:B:124[B]:MET:HA	1:B:124[B]:MET:CE	2.15	0.74
1:B:221:LEU:HD21	1:B:278:VAL:HG13	1.73	0.69
1:B:200:LEU:HD22	1:B:220:PHE:CE2	2.31	0.66
1:B:124[B]:MET:CA	1:B:124[B]:MET:CE	2.74	0.65
1:B:220:PHE:CD1	1:B:220:PHE:C	2.80	0.59
1:B:217:VAL:O	1:B:221:LEU:HG	2.05	0.57
1:B:200:LEU:HD22	1:B:220:PHE:HE2	1.74	0.51
1:B:221:LEU:HD21	1:B:278:VAL:CG1	2.40	0.50
1:B:221:LEU:HD11	1:B:278:VAL:CG1	2.33	0.49
1:B:221:LEU:CD1	1:B:278:VAL:HG13	2.34	0.47
1:A:272:HIS:HA	1:A:336:SER:CB	2.45	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:218:SER:HB2	1:B:229:ILE:HB	1.96	0.46
1:B:266:PRO:HB2	1:B:367:THR:HG22	1.98	0.45
1:B:183:LYS:O	1:B:186:PRO:HD2	2.18	0.44
1:B:102:MET:HE3	1:B:102:MET:HA	2.00	0.44
1:B:125:TYR:OH	1:B:267:GLU:OE2	2.30	0.43
1:B:215:GLU:OE2	1:B:219:ASN:ND2	2.51	0.43
1:B:181:LEU:O	1:B:185:TYR:N	2.51	0.43
1:B:222:LYS:O	1:B:226:GLY:HA2	2.19	0.43
1:B:200:LEU:HD22	1:B:220:PHE:CZ	2.54	0.43
1:A:380:GLU:OE1	1:A:383:LYS:NZ	2.45	0.43
1:B:248:ARG:NE	1:B:291:GLN:OE1	2.44	0.41
1:B:217:VAL:HG11	1:B:278:VAL:HG21	2.03	0.41
1:A:122:VAL:HG21	1:A:233:ALA:HB3	2.02	0.41
1:B:188:HIS:O	1:B:329:VAL:HG12	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/307 (95%)	288 (99%)	3 (1%)	0	100	100
1	B	233/307 (76%)	226 (97%)	7 (3%)	0	100	100
All	All	524/614 (85%)	514 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/273 (91%)	248 (100%)	0	100	100
1	B	160/273 (59%)	156 (98%)	4 (2%)	42	37
All	All	408/546 (75%)	404 (99%)	4 (1%)	68	70

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	187	THR
1	B	220	PHE
1	B	227	PHE
1	B	228	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	179	GLN
1	B	117	HIS
1	B	251	HIS
1	B	258	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	TRS	A	502	2	7,7,7	0.42	0	9,9,9	0.41	0
4	GOL	B	503	-	5,5,5	0.61	0	5,5,5	0.57	0
3	TRS	B	502	2	7,7,7	0.51	0	9,9,9	0.69	0
4	GOL	A	503	-	5,5,5	0.30	0	5,5,5	0.42	0
5	6H5	A	504	-	40,43,43	0.84	1 (2%)	50,58,58	2.40	5 (10%)
5	6H5	B	504	-	40,43,43	0.79	1 (2%)	50,58,58	2.53	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	A	502	2	-	1/9/9/9	-
4	GOL	B	503	-	-	2/4/4/4	-
3	TRS	B	502	2	-	0/9/9/9	-
4	GOL	A	503	-	-	2/4/4/4	-
5	6H5	A	504	-	-	4/31/39/39	0/4/4/4
5	6H5	B	504	-	-	2/31/39/39	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	504	6H5	C8-N1	-3.18	1.28	1.36
5	B	504	6H5	C8-N1	-2.95	1.29	1.36

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	504	6H5	C10-C22-N4	13.77	126.70	118.63
5	B	504	6H5	C10-C22-N4	13.22	126.37	118.63
5	B	504	6H5	C27-N4-C23	6.13	125.35	111.57
5	A	504	6H5	C27-N4-C23	5.84	124.71	111.57

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	504	6H5	C21-C22-N4	-5.32	114.08	122.42
5	B	504	6H5	C21-C22-N4	-4.84	114.84	122.42
5	B	504	6H5	C15-O2-C14	4.60	127.37	117.50
5	B	504	6H5	C6-C7-N1	3.69	121.57	116.25
5	B	504	6H5	C11-O1-C12	3.30	125.93	117.95
5	A	504	6H5	C11-O1-C12	2.39	123.75	117.95
5	A	504	6H5	C6-C7-N1	2.06	119.21	116.25

There are no chirality outliers.


All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	503	GOL	C1-C2-C3-O3
4	B	503	GOL	C1-C2-C3-O3
5	A	504	6H5	C13-C14-O2-C15
5	A	504	6H5	C16-C14-O2-C15
4	A	503	GOL	O2-C2-C3-O3
4	B	503	GOL	O2-C2-C3-O3
5	B	504	6H5	C21-C22-N4-C23
5	A	504	6H5	C21-C22-N4-C27
5	A	504	6H5	C10-C22-N4-C27
5	B	504	6H5	C10-C22-N4-C23
3	A	502	TRS	C1-C-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	292/307 (95%)	0.25	12 (4%) 41 44	22, 33, 54, 82	1 (0%)
1	B	248/307 (80%)	2.20	126 (50%) 0 0	27, 71, 96, 108	1 (0%)
All	All	540/614 (87%)	1.15	138 (25%) 1 1	22, 44, 88, 108	2 (0%)

All (138) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	200	LEU	6.3
1	B	286	PHE	6.2
1	B	199	LEU	6.1
1	B	188	HIS	5.7
1	B	220	PHE	5.5
1	B	274	LEU	5.4
1	B	221	LEU	5.2
1	B	177	VAL	5.2
1	B	194	LEU	4.8
1	B	338	ILE	4.7
1	B	173	THR	4.7
1	B	329	VAL	4.4
1	B	181	LEU	4.4
1	B	354	PHE	4.4
1	B	366	ILE	4.3
1	B	289	PHE	4.3
1	B	189	ALA	4.3
1	B	280	LEU	4.2
1	B	174	TRP	4.2
1	B	381	ASP	4.2
1	B	345	LEU	4.1
1	B	184	LEU	4.1
1	B	282	ALA	4.0
1	B	178	PHE	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	182	ASN	4.0
1	B	164	VAL	4.0
1	B	262	PRO	4.0
1	B	141	TYR	3.9
1	B	264	TYR	3.9
1	B	358	ILE	3.9
1	B	151	LEU	3.8
1	B	341	LEU	3.8
1	A	133	HIS	3.8
1	B	217	VAL	3.8
1	B	314	PHE	3.8
1	B	379	PHE	3.8
1	B	321	CYS	3.7
1	B	185	TYR	3.6
1	B	279	PRO	3.6
1	B	197	LEU	3.6
1	B	367	THR	3.6
1	B	190	CYS	3.5
1	B	371	ASP	3.5
1	B	281	LEU	3.4
1	B	147	TYR	3.4
1	B	255	TYR	3.4
1	B	275	LEU	3.4
1	B	265	THR	3.4
1	B	382	ALA	3.4
1	B	335	LEU	3.4
1	B	176	THR	3.4
1	B	193	TYR	3.3
1	B	278	VAL	3.3
1	B	216	ASP	3.3
1	B	287	ALA	3.2
1	A	134	PRO	3.2
1	B	227	PHE	3.2
1	B	336	SER	3.1
1	B	318	PHE	3.1
1	B	374	PHE	3.1
1	B	271	CYS	3.1
1	B	391	LYS	3.1
1	B	198	PRO	3.1
1	B	368	THR	3.0
1	B	142	ARG	3.0
1	B	330	PHE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	161	ILE	3.0
1	B	364	CYS	2.9
1	B	249	VAL	2.9
1	A	136	PHE	2.9
1	B	263	PHE	2.9
1	B	183	LYS	2.9
1	B	285	SER	2.9
1	B	351	VAL	2.9
1	B	369	PHE	2.9
1	A	102	MET	2.9
1	B	102	MET	2.9
1	B	112	ILE	2.9
1	B	229	ILE	2.9
1	B	320	LEU	2.9
1	B	191	ARG	2.8
1	B	344	ALA	2.8
1	B	247	PHE	2.8
1	B	337	SER	2.8
1	A	132	ASP	2.7
1	B	196	ASN	2.7
1	B	130	ASP	2.7
1	B	160	PRO	2.7
1	B	162	PRO	2.7
1	B	372	VAL	2.7
1	A	199	LEU	2.7
1	B	350	LYS	2.6
1	A	327	LEU	2.6
1	A	131	ALA	2.6
1	B	233	ALA	2.6
1	B	389	PHE	2.6
1	B	390	THR	2.6
1	B	226	GLY	2.6
1	B	154	ASN	2.6
1	A	169	GLU	2.5
1	B	266	PRO	2.5
1	B	334	LEU	2.5
1	B	257	ARG	2.5
1	A	201	SER	2.4
1	B	187	THR	2.4
1	B	384	GLU	2.4
1	B	268	PRO	2.4
1	B	195	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	365	LEU	2.4
1	B	356	PRO	2.4
1	B	140	VAL	2.4
1	B	319	GLY	2.4
1	B	186	PRO	2.3
1	B	155	TYR	2.3
1	B	149	ALA	2.3
1	B	346	SER	2.3
1	B	144	ARG	2.3
1	B	254	GLN	2.3
1	B	214	LEU	2.3
1	B	117	HIS	2.3
1	B	328	ARG	2.2
1	A	393	ILE	2.2
1	B	304	ALA	2.2
1	B	308	LEU	2.2
1	B	103	GLU	2.1
1	B	125	TYR	2.1
1	A	326	GLN	2.1
1	B	253	THR	2.1
1	B	223	GLU	2.1
1	B	342	LYS	2.1
1	B	180	GLU	2.1
1	B	175	GLY	2.1
1	B	316	VAL	2.1
1	B	270	THR	2.0
1	B	129	LEU	2.0
1	B	259	SER	2.0
1	B	225	THR	2.0
1	B	256	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	A	503	6/6	0.80	0.15	39,47,51,52	0
5	6H5	B	504	40/40	0.89	0.12	43,49,62,64	0
3	TRS	B	502	8/8	0.93	0.09	46,48,50,55	0
4	GOL	B	503	6/6	0.94	0.08	40,43,46,47	0
5	6H5	A	504	40/40	0.95	0.07	25,31,37,49	0
3	TRS	A	502	8/8	0.95	0.08	24,24,26,26	0
2	FE	B	501	1/1	0.99	0.04	48,48,48,48	0
2	FE	A	501	1/1	1.00	0.04	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 6H5 A 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.