



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:09 AM UTC

PDB ID : 5J6F / pdb_00005j6f
Title : Crystal structure of DAH7PS-CM complex from Geobacillus sp. with prephenate
Authors : Nazmi, A.R.; Othman, M.; Lang, E.J.M.; Bai, Y.; Allison, T.M.; Panjkar, S.; Arcus, V.L.; Parker, E.J.
Deposited on : 2016-04-04
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	 79% 15% . .
1	B	368	 76% 14% . 8%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase, chorismate mutase-isozyme 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	352	Total	C	N	O	S	0	0	0
			2741	1730	485	516	10			
1	B	340	Total	C	N	O	S	0	0	0
			2625	1657	463	494	11			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	GLY	SER	conflict	UNP L8A208
A	359	ARG	GLN	conflict	UNP L8A208
A	361	LEU	-	expression tag	UNP L8A208
A	362	GLU	-	expression tag	UNP L8A208
A	363	HIS	-	expression tag	UNP L8A208
A	364	HIS	-	expression tag	UNP L8A208
A	365	HIS	-	expression tag	UNP L8A208
A	366	HIS	-	expression tag	UNP L8A208
A	367	HIS	-	expression tag	UNP L8A208
A	368	HIS	-	expression tag	UNP L8A208
B	2	GLY	SER	conflict	UNP L8A208
B	359	ARG	GLN	conflict	UNP L8A208
B	361	LEU	-	expression tag	UNP L8A208
B	362	GLU	-	expression tag	UNP L8A208
B	363	HIS	-	expression tag	UNP L8A208
B	364	HIS	-	expression tag	UNP L8A208
B	365	HIS	-	expression tag	UNP L8A208
B	366	HIS	-	expression tag	UNP L8A208
B	367	HIS	-	expression tag	UNP L8A208
B	368	HIS	-	expression tag	UNP L8A208

- Molecule 2 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mn	0	0
			1	1		
2	B	1	Total	Mn	0	0
			1	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is PREPHENIC ACID (CCD ID: PRE) (formula: C₁₀H₁₀O₆).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 16 10 6	0	0
4	B	1	Total C O 16 10 6	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase, chorismate mutase-isozyme 3



- Molecule 1: 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase, chorismate mutase-isozyme 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	95.44Å 95.44Å 167.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	82.65 – 2.75 82.65 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.0 (82.65-2.75) 99.0 (82.65-2.75)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.75Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.182 , 0.235 0.186 , 0.232	Depositor DCC
R_{free} test set	1054 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å ²)	74.8	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 30.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.095 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5420	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PRE, MN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.37	9/2781 (0.3%)	1.32	13/3755 (0.3%)
1	B	1.31	6/2661 (0.2%)	1.31	15/3593 (0.4%)
All	All	1.35	15/5442 (0.3%)	1.32	28/7348 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	353	PHE	N-CA	7.10	1.53	1.46
1	B	86	GLN	CA-C	6.54	1.61	1.52
1	A	320	MET	N-CA	-6.28	1.38	1.46
1	B	311	ALA	N-CA	-6.01	1.39	1.46
1	A	45	ARG	CA-C	6.00	1.60	1.52
1	A	287	HIS	N-CA	-5.72	1.39	1.46
1	A	61	HIS	CA-C	5.62	1.59	1.52
1	A	22	LYS	N-CA	-5.62	1.39	1.46
1	B	284	LYS	N-CA	-5.50	1.39	1.46
1	A	149	LEU	CA-C	-5.50	1.46	1.52
1	B	12	ARG	C-O	-5.35	1.17	1.24
1	A	158	THR	N-CA	-5.26	1.40	1.46
1	A	271	ARG	CA-CB	5.22	1.61	1.53
1	B	185	VAL	N-CA	-5.07	1.40	1.46
1	B	45	ARG	CD-NE	5.01	1.53	1.46

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	286	THR	N-CA-C	-9.01	96.03	109.81
1	A	58	ILE	N-CA-C	-7.78	102.75	110.30
1	A	80	LYS	CA-CB-CG	7.16	128.42	114.10
1	A	332	SER	CB-CA-C	-6.90	97.53	110.16
1	B	104	ASN	N-CA-C	6.76	120.51	109.76
1	A	103	GLU	N-CA-C	-6.68	101.53	110.55
1	A	59	SER	N-CA-C	6.68	120.30	111.75
1	A	353	PHE	N-CA-C	6.64	120.08	112.57
1	B	64	GLY	N-CA-C	-6.44	104.27	112.00
1	B	238	ILE	CB-CA-C	-6.12	104.13	111.97
1	B	334	SER	N-CA-C	6.04	118.64	111.33
1	B	85	LEU	N-CA-C	6.02	118.63	111.71
1	A	301	ARG	CG-CD-NE	5.96	125.11	112.00
1	A	76	LYS	CB-CA-C	-5.90	101.61	110.88
1	B	126	CYS	N-CA-C	-5.87	104.58	110.97
1	B	161	TYR	N-CA-C	-5.85	106.06	112.72
1	A	33	GLU	CB-CA-C	5.72	120.40	110.68
1	A	104	ASN	CB-CA-C	5.65	118.97	109.48
1	A	271	ARG	CB-CG-CD	5.61	124.21	111.30
1	B	195	ILE	CB-CA-C	-5.55	104.77	111.88
1	B	59	SER	N-CA-C	5.45	119.03	112.38
1	B	251	GLN	N-CA-C	5.41	119.58	112.92
1	A	80	LYS	CB-CA-C	-5.24	102.43	110.81
1	B	287	HIS	CB-CA-C	5.21	120.36	111.46
1	B	132	GLU	N-CA-C	-5.19	105.70	111.36
1	A	271	ARG	N-CA-C	-5.17	105.53	111.07
1	B	117	GLY	N-CA-C	-5.01	107.74	114.25
1	B	157	ARG	N-CA-C	5.00	118.16	107.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	156	PRO	Peptide
1	B	352	ALA	Peptide
1	B	353	PHE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2741	0	2773	23	0
1	B	2625	0	2650	10	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	10	0	0	0	0
3	B	10	0	0	1	0
4	A	16	0	8	0	0
4	B	16	0	8	0	0
All	All	5420	0	5439	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:ARG:HH12	1:A:104:ASN:HB3	1.52	0.74
1:A:41:GLN:HG3	1:A:43:THR:HG23	1.73	0.71
1:A:276:ILE:HD13	1:B:276:ILE:HD11	1.73	0.71
1:A:94:LEU:HD22	1:A:246:GLU:HG3	1.74	0.69
1:B:261:ARG:NE	3:B:402:SO4:O1	2.27	0.67
1:A:41:GLN:HG3	1:A:43:THR:CG2	2.33	0.59
1:A:123:MET:O	1:A:149:LEU:HD12	2.03	0.58
1:B:123:MET:O	1:B:149:LEU:HD12	2.04	0.57
1:A:101:HIS:HB3	1:A:287:HIS:CD2	2.40	0.56
1:A:305:ILE:HB	1:A:306:PRO:HD3	1.89	0.54
1:B:305:ILE:HB	1:B:306:PRO:HD3	1.90	0.53
1:B:188:GLU:HA	1:B:206:GLN:HB3	1.93	0.50
1:A:39:GLU:OE2	1:A:90:HIS:ND1	2.43	0.49
1:A:157:ARG:NH2	1:A:331:LEU:O	2.47	0.47
1:A:95:LEU:N	1:A:246:GLU:OE2	2.48	0.47
1:A:188:GLU:HA	1:A:206:GLN:HB3	1.97	0.46
1:B:194:ASP:HA	1:B:197:ILE:HD12	1.98	0.46
1:A:122:VAL:HB	1:A:320:MET:HG3	1.99	0.45
1:A:222:GLY:HA3	1:A:251:GLN:HB2	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:TYR:CD1	1:A:154:TYR:N	2.85	0.44
1:A:5:ARG:HA	1:A:8:GLU:HG2	1.97	0.44
1:B:302:ASP:OD1	1:B:302:ASP:N	2.42	0.43
1:A:194:ASP:HA	1:A:197:ILE:HD12	1.99	0.43
1:A:131:TYR:HH	1:A:180:GLU:CD	2.27	0.43
1:A:276:ILE:O	1:A:277:SER:C	2.62	0.42
1:A:350:VAL:O	1:A:351:ARG:C	2.63	0.42
1:B:5:ARG:HA	1:B:8:GLU:HG2	2.01	0.41
1:A:55:LEU:HD22	1:A:72:GLN:HG2	2.01	0.41
1:A:302:ASP:OD1	1:A:302:ASP:N	2.41	0.41
1:B:122:VAL:HB	1:B:320:MET:HG3	2.02	0.41
1:A:339:ASP:C	1:A:339:ASP:OD1	2.64	0.41
1:B:284:LYS:HD3	1:B:284:LYS:HA	1.92	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/368 (95%)	343 (98%)	7 (2%)	0	100	100
1	B	336/368 (91%)	332 (99%)	4 (1%)	0	100	100
All	All	686/736 (93%)	675 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/308 (94%)	272 (94%)	16 (6%)	19	36
1	B	273/308 (89%)	246 (90%)	27 (10%)	7	15
All	All	561/616 (91%)	518 (92%)	43 (8%)	12	22

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	LEU
1	A	57	LEU
1	A	63	ASP
1	A	107	VAL
1	A	109	VAL
1	A	113	ARG
1	A	141	VAL
1	A	150	ARG
1	A	191	THR
1	A	271	ARG
1	A	276	ILE
1	A	277	SER
1	A	284	LYS
1	A	302	ASP
1	A	340	ILE
1	A	350	VAL
1	B	12	ARG
1	B	15	GLU
1	B	21	LEU
1	B	45	ARG
1	B	53	LYS
1	B	63	ASP
1	B	67	GLU
1	B	107	VAL
1	B	109	VAL
1	B	132	GLU
1	B	141	VAL
1	B	150	ARG
1	B	157	ARG
1	B	158	THR
1	B	164	GLN
1	B	180	GLU
1	B	185	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	191	THR
1	B	239	GLU
1	B	271	ARG
1	B	276	ILE
1	B	277	SER
1	B	302	ASP
1	B	348	GLU
1	B	350	VAL
1	B	351	ARG
1	B	354	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	118	ASN
1	A	164	GLN
1	A	206	GLN
1	A	223	GLN
1	A	255	GLN
1	B	19	GLN
1	B	225	ASN
1	B	287	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	SO4	B	402	-	4,4,4	0.43	0	6,6,6	0.45	0
3	SO4	B	403	-	4,4,4	0.46	0	6,6,6	0.35	0
4	PRE	A	404	-	16,16,16	2.89	7 (43%)	14,23,23	6.00	6 (42%)
3	SO4	A	402	-	4,4,4	0.62	0	6,6,6	0.42	0
3	SO4	A	403	-	4,4,4	0.49	0	6,6,6	0.36	0
4	PRE	B	404	-	16,16,16	2.83	6 (37%)	14,23,23	5.89	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PRE	A	404	-	-	9/15/27/27	0/1/1/1
4	PRE	B	404	-	-	2/15/27/27	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	404	PRE	C1-C6	-6.35	1.42	1.50
4	B	404	PRE	C1-C6	-6.35	1.42	1.50
4	A	404	PRE	C1-C2	-6.10	1.42	1.50
4	B	404	PRE	C1-C2	-6.06	1.42	1.50
4	A	404	PRE	C8-C1	-3.55	1.49	1.56
4	A	404	PRE	O72-C7	-3.34	1.18	1.30
4	B	404	PRE	C8-C1	-3.25	1.50	1.56
4	B	404	PRE	O72-C7	-3.25	1.18	1.30
4	A	404	PRE	C5-C6	3.12	1.37	1.32
4	B	404	PRE	C5-C6	2.81	1.36	1.32
4	B	404	PRE	C3-C2	2.81	1.36	1.32
4	A	404	PRE	C3-C2	2.80	1.36	1.32
4	A	404	PRE	C1-C7	-2.41	1.50	1.53

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	404	PRE	O71-C7-C1	-16.80	95.98	121.96
4	A	404	PRE	O71-C7-C1	-16.67	96.18	121.96
4	A	404	PRE	O72-C7-O71	11.61	161.00	123.86
4	B	404	PRE	O72-C7-O71	11.00	159.04	123.86
4	A	404	PRE	O72-C7-C1	-6.43	102.82	114.86
4	B	404	PRE	C4-C5-C6	-5.44	118.93	123.40
4	B	404	PRE	O72-C7-C1	-5.28	104.97	114.86
4	B	404	PRE	C4-C3-C2	-4.40	119.78	123.40
4	A	404	PRE	C4-C5-C6	-4.02	120.09	123.40
4	A	404	PRE	O'L-C2'-C1'	-3.62	117.31	121.81
4	A	404	PRE	C4-C3-C2	-3.49	120.53	123.40

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	404	PRE	C6-C1-C7-O71
4	A	404	PRE	C6-C1-C7-O72
4	A	404	PRE	C8-C1-C7-O71
4	A	404	PRE	C8-C1-C7-O72
4	A	404	PRE	C8-C1'-C2'-O'M
4	B	404	PRE	C2-C1-C8-C1'
4	B	404	PRE	C7-C1-C8-C1'
4	A	404	PRE	O1'-C1'-C2'-O'L
4	A	404	PRE	C8-C1'-C2'-O'L
4	A	404	PRE	C2-C1-C7-O71
4	A	404	PRE	O1'-C1'-C2'-O'M

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	352/368 (95%)	-0.72	0	100 100	54, 74, 96, 118	0
1	B	340/368 (92%)	-0.58	3 (0%)	81 81	59, 77, 107, 161	0
All	All	692/736 (94%)	-0.65	3 (0%)	88 89	54, 75, 99, 161	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	102	PRO	2.5
1	B	161	TYR	2.2
1	B	160	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PRE	A	404	16/16	0.91	0.11	62,107,122,126	0
4	PRE	B	404	16/16	0.92	0.09	63,79,84,86	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	B	402	5/5	0.94	0.06	76,76,81,83	0
3	SO4	B	403	5/5	0.94	0.11	90,91,100,107	0
3	SO4	A	402	5/5	0.95	0.08	73,78,83,91	0
3	SO4	A	403	5/5	0.97	0.09	69,70,84,88	0
2	MN	A	401	1/1	0.99	0.02	74,74,74,74	0
2	MN	B	401	1/1	0.99	0.07	100,100,100,100	0

6.5 Other polymers [i](#)

There are no such residues in this entry.