



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:00 AM UTC

PDB ID : 8J66 / pdb_00008j66
Title : Crystal structure of glycosyltransferase SgUGT94-289-3 in complex with M3, state 2
Authors : Li, M.; Zhang, S.; Cui, S.
Deposited on : 2023-04-24
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	459	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3810 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycosyltransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	436	3464	2242	585	619	18	0	4	0

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (CCD ID: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	25	9	2	12	2	0	0

- Molecule 3 is (2S,3S,4S,5R,6R)-2-(hydroxymethyl)-6-[[[(2R,3S,4S,5R,6S)-6-[(3R,6S)-6-[(3S,8S,9R,10R,11S,13R,14S,17S)-3-[(2R,3R,4S,5S,6S)-6-(hydroxymethyl)-3,4,5-tris(oxidanyl)oxan-2-yl]oxy-4,4,9,13,14-pentamethyl-11-oxidanyl-2,3,7,8,10,11,12,15,16,17-decahydro-1H-cyclopenta[a]phenanthren-17-yl]-2-methyl-2-oxidanyl-heptan-3-yl]oxy-3,4,5-tris(oxidanyl)oxan-2-yl]methoxy]oxane-3,4,5-triol (CCD ID: M2Q) (formula: $C_{48}H_{82}O_{19}$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			67	48	19		
3	A	1	Total	C	O	0	0
			67	48	19		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	187	Total	O	0	0
			187	187		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glycosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.07Å 78.13Å 94.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.33 – 2.27 47.33 – 2.27	Depositor EDS
% Data completeness (in resolution range)	98.1 (47.33-2.27) 98.1 (47.33-2.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.91 (at 2.27Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.207 , 0.234 0.206 , 0.234	Depositor DCC
R_{free} test set	1225 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	27.1	Xtriage
Anisotropy	0.111	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3810	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, M2Q

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/3565	0.76	9/4836 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	145	GLY	CA-C-N	7.03	131.79	120.55
1	A	145	GLY	C-N-CA	7.03	131.79	120.55
1	A	26	PHE	CA-CB-CG	6.13	119.93	113.80
1	A	234	VAL	N-CA-C	-5.57	106.88	112.17
1	A	255	ASN	CB-CA-C	5.43	119.81	110.79
1	A	350	HIS	CA-C-N	5.43	129.84	122.19
1	A	350	HIS	C-N-CA	5.43	129.84	122.19
1	A	147	PHE	CA-CB-CG	5.14	118.94	113.80
1	A	421	ARG	O-C-N	5.13	127.36	122.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3464	0	3403	38	0
2	A	25	0	11	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	134	0	0	0	0
4	A	187	0	0	6	0
All	All	3810	0	3414	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:TYR:CE2	1:A:371:MET:HE2	2.06	0.91
1:A:236:PRO:HG3	1:A:431:LEU:HD22	1.58	0.85
1:A:188:ARG:N	4:A:604:HOH:O	2.20	0.74
1:A:144:THR:OG1	4:A:601:HOH:O	1.87	0.74
1:A:274:TYR:CZ	1:A:371:MET:HE2	2.31	0.65
1:A:86:GLY:HA2	1:A:275:PHE:HZ	1.64	0.63
1:A:365:PRO:HB3	1:A:420:VAL:HG12	1.81	0.62
1:A:13:LEU:HD13	1:A:110:LEU:HD23	1.81	0.61
1:A:146:VAL:HG21	1:A:220:MET:HG3	1.83	0.61
1:A:280:GLU:HG2	1:A:399:ILE:HD12	1.86	0.58
1:A:235:GLY:HA3	1:A:438:LYS:HD3	1.87	0.55
1:A:421:ARG:HD3	1:A:425:ARG:HH21	1.71	0.54
1:A:31:LYS:O	1:A:34:SER:HB3	2.10	0.52
1:A:146:VAL:HG21	1:A:220:MET:CG	2.39	0.52
1:A:23:LEU:HD11	1:A:48:ASN:CG	2.36	0.51
1:A:13:LEU:HD13	1:A:110:LEU:CD2	2.41	0.50
1:A:305:ASP:OD2	1:A:308:SER:OG	2.29	0.50
1:A:188:ARG:O	1:A:192:ARG:HG3	2.13	0.48
1:A:146:VAL:HG11	1:A:220:MET:HA	1.95	0.48
1:A:417:ARG:NH1	4:A:610:HOH:O	2.30	0.48
1:A:218:LYS:HG2	4:A:654:HOH:O	2.12	0.48
1:A:301:PHE:CD2	1:A:308:SER:OG	2.64	0.47
1:A:301:PHE:CE2	1:A:308:SER:HB2	2.50	0.46
1:A:86:GLY:O	1:A:303:GLN:HG3	2.15	0.46
1:A:148:VAL:HG23	1:A:377:PHE:CD2	2.51	0.46
1:A:174:TRP:HD1	1:A:177:MET:HE2	1.79	0.46
1:A:236:PRO:HA	1:A:357[A]:MET:HE3	1.98	0.46
1:A:274:TYR:HE2	1:A:371:MET:HE2	1.74	0.45
1:A:281:MET:HE1	1:A:299:VAL:HG13	1.99	0.45
1:A:310:ILE:N	4:A:603:HOH:O	2.16	0.45
1:A:360:MET:CE	1:A:387:VAL:HG21	2.47	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:360:MET:O	1:A:428:SER:HB2	2.18	0.44
1:A:215:LEU:HD13	1:A:360:MET:HE1	2.00	0.44
1:A:270:PHE:HA	1:A:350:HIS:HB3	1.99	0.43
1:A:14:MET:HE2	1:A:26:PHE:CD2	2.55	0.42
1:A:43:CYS:HA	1:A:67:VAL:O	2.20	0.41
1:A:170:LEU:HD23	1:A:175:LYS:HG2	2.03	0.41
1:A:144:THR:HB	4:A:702:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	434/459 (95%)	428 (99%)	5 (1%)	1 (0%)	43 53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	373	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	377/399 (94%)	374 (99%)	3 (1%)	73 84

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	209[A]	ILE
1	A	209[B]	ILE
1	A	307	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	HIS
1	A	65	GLN
1	A	172	ASN
1	A	242	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	M2Q	A	503	-	73,73,73	4.74	18 (24%)	111,117,117	2.17	29 (26%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UDP	A	501	-	25,26,26	1.23	1 (4%)	38,40,40	0.72	0
3	M2Q	A	502	-	73,73,73	4.74	24 (32%)	111,117,117	3.54	28 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	M2Q	A	503	-	-	10/32/168/168	0/7/7/7
2	UDP	A	501	-	-	3/16/32/32	0/2/2/2
3	M2Q	A	502	-	-	11/32/168/168	0/7/7/7

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	503	M2Q	C20-C22	-28.47	1.13	1.56
3	A	502	M2Q	C20-C22	-27.91	1.14	1.56
3	A	502	M2Q	C23-C22	-19.28	1.22	1.56
3	A	503	M2Q	C23-C22	-19.24	1.22	1.56
3	A	503	M2Q	C37-C34	-9.24	1.16	1.33
3	A	502	M2Q	C37-C34	-8.81	1.17	1.33
3	A	502	M2Q	O01-C27	6.79	1.54	1.43
3	A	502	M2Q	C39-C40	6.73	1.65	1.51
3	A	503	M2Q	C39-C40	6.48	1.64	1.51
3	A	502	M2Q	C39-C36	6.46	1.68	1.52
3	A	503	M2Q	O01-C27	6.38	1.54	1.43
3	A	503	M2Q	C39-C36	6.30	1.68	1.52
3	A	503	M2Q	O05-C48	-6.23	1.34	1.44
3	A	502	M2Q	O05-C48	-6.01	1.34	1.44
2	A	501	UDP	PA-O3A	5.30	1.65	1.59
3	A	502	M2Q	O02-C40	5.15	1.52	1.45
3	A	503	M2Q	O02-C40	5.01	1.52	1.45
3	A	503	M2Q	C26-C20	-4.74	1.42	1.54
3	A	502	M2Q	C52-C48	4.35	1.59	1.52
3	A	502	M2Q	C28-C34	3.91	1.59	1.52
3	A	502	M2Q	C33-C24	-3.84	1.47	1.54
3	A	502	M2Q	C38-C40	-3.75	1.48	1.54
3	A	503	M2Q	C28-C34	3.73	1.59	1.52
3	A	502	M2Q	C35-C23	3.50	1.59	1.54
3	A	503	M2Q	C35-C23	3.48	1.59	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	M2Q	C26-C20	-3.44	1.45	1.54
3	A	503	M2Q	C33-C24	-3.36	1.48	1.54
3	A	502	M2Q	C30-C37	3.33	1.57	1.50
3	A	503	M2Q	C38-C40	-3.25	1.49	1.54
3	A	503	M2Q	C30-C37	3.19	1.56	1.50
3	A	503	M2Q	C52-C48	3.17	1.57	1.52
3	A	502	M2Q	O08-C55	2.87	1.49	1.41
3	A	502	M2Q	C25-C21	-2.70	1.49	1.54
3	A	503	M2Q	C25-C21	-2.62	1.49	1.54
3	A	502	M2Q	O04-C47	2.59	1.48	1.41
3	A	502	M2Q	C21-C24	-2.55	1.50	1.54
3	A	503	M2Q	O08-C55	2.38	1.48	1.41
3	A	502	M2Q	C25-C27	-2.34	1.49	1.53
3	A	502	M2Q	O04-C53	2.28	1.49	1.44
3	A	502	M2Q	O10-C56	2.21	1.48	1.43
3	A	503	M2Q	C66-C64	-2.19	1.46	1.52
3	A	502	M2Q	C36-C28	2.06	1.57	1.53
3	A	502	M2Q	O15-C62	2.02	1.47	1.41

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	M2Q	C32-C21-C25	-22.79	79.58	111.08
3	A	502	M2Q	C32-C21-C24	-16.32	80.18	111.72
3	A	502	M2Q	C38-C34-C28	-12.04	105.82	115.90
3	A	503	M2Q	C38-C34-C28	-10.16	107.40	115.90
3	A	502	M2Q	C21-C24-C33	-7.56	110.38	119.28
3	A	503	M2Q	C32-C21-C24	-7.19	97.83	111.72
3	A	502	M2Q	C32-C21-C20	-7.16	100.82	112.36
3	A	502	M2Q	C21-C20-C22	6.53	116.32	109.86
3	A	502	M2Q	C22-C23-C27	5.83	118.56	107.52
3	A	503	M2Q	C31-C20-C22	5.60	118.95	111.49
3	A	503	M2Q	C22-C23-C27	5.36	117.66	107.52
3	A	502	M2Q	C25-C21-C20	4.91	116.70	108.31
3	A	503	M2Q	C21-C20-C22	4.59	114.40	109.86
3	A	502	M2Q	C20-C21-C24	4.34	110.46	100.94
3	A	502	M2Q	C25-C21-C24	4.06	131.22	117.43
3	A	502	M2Q	C47-O02-C40	-4.05	107.72	114.92
3	A	502	M2Q	C21-C25-C27	-4.05	107.40	113.20
3	A	503	M2Q	C26-C20-C21	-3.90	97.19	101.83
3	A	503	M2Q	C30-C37-C34	-3.89	118.23	125.14
3	A	502	M2Q	C20-C22-C23	3.85	124.45	115.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	M2Q	C45-C46-C48	-3.83	109.78	115.45
3	A	503	M2Q	C31-C20-C26	-3.77	97.73	111.63
3	A	503	M2Q	C42-C33-C41	-3.61	104.75	110.34
3	A	503	M2Q	C20-C22-C23	3.59	123.85	115.58
3	A	503	M2Q	C29-C26-C20	-3.47	98.01	104.23
3	A	502	M2Q	C30-C37-C34	-3.47	118.97	125.14
3	A	502	M2Q	C31-C20-C22	3.28	115.86	111.49
3	A	503	M2Q	O02-C40-C39	-3.21	105.94	109.95
3	A	503	M2Q	O03-C46-C48	3.19	110.30	106.43
3	A	502	M2Q	C35-C23-C22	-3.08	105.48	112.81
3	A	503	M2Q	C22-C30-C37	3.04	118.56	112.99
3	A	503	M2Q	C35-C23-C28	-2.93	103.44	110.17
3	A	502	M2Q	O04-C53-C54	2.88	114.89	109.70
3	A	503	M2Q	C47-O02-C40	-2.76	110.01	114.92
3	A	502	M2Q	C31-C20-C21	-2.72	107.98	112.36
3	A	503	M2Q	C30-C22-C23	2.64	113.81	110.55
3	A	502	M2Q	C45-C41-C33	-2.49	108.99	115.37
3	A	503	M2Q	C20-C21-C24	2.47	106.37	100.94
3	A	503	M2Q	C35-C23-C27	-2.46	104.19	108.89
3	A	503	M2Q	C39-C36-C28	-2.45	105.93	112.06
3	A	502	M2Q	C35-C23-C27	-2.40	104.31	108.89
3	A	503	M2Q	C29-C24-C21	-2.39	100.64	103.70
3	A	502	M2Q	O03-C46-C48	2.36	109.29	106.43
3	A	503	M2Q	C26-C20-C22	-2.29	109.66	113.94
3	A	502	M2Q	C26-C20-C22	-2.28	109.68	113.94
3	A	502	M2Q	C31-C20-C26	-2.22	103.46	111.63
3	A	503	M2Q	O04-C53-C54	2.19	113.65	109.70
3	A	502	M2Q	C35-C23-C28	-2.18	105.16	110.17
3	A	502	M2Q	C22-C30-C37	2.17	116.97	112.99
3	A	503	M2Q	C25-C21-C24	2.15	124.75	117.43
3	A	503	M2Q	C36-C28-C23	-2.15	110.77	114.64
3	A	503	M2Q	C39-C40-C38	-2.11	111.69	113.28
3	A	503	M2Q	C25-C21-C20	-2.08	104.76	108.31
3	A	503	M2Q	C47-C49-C50	2.05	114.33	110.01
3	A	502	M2Q	C42-C33-C24	-2.05	109.81	112.88
3	A	503	M2Q	C45-C46-C48	-2.05	112.42	115.45
3	A	502	M2Q	C57-C53-C54	-2.03	108.03	113.02

There are no chirality outliers.



All (24) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	UDP	PB-O3A-PA-O5'
3	A	502	M2Q	O08-C59-C61-O14
3	A	502	M2Q	O15-C65-C67-O19
3	A	503	M2Q	C42-C33-C41-C45
3	A	502	M2Q	C66-C65-C67-O19
3	A	503	M2Q	C24-C33-C41-C45
3	A	502	M2Q	O08-C55-O03-C46
3	A	502	M2Q	C60-C59-C61-O14
3	A	503	M2Q	O04-C53-C57-O11
3	A	503	M2Q	O03-C46-C48-C52
3	A	503	M2Q	C45-C46-C48-C52
3	A	502	M2Q	C21-C24-C33-C42
3	A	502	M2Q	C21-C24-C33-C41
3	A	503	M2Q	C45-C46-C48-O05
3	A	503	M2Q	O15-C65-C67-O19
2	A	501	UDP	O4'-C4'-C5'-O5'
3	A	503	M2Q	C48-C46-O03-C55
3	A	502	M2Q	C29-C24-C33-C42
3	A	502	M2Q	C41-C45-C46-O03
3	A	503	M2Q	O03-C46-C48-O05
3	A	502	M2Q	C29-C24-C33-C41
2	A	501	UDP	C3'-C4'-C5'-O5'
3	A	503	M2Q	C21-C24-C33-C42
3	A	502	M2Q	O03-C46-C48-C51

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	436/459 (94%)	0.33	26 (5%)	27 28	10, 26, 49, 74	5 (1%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	308	SER	4.9
1	A	235	GLY	4.7
1	A	307	THR	4.7
1	A	188	ARG	4.0
1	A	178	TYR	3.8
1	A	457	HIS	3.5
1	A	179	SER	3.5
1	A	351	CYS	3.4
1	A	324	GLU	3.2
1	A	242	ASN	3.2
1	A	249	GLY	2.8
1	A	165	PHE	2.8
1	A	309	GLY	2.7
1	A	248	GLU	2.7
1	A	160	HIS	2.6
1	A	301	PHE	2.5
1	A	189	THR	2.4
1	A	374	ASP	2.4
1	A	456	HIS	2.4
1	A	316	LYS	2.3
1	A	402	ASP	2.2
1	A	312	ASP	2.2
1	A	454	HIS	2.1
1	A	304	GLY	2.0
1	A	303	GLN	2.0
1	A	305	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	M2Q	A	502	67/67	0.84	0.15	17,30,85,89	0
3	M2Q	A	503	67/67	0.85	0.12	25,35,43,47	0
2	UDP	A	501	25/25	0.97	0.07	19,23,26,31	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around M2Q A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around M2Q A 503:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UDP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.