



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 10:40 PM UTC

PDB ID : 5J7X / pdb_00005j7x
Title : Baeyer-Villiger monooxygenase BVMOAFL838 from *Aspergillus flavus*
Authors : Ferroni, F.M.; Tolmie, C.; Smit, M.S.; Opperman, D.J.
Deposited on : 2016-04-07
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	549	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	TAM	A	603	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4659 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dimethylaniline monooxygenase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	527	4229	2712	713	790	14	0	1	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	544	HIS	-	expression tag	UNP B8N653
A	545	HIS	-	expression tag	UNP B8N653
A	546	HIS	-	expression tag	UNP B8N653
A	547	HIS	-	expression tag	UNP B8N653
A	548	HIS	-	expression tag	UNP B8N653
A	549	HIS	-	expression tag	UNP B8N653

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is TRIS(HYDROXYETHYL)AMINOMETHANE (CCD ID: TAM) (formula: C₇H₁₇NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			11	7	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	361	Total	O	0	0
			361	361		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dimethylaniline monooxygenase, putative



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	109.67Å 177.51Å 76.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.40 – 1.90 32.40 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (32.40-1.90) 99.5 (32.40-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.157 , 0.204 0.168 , 0.211	Depositor DCC
R_{free} test set	2981 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	23.8	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4659	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TAM, FAD, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.98	89/4346 (2.0%)	1.53	35/5906 (0.6%)

All (89) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	437	SER	CA-C	9.69	1.63	1.53
1	A	132	THR	N-CA	-9.26	1.34	1.46
1	A	439	GLN	C-O	9.14	1.36	1.23
1	A	389	GLY	CA-C	8.64	1.65	1.52
1	A	171	ALA	N-CA	8.33	1.56	1.46
1	A	170	SER	C-O	-8.25	1.13	1.24
1	A	269	GLU	CD-OE1	7.95	1.40	1.25
1	A	214	ARG	C-O	-7.93	1.14	1.23
1	A	474	THR	N-CA	7.75	1.55	1.46
1	A	405	VAL	C-O	-7.49	1.13	1.24
1	A	147	GLY	CA-C	7.36	1.61	1.51
1	A	504	MET	N-CA	7.23	1.54	1.46
1	A	26	TYR	C-O	7.22	1.32	1.24
1	A	429	PHE	N-CA	7.13	1.54	1.46
1	A	441	PRO	C-O	-7.08	1.14	1.24
1	A	516	LEU	N-CA	-7.05	1.37	1.46
1	A	278	TYR	CZ-OH	7.04	1.52	1.38
1	A	150	SER	CA-C	-7.02	1.43	1.52
1	A	64	CYS	C-O	-6.88	1.16	1.24
1	A	412	GLU	N-CA	-6.85	1.37	1.46
1	A	170	SER	CA-C	6.83	1.62	1.52
1	A	490	ILE	C-O	6.80	1.31	1.24
1	A	193	SER	CA-C	-6.76	1.44	1.52
1	A	61	ARG	CZ-NH2	-6.73	1.24	1.33
1	A	137	ALA	N-CA	6.71	1.54	1.45
1	A	533	LEU	N-CA	-6.67	1.37	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	525	TYR	C-O	6.66	1.31	1.24
1	A	206	ALA	N-CA	-6.56	1.37	1.45
1	A	171	ALA	CA-C	6.52	1.61	1.52
1	A	538	PHE	CA-C	6.52	1.60	1.52
1	A	114	ARG	CZ-NH2	-6.44	1.25	1.33
1	A	199	PHE	C-O	-6.44	1.16	1.24
1	A	31	LEU	C-O	6.39	1.31	1.24
1	A	526	ARG	CZ-NH1	6.37	1.41	1.32
1	A	79	SER	CA-C	-6.35	1.44	1.52
1	A	130	TRP	CA-C	6.34	1.60	1.52
1	A	459	THR	N-CA	-6.33	1.38	1.46
1	A	498	LEU	CA-C	-6.19	1.43	1.52
1	A	160	LEU	CA-C	-6.17	1.44	1.52
1	A	278	TYR	CG-CD2	-6.15	1.26	1.39
1	A	425	ALA	C-O	6.12	1.31	1.23
1	A	213	GLN	C-O	-6.07	1.17	1.24
1	A	143	LEU	CA-CB	-6.05	1.44	1.53
1	A	66	VAL	N-CA	-6.03	1.40	1.47
1	A	515	SER	C-O	-6.03	1.16	1.23
1	A	486	GLU	CD-OE2	6.02	1.36	1.25
1	A	446	ASN	CA-C	5.97	1.60	1.52
1	A	219	CYS	N-CA	5.95	1.53	1.46
1	A	413	ARG	CZ-NH2	-5.92	1.25	1.33
1	A	97	ASP	N-CA	5.88	1.53	1.46
1	A	398	LEU	CA-C	-5.79	1.44	1.52
1	A	418	MET	C-O	-5.78	1.16	1.24
1	A	54	ASN	N-CA	-5.75	1.39	1.46
1	A	96	PHE	C-O	5.63	1.30	1.24
1	A	255	TYR	CA-C	5.63	1.59	1.52
1	A	153	TYR	C-O	-5.62	1.17	1.24
1	A	254	GLU	C-O	-5.62	1.16	1.24
1	A	218	LEU	N-CA	-5.62	1.39	1.46
1	A	162	SER	CB-OG	5.58	1.53	1.42
1	A	464	LYS	N-CA	-5.54	1.39	1.46
1	A	484	ASN	C-O	-5.54	1.17	1.24
1	A	91	GLU	N-CA	-5.52	1.39	1.46
1	A	331	TYR	CZ-OH	5.50	1.49	1.38
1	A	34	GLU	N-CA	-5.49	1.39	1.46
1	A	78	GLN	CA-C	-5.48	1.45	1.52
1	A	402	LEU	CA-C	5.48	1.59	1.52
1	A	439	GLN	CA-C	-5.45	1.45	1.53
1	A	394	THR	CA-C	-5.43	1.45	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	213	GLN	CA-C	5.38	1.58	1.52
1	A	208	GLU	C-O	-5.38	1.17	1.23
1	A	148	PHE	C-O	-5.38	1.18	1.24
1	A	449	THR	CA-C	5.38	1.59	1.52
1	A	461	LEU	C-O	5.37	1.30	1.24
1	A	227	LEU	CA-C	5.37	1.59	1.52
1	A	269	GLU	CA-C	5.32	1.59	1.52
1	A	32	ARG	CZ-NH1	5.32	1.40	1.32
1	A	533	LEU	CA-C	5.32	1.60	1.52
1	A	97	ASP	C-O	-5.28	1.18	1.24
1	A	75	GLU	CD-OE2	5.24	1.35	1.25
1	A	220	LEU	CA-C	5.21	1.60	1.53
1	A	169	HIS	C-O	-5.19	1.18	1.24
1	A	215	THR	CA-C	-5.14	1.48	1.52
1	A	61	ARG	CA-C	-5.11	1.46	1.52
1	A	90	LYS	C-O	-5.11	1.18	1.24
1	A	261	ASN	CG-ND2	-5.07	1.22	1.33
1	A	350	LYS	CA-C	-5.06	1.46	1.52
1	A	360	SER	CA-CB	5.05	1.61	1.53
1	A	340	LEU	N-CA	-5.05	1.40	1.45
1	A	94	SER	CA-CB	5.00	1.61	1.53

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	504	MET	CG-SD-CE	-12.76	72.82	100.90
1	A	175	GLU	N-CA-C	-9.06	95.47	108.96
1	A	370	GLY	N-CA-C	7.46	121.32	110.80
1	A	392	ALA	N-CA-C	-7.27	101.47	110.41
1	A	215	THR	CB-CA-C	-7.17	97.68	109.15
1	A	526	ARG	NE-CZ-NH2	-7.00	112.90	119.20
1	A	328	GLU	CA-C-N	-6.83	114.74	119.66
1	A	328	GLU	C-N-CA	-6.83	114.74	119.66
1	A	545	HIS	CA-C-O	-6.43	109.86	120.80
1	A	160	LEU	CB-CA-C	-6.34	98.91	110.63
1	A	269	GLU	CB-CA-C	-6.32	100.96	110.88
1	A	23	SER	N-CA-C	6.31	117.82	111.07
1	A	321	ARG	N-CA-CB	-6.30	100.86	110.12
1	A	30	LYS	CD-CE-NZ	6.15	131.59	111.90
1	A	394	THR	N-CA-C	6.15	117.98	111.28
1	A	360	SER	CB-CA-C	-6.07	101.31	110.90
1	A	526	ARG	CB-CA-C	-6.04	101.15	110.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	390	PHE	CA-CB-CG	-6.03	107.77	113.80
1	A	390	PHE	N-CA-C	5.79	119.52	110.32
1	A	267	GLU	CA-CB-CG	-5.72	102.65	114.10
1	A	201	GLU	CB-CA-C	-5.71	101.92	110.88
1	A	229	ALA	CA-C-O	-5.68	111.14	120.80
1	A	461	LEU	CD1-CG-CD2	-5.68	98.30	110.80
1	A	511	LYS	N-CA-C	-5.62	99.77	109.15
1	A	476	THR	N-CA-CB	5.59	116.80	110.03
1	A	394	THR	CA-CB-OG1	-5.52	101.33	109.60
1	A	429	PHE	CA-C-N	-5.51	114.91	120.31
1	A	429	PHE	C-N-CA	-5.51	114.91	120.31
1	A	412	GLU	N-CA-CB	-5.37	102.20	110.20
1	A	269	GLU	CG-CD-OE2	-5.34	106.13	118.40
1	A	442	THR	N-CA-C	5.28	122.05	110.80
1	A	418	MET	N-CA-C	-5.25	101.13	109.59
1	A	200	GLN	CB-CA-C	-5.09	102.66	110.81
1	A	54	ASN	N-CA-C	5.08	117.93	111.69
1	A	466	GLU	N-CA-C	-5.07	105.83	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4229	0	4089	28	0
2	A	53	0	31	0	0
3	A	5	0	0	0	0
4	A	11	0	15	2	0
5	A	361	0	0	1	2
All	All	4659	0	4135	28	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:GLN:HA	1:A:78:GLN:OE1	1.76	0.82
1:A:393:VAL:HG11	1:A:438:LEU:HB3	1.65	0.77
1:A:214:ARG:HB3	4:A:603:TAM:H62	1.76	0.67
1:A:500:ASP:O	1:A:504:MET:HG2	1.97	0.65
1:A:175:GLU:O	1:A:176:ALA:CB	2.48	0.61
1:A:78:GLN:OE1	1:A:78:GLN:CA	2.50	0.60
1:A:75:GLU:H	1:A:75:GLU:CD	2.15	0.55
1:A:414:TRP:CH2	1:A:438:LEU:HD21	2.42	0.55
1:A:79:SER:HB2	5:A:877:HOH:O	2.09	0.53
1:A:175:GLU:O	1:A:176:ALA:HB3	2.09	0.52
1:A:185:LYS:HE3	1:A:207:GLU:OE2	2.11	0.50
1:A:461:LEU:C	1:A:461:LEU:HD23	2.37	0.49
1:A:453:LEU:C	1:A:453:LEU:HD23	2.37	0.49
1:A:367:THR:HB	1:A:368:PRO:CD	2.43	0.49
1:A:457:TRP:CD1	1:A:529:CYS:HB3	2.48	0.48
1:A:504:MET:HE3	1:A:511:LYS:HE2	1.96	0.47
1:A:160:LEU:HD22	1:A:167:ILE:CD1	2.45	0.47
1:A:393:VAL:CG1	1:A:438:LEU:HB3	2.41	0.45
1:A:386:VAL:HG12	1:A:388:THR:HG23	1.99	0.45
1:A:74:PRO:O	1:A:78:GLN:HG2	2.18	0.43
1:A:367:THR:HB	1:A:368:PRO:HD2	1.99	0.43
1:A:245:GLU:HG3	1:A:246:CYS:N	2.34	0.42
1:A:352:ASN:OD1	1:A:352:ASN:N	2.42	0.42
1:A:17:ILE:HB	1:A:41:ILE:HG12	2.00	0.42
1:A:394:THR:O	1:A:398:LEU:HG	2.20	0.41
1:A:148:PHE:CE2	1:A:393:VAL:HB	2.56	0.41
1:A:214:ARG:HE	4:A:603:TAM:H52	1.86	0.41
1:A:166:THR:HG21	1:A:168:TYR:CZ	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:716:HOH:O	5:A:716:HOH:O[2_775]	0.94	1.26
5:A:1037:HOH:O	5:A:1037:HOH:O[3_756]	1.23	0.97

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	524/549 (95%)	509 (97%)	13 (2%)	2 (0%)	30	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	176	ALA
1	A	442	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	447/464 (96%)	438 (98%)	9 (2%)	48	46

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	GLU
1	A	78	GLN
1	A	158	LYS
1	A	175	GLU
1	A	205	GLU
1	A	355	ILE
1	A	394	THR
1	A	438	LEU
1	A	453	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	ASN
1	A	228	HIS
1	A	406	ASN
1	A	439	GLN
1	A	484	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	602	-	4,4,4	1.27	1 (25%)	6,6,6	0.89	0
2	FAD	A	601	-	58,58,58	1.85	16 (27%)	85,89,89	1.88	23 (27%)
4	TAM	A	603	-	10,10,10	2.10	3 (30%)	12,12,12	7.69	10 (83%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	601	-	-	5/34/50/50	0/6/6/6
4	TAM	A	603	-	-	7/12/12/12	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	FAD	C1'-C2'	5.89	1.60	1.52
2	A	601	FAD	C9A-C5X	5.00	1.49	1.41
4	A	603	TAM	C3-C	-4.51	1.47	1.53
2	A	601	FAD	C5'-C4'	4.08	1.57	1.51
2	A	601	FAD	C7M-C7	3.55	1.57	1.51
2	A	601	FAD	P-O3P	-3.45	1.55	1.59
2	A	601	FAD	C4-N3	-3.24	1.32	1.38
4	A	603	TAM	C3-C6	-3.20	1.46	1.52
2	A	601	FAD	C5X-N5	-2.65	1.34	1.39
2	A	601	FAD	C5A-N7A	-2.62	1.34	1.39
2	A	601	FAD	C6-C5X	-2.58	1.36	1.40
2	A	601	FAD	C9-C8	-2.51	1.36	1.39
4	A	603	TAM	C1-C	-2.49	1.50	1.53
2	A	601	FAD	O4-C4	2.46	1.28	1.23
2	A	601	FAD	C4A-N9A	-2.40	1.32	1.37
2	A	601	FAD	C2A-N3A	2.40	1.38	1.33
2	A	601	FAD	C6-C7	2.29	1.42	1.39
2	A	601	FAD	PA-O1A	-2.20	1.43	1.50
2	A	601	FAD	C8A-N7A	2.14	1.35	1.31
3	A	602	SO4	O2-S	2.11	1.57	1.44

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	603	TAM	C3-C-N	-16.22	67.51	108.22
4	A	603	TAM	C4-C1-C	-12.85	100.82	115.97
4	A	603	TAM	C5-C2-C	-9.45	104.83	115.97
4	A	603	TAM	C3-C-C1	6.70	122.32	110.50
4	A	603	TAM	O6-C6-C3	-5.84	94.93	111.33
4	A	603	TAM	C6-C3-C	-5.41	109.59	115.97
2	A	601	FAD	N9A-C8A-N7A	-5.08	106.72	113.94
4	A	603	TAM	C1-C-N	-5.01	95.64	108.22
4	A	603	TAM	C2-C-C1	5.01	119.34	110.50
2	A	601	FAD	C5A-C4A-N3A	-4.51	120.50	126.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FAD	C4A-N9A-C8A	4.37	110.33	105.74
4	A	603	TAM	C2-C-N	-4.19	97.69	108.22
4	A	603	TAM	C3-C-C2	3.95	117.47	110.50
2	A	601	FAD	O4-C4-C4X	-3.90	116.25	126.53
2	A	601	FAD	N3A-C4A-N9A	3.60	133.29	127.17
2	A	601	FAD	N3A-C2A-N1A	-3.51	123.27	128.58
2	A	601	FAD	C10-N1-C2	3.42	124.26	116.85
2	A	601	FAD	C5A-N7A-C8A	3.29	108.61	103.45
2	A	601	FAD	C4-C4X-N5	3.28	122.73	118.21
2	A	601	FAD	C6-C5X-N5	3.18	123.73	118.44
2	A	601	FAD	O2P-P-O1P	3.14	127.06	112.44
2	A	601	FAD	C4'-C3'-C2'	2.98	118.52	113.57
2	A	601	FAD	C4X-C4-N3	2.97	120.82	113.25
2	A	601	FAD	O3'-C3'-C4'	-2.77	102.65	108.93
2	A	601	FAD	C4A-N9A-C1B	-2.63	120.49	126.63
2	A	601	FAD	N6A-C6A-N1A	2.61	124.20	118.38
2	A	601	FAD	C4-C4X-C10	-2.47	112.70	116.93
2	A	601	FAD	C4A-C5A-N7A	-2.38	107.86	110.58
2	A	601	FAD	C9A-C5X-N5	-2.35	119.96	122.45
2	A	601	FAD	O2'-C2'-C3'	-2.34	103.77	109.25
2	A	601	FAD	C6-C5X-C9A	-2.30	115.89	119.05
2	A	601	FAD	O4B-C1B-C2B	-2.14	102.03	106.62
2	A	601	FAD	O2'-C2'-C1'	-2.05	101.77	110.20

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	FAD	N10-C1'-C2'-O2'
2	A	601	FAD	N10-C1'-C2'-C3'
4	A	603	TAM	C2-C-C1-C4
4	A	603	TAM	N-C-C1-C4
4	A	603	TAM	C1-C-C2-C5
4	A	603	TAM	C3-C-C2-C5
4	A	603	TAM	C1-C-C3-C6
4	A	603	TAM	C-C3-C6-O6
2	A	601	FAD	C2'-C3'-C4'-C5'
2	A	601	FAD	O3'-C3'-C4'-C5'
4	A	603	TAM	C-C2-C5-O5
2	A	601	FAD	O4B-C4B-C5B-O5B

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	603	TAM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	527/549 (95%)	-0.23	13 (2%) 58 62	13, 26, 45, 84	1 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	176	ALA	3.9
1	A	171	ALA	3.6
1	A	393	VAL	3.4
1	A	229	ALA	3.2
1	A	388	THR	3.2
1	A	9	GLY	2.9
1	A	240	ALA	2.6
1	A	389	GLY	2.6
1	A	394	THR	2.6
1	A	418	MET	2.3
1	A	243	LEU	2.1
1	A	170	SER	2.0
1	A	78	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	TAM	A	603	11/11	0.87	0.14	35,46,52,56	0
3	SO4	A	602	5/5	0.96	0.21	39,40,41,42	0
2	FAD	A	601	53/53	0.99	0.04	13,16,23,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.