



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 04:09 PM UTC

PDB ID : 5J8R / pdb_00005j8r
Title : Crystal Structure of the Catalytic Domain of Human Protein Tyrosine Phosphatase non-receptor Type 12 - K61R mutant
Authors : Li, H.; Yang, F.; Xu, Y.F.; Wang, W.J.; Xiao, P.; Yu, X.; Sun, J.P.
Deposited on : 2016-04-08
Resolution : 2.04 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	
1	B	311	
1	C	311	
1	D	311	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10076 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	S	0	1	0
			2477	1576	434	451	16			
1	B	293	Total	C	N	O	S	0	1	0
			2453	1563	429	445	16			
1	C	286	Total	C	N	O	S	0	0	0
			2396	1531	419	430	16			
1	D	284	Total	C	N	O	S	0	0	0
			2373	1518	412	427	16			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP Q05209
A	-4	HIS	-	expression tag	UNP Q05209
A	-3	HIS	-	expression tag	UNP Q05209
A	-2	HIS	-	expression tag	UNP Q05209
A	-1	HIS	-	expression tag	UNP Q05209
A	0	HIS	-	expression tag	UNP Q05209
A	61	ARG	LYS	engineered mutation	UNP Q05209
B	-5	HIS	-	expression tag	UNP Q05209
B	-4	HIS	-	expression tag	UNP Q05209
B	-3	HIS	-	expression tag	UNP Q05209
B	-2	HIS	-	expression tag	UNP Q05209
B	-1	HIS	-	expression tag	UNP Q05209
B	0	HIS	-	expression tag	UNP Q05209
B	61	ARG	LYS	engineered mutation	UNP Q05209
C	-5	HIS	-	expression tag	UNP Q05209
C	-4	HIS	-	expression tag	UNP Q05209
C	-3	HIS	-	expression tag	UNP Q05209
C	-2	HIS	-	expression tag	UNP Q05209
C	-1	HIS	-	expression tag	UNP Q05209
C	0	HIS	-	expression tag	UNP Q05209
C	61	ARG	LYS	engineered mutation	UNP Q05209

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-5	HIS	-	expression tag	UNP Q05209
D	-4	HIS	-	expression tag	UNP Q05209
D	-3	HIS	-	expression tag	UNP Q05209
D	-2	HIS	-	expression tag	UNP Q05209
D	-1	HIS	-	expression tag	UNP Q05209
D	0	HIS	-	expression tag	UNP Q05209
D	61	ARG	LYS	engineered mutation	UNP Q05209


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	130	Total 130	O 130	0	0
2	B	142	Total 142	O 142	0	0
2	C	51	Total 51	O 51	0	0
2	D	54	Total 54	O 54	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tyrosine-protein phosphatase non-receptor type 12




• Molecule 1: Tyrosine-protein phosphatase non-receptor type 12




• Molecule 1: Tyrosine-protein phosphatase non-receptor type 12





- Molecule 1: Tyrosine-protein phosphatase non-receptor type 12



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	129.94Å 129.94Å 278.76Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.79 – 2.04 37.79 – 2.04	Depositor EDS
% Data completeness (in resolution range)	90.3 (37.79-2.04) 90.4 (37.79-2.04)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.84 (at 2.05Å)	Xtriage
Refinement program	PHENIX 1.6.2_432	Depositor
R, R_{free}	0.239 , 0.275 0.233 , 0.266	Depositor DCC
R_{free} test set	5268 reflections (4.73%)	wwPDB-VP
Wilson B-factor (Å ²)	30.2	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.457 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10076	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	3/2536 (0.1%)	0.85	0/3422
1	B	0.55	0/2511	0.88	0/3388
1	C	0.59	0/2452	0.86	3/3307 (0.1%)
1	D	0.53	0/2427	0.81	0/3272
All	All	0.59	3/9926 (0.0%)	0.85	3/13389 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	72	HIS	CG-ND1	-7.32	1.30	1.38
1	A	71	ASP	C-O	-6.51	1.16	1.24
1	A	72	HIS	CD2-NE2	-5.97	1.31	1.37

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	199	ASP	N-CA-C	7.57	120.89	110.06
1	C	185	GLU	N-CA-C	-7.06	98.91	109.86
1	C	75	VAL	CB-CA-C	-5.17	105.64	111.23

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	184	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2477	0	2420	89	0
1	B	2453	0	2404	57	0
1	C	2396	0	2358	131	0
1	D	2373	0	2326	140	0
2	A	130	0	0	10	0
2	B	142	0	0	8	0
2	C	51	0	0	8	0
2	D	54	0	0	10	0
All	All	10076	0	9508	414	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 414 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:165:GLU:OE2	1:C:179:LEU:HD11	1.18	1.30
1:C:13:ARG:HH21	1:C:17:MET:HG3	1.11	1.12
1:C:165:GLU:OE2	1:C:179:LEU:CD1	2.00	1.07
1:B:155:ILE:C	1:B:161:LYS:HZ3	1.62	1.05
1:B:299:GLN:HA	1:B:299:GLN:OE1	1.60	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/311 (94%)	278 (95%)	12 (4%)	3 (1%)	12	6
1	B	290/311 (93%)	279 (96%)	8 (3%)	3 (1%)	12	6
1	C	280/311 (90%)	245 (88%)	27 (10%)	8 (3%)	3	0
1	D	274/311 (88%)	232 (85%)	36 (13%)	6 (2%)	5	1
All	All	1137/1244 (91%)	1034 (91%)	83 (7%)	20 (2%)	6	2

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	199	ASP
1	C	82	PRO
1	C	171	THR
1	D	171	THR
1	A	171	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/282 (95%)	245 (91%)	23 (9%)	10	4
1	B	265/282 (94%)	243 (92%)	22 (8%)	10	5
1	C	258/282 (92%)	229 (89%)	29 (11%)	6	1
1	D	255/282 (90%)	222 (87%)	33 (13%)	4	1
All	All	1046/1128 (93%)	939 (90%)	107 (10%)	7	2

5 of 107 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	76	LYS
1	C	235	CYS
1	D	199	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	84	GLN
1	C	167	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	299	GLN
1	D	27	ASN
1	D	183	GLN
1	B	22	HIS
1	B	15	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	296/311 (95%)	-1.32	0 100 100	16, 33, 53, 69	1 (0%)
1	B	293/311 (94%)	-1.33	0 100 100	15, 33, 52, 63	1 (0%)
1	C	286/311 (91%)	-0.97	0 100 100	30, 54, 74, 86	0
1	D	284/311 (91%)	-0.96	0 100 100	30, 54, 74, 84	0
All	All	1159/1244 (93%)	-1.15	0 100 100	15, 42, 71, 86	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.