



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2026 – 01:33 PM UTC

PDB ID : 6J8Y / pdb_00006j8y
Title : Crystal structure of the human RAD9-HUS1-RAD1-RHINO complex
Authors : Hara, K.; Iida, N.; Sakurai, H.; Hashimoto, H.
Deposited on : 2019-01-21
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	270	
2	B	286	
3	C	282	
4	D	20	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6267 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell cycle checkpoint control protein RAD9A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			1988	1270	343	359	16			

- Molecule 2 is a protein called Checkpoint protein HUS1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	264	Total	C	N	O	S	0	0	0
			2120	1356	366	384	14			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-5	MET	-	initiating methionine	UNP O60921
B	-4	HIS	-	expression tag	UNP O60921
B	-3	HIS	-	expression tag	UNP O60921
B	-2	HIS	-	expression tag	UNP O60921
B	-1	HIS	-	expression tag	UNP O60921
B	0	HIS	-	expression tag	UNP O60921
B	1	HIS	-	expression tag	UNP O60921

- Molecule 3 is a protein called Cell cycle checkpoint protein RAD1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	255	Total	C	N	O	S	0	0	0
			2012	1279	332	384	17			

- Molecule 4 is a protein called RAD9, HUS1, RAD1-interacting nuclear orphan protein 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	10	Total	C	N	O	0	0	0
			79	51	11	17			


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	38	Total 38	O 38	0	0
5	B	18	Total 18	O 18	0	0
5	C	10	Total 10	O 10	0	0
5	D	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cell cycle checkpoint control protein RAD9A




- Molecule 2: Checkpoint protein HUS1



- Molecule 3: Cell cycle checkpoint protein RAD1



- Molecule 4: RAD9, HUS1, RAD1-interacting nuclear orphan protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.94Å 136.19Å 154.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.80 – 2.40 19.80 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (19.80-2.40) 99.7 (19.80-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.41Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.218 , 0.258 0.221 , 0.261	Depositor DCC
R_{free} test set	2248 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	50.8	Xtriage
Anisotropy	0.322	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6267	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	0/2022	0.69	0/2724
2	B	0.26	0/2159	0.71	0/2919
3	C	0.28	0/2047	0.72	1/2766 (0.0%)
4	D	0.29	0/82	0.60	0/113
All	All	0.27	0/6310	0.71	1/8522 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	114	GLY	N-CA-C	5.64	118.24	110.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1988	0	2030	22	0
2	B	2120	0	2176	43	0
3	C	2012	0	1997	42	0
4	D	79	0	64	1	0
5	A	38	0	0	1	0
5	B	18	0	0	0	0
5	C	10	0	0	0	0
5	D	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6267	0	6267	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (104) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:197:GLU:HG3	2:B:198:LEU:HG	1.53	0.91
3:C:158:LEU:HB3	3:C:163:LEU:HD11	1.66	0.76
1:A:6:THR:HG22	1:A:96:LYS:HG2	1.68	0.75
3:C:15:TYR:HA	3:C:113:GLN:HB3	1.71	0.72
3:C:16:SER:H	3:C:113:GLN:HA	1.56	0.70
1:A:154:GLU:OE1	2:B:127:ARG:NH1	2.24	0.65
3:C:57:LYS:HA	3:C:143:LEU:HD11	1.79	0.65
1:A:174:ARG:NH1	1:A:207:LEU:O	2.30	0.64
2:B:33:ARG:HB3	2:B:40:ASN:HB2	1.79	0.63
2:B:61:GLU:H	2:B:61:GLU:CD	2.07	0.62
1:A:72:ARG:HD2	1:A:131:SER:HB2	1.81	0.62
2:B:187:ASP:HB2	2:B:225:GLU:HG2	1.82	0.61
2:B:43:LEU:HB2	2:B:54:MET:HB3	1.83	0.61
3:C:121:LEU:HB2	3:C:132:CYS:HB2	1.82	0.61
2:B:17:THR:O	2:B:21:ASN:ND2	2.34	0.60
1:A:123:ASN:HB2	3:C:198:SER:HB3	1.82	0.60
2:B:60:GLN:HG3	2:B:67:PHE:CG	2.37	0.59
3:C:44:ALA:HB3	3:C:80:VAL:HG13	1.84	0.59
1:A:30:GLU:HG3	1:A:72:ARG:HG3	1.85	0.58
3:C:163:LEU:HB3	3:C:167:PHE:CE2	2.40	0.57
2:B:-2:HIS:CD2	2:B:-2:HIS:N	2.73	0.57
2:B:173:MET:HE2	2:B:194:ILE:HG23	1.87	0.56
3:C:35:HIS:NE2	3:C:87:THR:OG1	2.35	0.56
3:C:151:ASN:N	3:C:151:ASN:OD1	2.38	0.56
3:C:24:VAL:HG11	3:C:123:LEU:HD23	1.88	0.55
2:B:4:ARG:NH1	2:B:66:GLU:OE1	2.37	0.54
3:C:167:PHE:HA	3:C:170:LEU:HD13	1.90	0.54
2:B:24:ALA:HA	2:B:85:SER:HB3	1.89	0.53
3:C:163:LEU:HB3	3:C:167:PHE:HE2	1.72	0.53
3:C:164:ARG:NH1	3:C:235:LEU:O	2.41	0.53
1:A:4:LEU:HB2	1:A:60:GLN:HB3	1.91	0.52
3:C:165:GLU:CD	3:C:165:GLU:H	2.17	0.52
3:C:24:VAL:HG12	3:C:108:LEU:HB2	1.92	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:60:GLN:HG3	2:B:67:PHE:CD2	2.46	0.51
2:B:250:LYS:HB3	2:B:266:LEU:HB3	1.92	0.51
2:B:81:LEU:HD23	2:B:135:ILE:HD13	1.92	0.51
1:A:42:ASN:HB2	1:A:47:ALA:H	1.75	0.50
2:B:14:ASN:O	2:B:18:ARG:HG3	2.11	0.50
2:B:23:ILE:HG12	2:B:54:MET:HE1	1.94	0.50
2:B:171:GLU:O	2:B:174:LYS:HE3	2.12	0.50
3:C:205:LYS:HA	3:C:210:MET:HG3	1.94	0.50
2:B:186:LEU:HB2	2:B:225:GLU:HG3	1.94	0.49
3:C:15:TYR:CE2	3:C:111:CYS:HB3	2.47	0.49
2:B:61:GLU:OE1	2:B:61:GLU:N	2.40	0.49
2:B:225:GLU:N	2:B:225:GLU:OE1	2.45	0.49
2:B:185:ASN:HB3	2:B:187:ASP:H	1.78	0.49
1:A:155:ALA:HA	1:A:197:MET:HE3	1.95	0.48
1:A:242:ALA:HB3	1:A:245:ARG:HD2	1.96	0.48
2:B:172:LYS:HB3	3:C:95:ILE:HD12	1.96	0.48
3:C:145:PHE:O	3:C:147:PHE:N	2.45	0.47
1:A:158:PRO:HG2	1:A:197:MET:HE2	1.97	0.47
2:B:233:ASP:OD2	2:B:235:ARG:NH1	2.47	0.47
3:C:151:ASN:HB2	3:C:247:ASN:ND2	2.29	0.47
1:A:9:ASN:HA	1:A:12:VAL:HG13	1.97	0.47
3:C:151:ASN:HB2	3:C:247:ASN:HD21	1.80	0.47
2:B:55:TRP:CE2	2:B:148:GLU:HB3	2.49	0.47
1:A:165:GLU:HB2	1:A:181:TYR:HB3	1.97	0.47
3:C:59:VAL:HG21	3:C:227:LEU:HB3	1.96	0.46
3:C:34:ILE:O	3:C:53:VAL:HG21	2.15	0.46
3:C:176:VAL:HG21	3:C:222:ARG:NH2	2.30	0.46
2:B:28:LYS:HD3	2:B:28:LYS:HA	1.74	0.46
1:A:69:ASP:HB3	1:A:70:LEU:H	1.55	0.46
3:C:179:ILE:HG12	3:C:190:LEU:HD12	1.98	0.46
2:B:179:HIS:HB3	2:B:231:HIS:NE2	2.31	0.45
3:C:238:SER:HB3	3:C:255:TYR:HB3	1.98	0.45
3:C:111:CYS:HB2	3:C:120:MET:HB2	1.97	0.45
3:C:165:GLU:CD	3:C:165:GLU:N	2.75	0.45
2:B:102:LYS:HB2	2:B:116:SER:HB2	1.99	0.44
2:B:249:THR:HG22	2:B:267:HIS:HA	2.00	0.44
3:C:159:GLN:O	3:C:163:LEU:HG	2.16	0.44
2:B:105:LEU:HD12	2:B:105:LEU:HA	1.86	0.44
3:C:164:ARG:HD3	3:C:236:VAL:HA	2.00	0.44
1:A:90:LEU:HD23	1:A:94:VAL:HG21	1.99	0.44
2:B:82:GLU:HB3	2:B:136:LYS:HB3	1.98	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:21:LEU:HD23	3:C:70:PHE:HA	1.99	0.44
2:B:166:MET:HE2	2:B:241:LEU:HD11	1.99	0.43
2:B:-2:HIS:CD2	2:B:-2:HIS:H3	2.36	0.43
1:A:229:ALA:HB2	1:A:251:ILE:HG21	2.00	0.43
4:D:57:VAL:O	4:D:59:PRO:HD3	2.18	0.43
3:C:166:ALA:HB1	3:C:201:LEU:HD21	2.00	0.43
2:B:69:MET:HE2	2:B:69:MET:HB2	1.95	0.42
2:B:15:HIS:CE1	2:B:267:HIS:HD2	2.38	0.42
3:C:254:GLN:HG2	3:C:268:GLU:HG2	2.01	0.42
1:A:100:SER:OG	1:A:101:LEU:N	2.52	0.42
3:C:57:LYS:HG3	3:C:143:LEU:HD21	2.01	0.42
1:A:225:LEU:HD13	1:A:260:PHE:CE1	2.54	0.42
3:C:24:VAL:CG1	3:C:108:LEU:HB2	2.49	0.42
3:C:59:VAL:CG2	3:C:227:LEU:HB3	2.50	0.42
3:C:81:THR:HG22	3:C:139:PRO:HG2	2.02	0.42
1:A:78:LYS:HA	1:A:78:LYS:HD2	1.79	0.41
1:A:83:VAL:HG11	1:A:110:VAL:HG11	2.01	0.41
2:B:26:LEU:HD12	2:B:54:MET:HE2	2.02	0.41
3:C:116:GLY:C	3:C:117:TYR:HD1	2.28	0.41
3:C:161:GLU:C	3:C:163:LEU:H	2.28	0.41
3:C:242:SER:HB3	3:C:254:GLN:HB2	2.02	0.41
2:B:33:ARG:NH1	2:B:40:ASN:OD1	2.53	0.41
2:B:161:PRO:HG2	2:B:166:MET:HG3	2.03	0.41
1:A:194:VAL:HG22	2:B:132:ASP:HB2	2.02	0.41
2:B:279:LEU:O	2:B:280:SER:OG	2.32	0.41
3:C:187:TYR:HD2	3:C:204:PRO:HA	1.86	0.41
1:A:174:ARG:NH2	5:A:302:HOH:O	2.28	0.40
2:B:252:LEU:HD23	2:B:252:LEU:HA	1.96	0.40
2:B:43:LEU:HD22	2:B:54:MET:HE3	2.04	0.40
2:B:174:LYS:HE3	2:B:174:LYS:HB3	1.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/270 (92%)	246 (99%)	3 (1%)	0	100	100
2	B	256/286 (90%)	243 (95%)	12 (5%)	1 (0%)	30	43
3	C	249/282 (88%)	240 (96%)	9 (4%)	0	100	100
4	D	8/20 (40%)	6 (75%)	2 (25%)	0	100	100
All	All	762/858 (89%)	735 (96%)	26 (3%)	1 (0%)	48	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	268	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/229 (96%)	212 (97%)	7 (3%)	34	56
2	B	245/263 (93%)	235 (96%)	10 (4%)	27	46
3	C	230/256 (90%)	218 (95%)	12 (5%)	21	36
4	D	9/19 (47%)	9 (100%)	0	100	100
All	All	703/767 (92%)	674 (96%)	29 (4%)	27	46

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	VAL
1	A	73	CYS
1	A	85	ARG
1	A	107	ARG
1	A	135	VAL
1	A	178	LEU
1	A	191	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	-2	HIS
2	B	18	ARG
2	B	61	GLU
2	B	76	ASN
2	B	105	LEU
2	B	151	VAL
2	B	172	LYS
2	B	174	LYS
2	B	180	LEU
2	B	182	ILE
3	C	38	GLU
3	C	59	VAL
3	C	75	VAL
3	C	78	GLU
3	C	80	VAL
3	C	84	ILE
3	C	104	THR
3	C	113	GLN
3	C	151	ASN
3	C	165	GLU
3	C	169	GLU
3	C	211	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	-2	HIS
2	B	15	HIS
2	B	21	ASN
2	B	97	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/270 (95%)	0.07	8 (3%) 51 47	30, 51, 87, 106	0
2	B	264/286 (92%)	0.59	25 (9%) 14 11	42, 63, 98, 119	0
3	C	255/282 (90%)	0.62	15 (5%) 28 24	45, 65, 102, 116	0
4	D	10/20 (50%)	0.77	1 (10%) 12 9	59, 72, 84, 94	0
All	All	786/858 (91%)	0.43	49 (6%) 26 23	30, 61, 97, 119	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	95	ALA	6.5
2	B	120	LEU	4.4
1	A	69	ASP	4.3
3	C	196	ALA	4.1
3	C	100	PRO	4.1
2	B	280	SER	3.9
2	B	213	LEU	3.3
3	C	142	THR	3.3
1	A	42	ASN	3.1
2	B	224	VAL	3.1
2	B	269	ASP	3.0
3	C	166	ALA	3.0
2	B	246	VAL	2.9
2	B	245	GLN	2.9
3	C	167	PHE	2.8
3	C	197	GLY	2.8
2	B	52	VAL	2.8
2	B	44	CYS	2.8
2	B	-2	HIS	2.8
1	A	266	SER	2.7
4	D	64	ALA	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	137	GLN	2.6
3	C	117	TYR	2.6
2	B	249	THR	2.5
1	A	174	ARG	2.5
1	A	189	THR	2.5
2	B	151	VAL	2.4
2	B	74	ALA	2.4
3	C	104	THR	2.4
2	B	223	ASN	2.4
2	B	46	LYS	2.3
2	B	12	CYS	2.3
2	B	0	HIS	2.2
3	C	198	SER	2.2
2	B	248	PRO	2.2
2	B	247	ASN	2.1
1	A	105	SER	2.1
2	B	126	SER	2.1
3	C	145	PHE	2.1
3	C	15	TYR	2.1
2	B	45	ASP	2.1
1	A	66	PRO	2.1
3	C	170	LEU	2.1
2	B	153	ASP	2.1
3	C	150	THR	2.1
1	A	73	CYS	2.0
2	B	172	LYS	2.0
3	C	46	LYS	2.0
2	B	267	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.