



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:14 AM UTC

PDB ID : 2J9G / pdb_00002j9g
Title : Crystal structure of Biotin carboxylase from E. coli in complex with AMPPNP and ADP
Authors : Mochalkin, I.
Deposited on : 2008-03-21
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	449	6% 96% ..
1	B	449	16% 94% 5% .

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7604 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BIOTIN CARBOXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	444	Total	C	N	O	S	0	3	0
			3441	2166	616	637	22			
1	B	442	Total	C	N	O	S	8	3	1
			3424	2156	612	634	22			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	367	Total	O	0	0
			367	367		
6	B	302	Total	O	0	0
			302	302		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BIOTIN CARBOXYLASE



• Molecule 1: BIOTIN CARBOXYLASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.70Å 112.25Å 121.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	82.48 – 2.05 82.48 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.7 (82.48-2.05) 97.3 (82.48-2.05)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.63 (at 2.04Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.190 , 0.236 0.206 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	26.5	Xtriage
Anisotropy	0.357	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7604	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, MG, ANP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.48	0/3513	0.71	2/4738 (0.0%)
1	B	0.49	1/3484 (0.0%)	0.68	0/4700
All	All	0.49	1/6997 (0.0%)	0.70	2/9438 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	446	LEU	C-N	-6.56	1.24	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	348	LEU	CA-C-N	6.18	127.57	119.84
1	A	348	LEU	C-N-CA	6.18	127.57	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3441	0	3464	9	0
1	B	3424	0	3449	14	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	31	0	13	4	0
5	B	27	0	12	0	0
6	A	367	0	0	0	0
6	B	302	0	0	3	0
All	All	7604	0	6938	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:HIS:HD1	1:A:81:GLY:H	1.37	0.69
1:B:79:HIS:HD1	1:B:81:GLY:H	1.44	0.63
1:A:116:LYS:HZ1	4:A:1449:ANP:PB	2.26	0.59
1:B:1[B]:MET:HE3	1:B:319:GLN:OE1	2.04	0.58
1:B:228:ARG:HH21	1:B:294:GLN:NE2	2.03	0.56
1:A:116:LYS:NZ	4:A:1449:ANP:PB	2.78	0.56
1:B:1[B]:MET:HA	1:B:1[B]:MET:HE2	1.90	0.52
1:B:157:ILE:N	6:B:2122:HOH:O	2.41	0.51
1:A:116:LYS:HZ3	4:A:1449:ANP:PG	2.35	0.50
1:B:340:ASN:HD22	1:B:384:MET:HA	1.76	0.49
1:A:131:VAL:HG22	1:A:285:TYR:HB3	1.96	0.48
1:B:1[B]:MET:HE1	1:B:314:ARG:HA	1.96	0.47
1:B:57[A]:VAL:HG23	6:B:2050:HOH:O	2.16	0.46
1:A:169:MET:HE1	4:A:1449:ANP:O4'	2.17	0.45
1:A:290:ASN:ND2	1:A:294:GLN:OE1	2.37	0.45
1:B:343:ASP:O	1:B:347:PHE:HA	2.18	0.44
1:B:298:PRO:O	1:B:302:MET:HG2	2.18	0.43
1:A:82:TYR:CZ	1:A:295:VAL:HG22	2.53	0.43
1:B:157:ILE:HG22	6:B:2122:HOH:O	2.19	0.42
1:B:2:LEU:HD21	1:B:101:ILE:HD12	2.02	0.42
1:B:290:ASN:ND2	1:B:294:GLN:OE1	2.49	0.41
1:A:63:ILE:HB	1:A:64:PRO:HD3	2.02	0.41
1:B:297:HIS:N	1:B:298:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	443/449 (99%)	426 (96%)	16 (4%)	1 (0%)	43	37
1	B	438/449 (98%)	424 (97%)	14 (3%)	0	100	100
All	All	881/898 (98%)	850 (96%)	30 (3%)	1 (0%)	48	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	349	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	361/361 (100%)	360 (100%)	1 (0%)	86	90
1	B	358/361 (99%)	354 (99%)	4 (1%)	65	68
All	All	719/722 (100%)	714 (99%)	5 (1%)	78	80

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	206	ASN
1	B	1[A]	MET
1	B	1[B]	MET
1	B	206	ASN
1	B	259	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	GLN
1	A	180	GLN
1	A	206	ASN
1	A	236	HIS
1	A	237	GLN
1	A	312	GLN
1	A	358	HIS
1	A	404	ASN
1	A	426	ASN
1	B	145	ASN
1	B	180	GLN
1	B	206	ASN
1	B	236	HIS
1	B	290	ASN
1	B	294	GLN
1	B	312	GLN
1	B	340	ASN
1	B	404	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ADP	B	1449	2	28,29,29	1.49	5 (17%)	43,45,45	1.80	9 (20%)
3	SO4	A	1448	-	4,4,4	0.24	0	6,6,6	0.15	0
4	ANP	A	1449	-	33,33,33	2.09	10 (30%)	45,52,52	2.11	14 (31%)
3	SO4	B	1448	-	4,4,4	0.29	0	6,6,6	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ADP	B	1449	2	-	2/16/32/32	0/3/3/3
4	ANP	A	1449	-	-	5/18/38/38	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1449	ANP	PB-N3B	5.05	1.76	1.63
4	A	1449	ANP	PG-N3B	4.96	1.76	1.63
5	B	1449	ADP	C5-C4	4.87	1.47	1.39
4	A	1449	ANP	C5-C4	4.54	1.47	1.39
4	A	1449	ANP	PB-O1B	3.73	1.51	1.46
4	A	1449	ANP	PG-O1G	3.11	1.50	1.46
5	B	1449	ADP	C5-C6	2.70	1.48	1.41
5	B	1449	ADP	PA-O3A	2.68	1.62	1.59
4	A	1449	ANP	C5-C6	2.59	1.48	1.41
4	A	1449	ANP	PB-O3A	2.43	1.62	1.59
5	B	1449	ADP	C8-N7	2.37	1.36	1.31
4	A	1449	ANP	C5-N7	-2.36	1.34	1.39
5	B	1449	ADP	C5-N7	-2.35	1.34	1.39
4	A	1449	ANP	C8-N7	2.17	1.35	1.31
4	A	1449	ANP	PA-O3A	2.17	1.61	1.59

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1449	ANP	C5-C4-N3	-5.86	118.65	126.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1449	ADP	C5-C4-N3	-5.74	118.81	126.72
4	A	1449	ANP	O1G-PG-N3B	-5.32	103.93	111.77
4	A	1449	ANP	N3-C4-N9	4.70	135.15	127.17
5	B	1449	ADP	N3-C4-N9	4.49	134.80	127.17
4	A	1449	ANP	O2B-PB-O1B	3.92	118.28	109.87
4	A	1449	ANP	C2-N3-C4	3.90	121.35	111.83
5	B	1449	ADP	C2-N3-C4	3.77	121.05	111.83
4	A	1449	ANP	N3-C2-N1	-3.64	123.07	128.58
5	B	1449	ADP	N3-C2-N1	-3.57	123.18	128.58
5	B	1449	ADP	C4-C5-N7	-3.43	106.66	110.58
4	A	1449	ANP	C4-C5-N7	-3.36	106.74	110.58
4	A	1449	ANP	O2G-PG-O3G	2.65	114.72	107.59
4	A	1449	ANP	C5-N7-C8	2.64	107.60	103.45
4	A	1449	ANP	C4-N9-C8	2.58	108.44	105.74
5	B	1449	ADP	C5-N7-C8	2.55	107.46	103.45
5	B	1449	ADP	C4-N9-C8	2.47	108.33	105.74
4	A	1449	ANP	O4'-C1'-N9	2.35	112.60	108.09
4	A	1449	ANP	C6-C5-N7	2.23	136.38	132.09
4	A	1449	ANP	C2'-C1'-N9	-2.16	107.94	113.30
5	B	1449	ADP	C6-C5-N7	2.13	136.20	132.09
4	A	1449	ANP	N9-C8-N7	-2.05	111.02	113.94
5	B	1449	ADP	C2-N1-C6	2.02	122.05	118.73

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1449	ANP	C5'-O5'-PA-O1A
4	A	1449	ANP	C5'-O5'-PA-O3A
5	B	1449	ADP	C5'-O5'-PA-O1A
4	A	1449	ANP	C3'-C4'-C5'-O5'
4	A	1449	ANP	O4'-C4'-C5'-O5'
4	A	1449	ANP	PB-N3B-PG-O1G
5	B	1449	ADP	PB-O3A-PA-O2A


There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1449	ANP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	444/449 (98%)	0.19	27 (6%)	27 27	13, 24, 43, 56	5 (1%)
1	B	441/449 (98%)	0.67	71 (16%)	4 4	11, 24, 69, 127	4 (0%)
All	All	885/898 (98%)	0.43	98 (11%)	10 10	11, 24, 64, 127	9 (1%)

All (98) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	182	ILE	9.2
1	B	178	LEU	9.1
1	B	179	ALA	9.0
1	B	183	SER	7.5
1	B	172	VAL	7.3
1	B	191	ALA	7.1
1	B	187	ALA	6.9
1	B	189	ALA	6.8
1	B	171	VAL	6.7
1	B	176	ALA	6.1
1	B	156	VAL	5.9
1	B	152	ILE	5.8
1	B	139	GLY	5.4
1	B	198	VAL	5.3
1	B	170	ARG	5.2
1	B	181	SER	5.1
1	B	200	MET	5.1
1	B	185	THR	5.0
1	B	158	ILE	5.0
1	B	157	ILE	4.7
1	B	149	ALA	4.7
1	A	349	PRO	4.3
1	B	206	ASN	4.2
1	B	148	ILE	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	137	PRO	4.1
1	A	348	LEU	4.0
1	B	177	GLU	3.9
1	B	184	MET	3.9
1	B	160	ALA	3.9
1	B	138	LEU	3.9
1	B	155	PRO	3.8
1	B	180	GLN	3.8
1	B	140	ASP	3.7
1	A	446	LEU	3.7
1	B	196	ASP	3.7
1	A	163	GLY	3.6
1	B	147	ALA	3.5
1	A	347	PHE	3.5
1	B	348	LEU	3.4
1	B	197	MET	3.4
1	B	154	TYR	3.4
1	A	194	SER	3.4
1	A	184	MET	3.4
1	B	167	ARG	3.3
1	A	445	GLY	3.3
1	B	201	GLU	3.3
1	A	188	GLU	3.2
1	A	1	MET	3.2
1	B	150	LYS	3.2
1	A	189	ALA	3.2
1	B	446	LEU	3.2
1	B	199	TYR	3.1
1	B	188	GLU	3.1
1	B	159	LYS	3.1
1	B	142	MET	3.1
1	A	187	ALA	3.1
1	A	346	THR	3.0
1	B	346	THR	3.0
1	A	165	GLY	3.0
1	A	140	ASP	3.0
1	B	175	ASP	3.0
1	B	144	LYS	3.0
1	B	135	ASP	2.9
1	B	132	PRO	2.9
1	B	174	GLY	2.9
1	B	186	ARG	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	145	ASN	2.9
1	B	190	LYS	2.8
1	A	344	PRO	2.8
1	A	164	GLY	2.7
1	A	167	ARG	2.7
1	B	173	ARG	2.6
1	A	162	GLY	2.6
1	B	133	GLY	2.6
1	B	161	SER	2.6
1	A	193	PHE	2.5
1	A	180	GLN	2.5
1	A	235	ARG	2.5
1	B	146	ARG	2.5
1	B	195	ASN	2.5
1	A	190	LYS	2.4
1	B	345	ASN	2.4
1	B	143	ASP	2.4
1	B	97	ARG	2.2
1	A	141	ASP	2.2
1	B	126	ALA	2.2
1	B	344	PRO	2.2
1	B	141	ASP	2.2
1	B	114	GLY	2.2
1	B	382	ASP	2.2
1	A	166	GLY	2.2
1	B	164	GLY	2.2
1	B	129	PRO	2.1
1	B	127	GLY	2.1
1	B	153	GLY	2.1
1	B	93	GLU	2.1
1	A	442	LYS	2.1
1	A	382	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	1448	5/5	0.73	0.23	47,48,48,49	0
4	ANP	A	1449	31/31	0.79	0.13	34,40,62,62	0
3	SO4	B	1448	5/5	0.80	0.15	51,52,53,53	0
5	ADP	B	1449	27/27	0.89	0.10	37,38,40,40	0
2	MG	A	1447	1/1	0.94	0.05	42,42,42,42	0
2	MG	B	1447	1/1	0.96	0.04	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around ADP B 1449:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.