



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 02:14 PM UTC

PDB ID : 2JA5 / pdb_00002ja5
Title : CPD lesion containing RNA Polymerase II elongation complex A
Authors : Brueckner, F.; Hennecke, U.; Carell, T.; Cramer, P.
Deposited on : 2006-11-23
Resolution : 3.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1065 (3.96-3.64)
Clashscore	190562	1012 (3.94-3.66)
Ramachandran outliers	187476	1048 (3.96-3.64)
Sidechain outliers	187428	1043 (3.96-3.64)
RSRZ outliers	180081	1064 (3.96-3.64)
RNA backbone	3983	1007 (4.50-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1733	2% 23% 45% 13% • 18%
2	B	1224	5% 24% 52% 14% • 9%
3	C	318	% 23% 47% 11% • 16%
4	D	221	4% 22% 40% 15% • 20%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	215	
6	F	155	
7	G	171	
8	H	146	
9	I	122	
10	J	70	
11	K	120	
12	L	70	
13	P	11	
14	T	25	

2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 31660 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1421	Total	C	N	O	S	0	0	0
			11186	7048	1958	2118	62			

- Molecule 2 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1115	Total	C	N	O	S	0	0	0
			8866	5614	1553	1644	55			

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	267	Total	C	N	O	S	0	0	0
			2101	1320	349	419	13			

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	177	Total	C	N	O	S	0	0	0
			1427	882	256	287	2			

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	214	Total	C	N	O	S	0	0	0
			1752	1111	309	321	11			

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	87	Total	C	N	O	S	0	0	0
			705	451	119	132	3			

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	171	Total	C	N	O	S	0	0	0
			1340	861	222	249	8			

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	135	Total	C	N	O	S	0	0	0
			1084	683	183	214	4			

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	116	Total	C	N	O	S	0	0	0
			944	581	172	181	10			

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	65	Total	C	N	O	S	0	0	0
			532	339	93	94	6			

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	114	Total	C	N	O	S	0	0	0
			919	590	156	171	2			

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	46	Total	C	N	O	S	0	0	0
			364	224	72	64	4			

- Molecule 13 is a RNA chain called 5'-R(*UP*UP*CP*GP*AP*CP*CP*AP*GP*GP*AP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	P	10	Total	C	N	O	P	0	0	0
			212	96	41	66	9			

- Molecule 14 is a DNA chain called 5'-D(*AP*GP*CP*TP*CP*AP*AP*GP*TP*AP *CP*

TP*TTP*TP*TP*CP*CP*BRUP*GP*GP*TP*CP*AP*TP*T)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
14	T	11	Total	Br	C	N	O	P	0	0	0
			219	1	106	34	68	10			

- Molecule 15 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
15	A	1	Total	Mg	0	0
			1	1		


- Molecule 16 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	8	Total	Zn	0	0
			8	8		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA-directed RNA polymerase II subunit RPB1






Chain B: 5% 24% 52% 14% 9%







• Molecule 3: DNA-directed RNA polymerase II subunit RPB3




• Molecule 4: DNA-directed RNA polymerase II subunit RPB4




- Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC1




- Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2




- Molecule 7: DNA-directed RNA polymerase II subunit RPB7







• Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3




• Molecule 9: DNA-directed RNA polymerase II subunit RPB9




• Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5




• Molecule 11: DNA-directed RNA polymerase II subunit RPB11




- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4



- Molecule 13: 5'-R(*UP*UP*CP*GP*AP*CP*CP*AP*GP*GP*AP)-3'



- Molecule 14: 5'-D(*AP*GP*CP*TP*CP*AP*AP*GP*TP*AP *CP*TP*TTP*TP*TP*CP*CP *BRUP*GP*GP*TP*CP*AP*TP*T)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	222.21Å 392.21Å 284.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.80 50.00 – 3.80	Depositor EDS
% Data completeness (in resolution range)	97.4 (50.00-3.80) 97.4 (50.00-3.80)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 3.77Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.270 , 0.280 0.215 , 0.235	Depositor DCC
R_{free} test set	4681 reflections (1.98%)	wwPDB-VP
Wilson B-factor (Å ²)	99.7	Xtriage
Anisotropy	0.551	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 81.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.024 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.027 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	31660	wwPDB-VP
Average B, all atoms (Å ²)	87.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG, BRU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.59	0/11385	1.11	79/15393 (0.5%)
2	B	0.56	1/9037 (0.0%)	1.08	62/12181 (0.5%)
3	C	0.59	0/2138	1.10	14/2896 (0.5%)
4	D	0.47	0/1437	1.11	17/1925 (0.9%)
5	E	0.51	0/1788	0.98	11/2406 (0.5%)
6	F	0.65	0/716	1.14	3/964 (0.3%)
7	G	0.54	0/1368	1.13	13/1844 (0.7%)
8	H	0.48	0/1102	1.04	6/1492 (0.4%)
9	I	0.47	0/962	1.08	6/1295 (0.5%)
10	J	0.56	0/541	1.01	1/727 (0.1%)
11	K	1.27	9/937 (1.0%)	1.43	16/1265 (1.3%)
12	L	0.55	0/366	0.94	0/485
13	P	0.80	0/237	1.28	4/368 (1.1%)
14	T	0.61	0/220	1.62	8/335 (2.4%)
All	All	0.60	10/32234 (0.0%)	1.10	240/43576 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
14	T	0	2
All	All	0	4

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	K	112	GLN	CA-C	19.29	1.78	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	K	113	THR	CA-C	13.26	1.69	1.53
11	K	113	THR	N-CA	12.98	1.64	1.46
11	K	112	GLN	N-CA	12.71	1.62	1.46
11	K	112	GLN	CB-CG	8.18	1.76	1.52

The worst 5 of 240 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	K	113	THR	N-CA-C	21.59	136.74	108.24
4	D	26	THR	N-CA-C	-13.01	96.72	112.59
2	B	1185	CYS	N-CA-C	-12.95	97.49	113.28
2	B	473	MET	N-CA-C	-11.50	98.90	113.16
11	K	112	GLN	N-CA-C	11.01	134.26	110.80

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	303	TYR	Sidechain
2	B	486	TYR	Sidechain
14	T	19	DT	Sidechain
14	T	20	DC	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11186	0	11266	1395	0
2	B	8866	0	8898	1083	0
3	C	2101	0	2055	285	0
4	D	1427	0	1451	151	0
5	E	1752	0	1776	140	0
6	F	705	0	730	89	0
7	G	1340	0	1357	172	0
8	H	1084	0	1057	130	0
9	I	944	0	899	110	0
10	J	532	0	542	104	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	K	919	0	929	120	0
12	L	364	0	386	44	0
13	P	212	0	109	20	0
14	T	219	0	125	31	0
15	A	1	0	0	0	0
16	A	8	0	0	0	0
All	All	31660	0	31580	3568	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 56.

The worst 5 of 3568 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:K:112:GLN:CB	11:K:112:GLN:CG	1.77	1.62
11:K:112:GLN:C	11:K:112:GLN:CA	1.78	1.56
2:B:343:ILE:HG23	2:B:347:LYS:HB2	1.18	1.17
2:B:273:LEU:HB2	2:B:276:ILE:HD12	1.26	1.17
1:A:1445:ILE:H	1:A:1445:ILE:HD12	1.12	1.15

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1410/1733 (81%)	975 (69%)	289 (20%)	146 (10%)	0	6
2	B	1096/1224 (90%)	781 (71%)	200 (18%)	115 (10%)	0	6
3	C	264/318 (83%)	171 (65%)	65 (25%)	28 (11%)	0	6
4	D	173/221 (78%)	124 (72%)	31 (18%)	18 (10%)	0	6
5	E	212/215 (99%)	154 (73%)	44 (21%)	14 (7%)	1	13

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
6	F	84/155 (54%)	69 (82%)	11 (13%)	4 (5%)	2	17
7	G	169/171 (99%)	131 (78%)	28 (17%)	10 (6%)	1	15
8	H	131/146 (90%)	82 (63%)	32 (24%)	17 (13%)	0	3
9	I	114/122 (93%)	80 (70%)	23 (20%)	11 (10%)	0	7
10	J	63/70 (90%)	37 (59%)	12 (19%)	14 (22%)	0	1
11	K	112/120 (93%)	85 (76%)	16 (14%)	11 (10%)	0	7
12	L	44/70 (63%)	18 (41%)	17 (39%)	9 (20%)	0	1
All	All	3872/4565 (85%)	2707 (70%)	768 (20%)	397 (10%)	0	6

5 of 397 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	THR
1	A	48	ALA
1	A	57	ARG
1	A	62	ASP
1	A	65	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1244/1520 (82%)	1116 (90%)	128 (10%)	7	27
2	B	967/1061 (91%)	881 (91%)	86 (9%)	9	32
3	C	235/274 (86%)	212 (90%)	23 (10%)	7	29
4	D	159/200 (80%)	133 (84%)	26 (16%)	2	14
5	E	196/197 (100%)	191 (97%)	5 (3%)	40	60
6	F	77/137 (56%)	70 (91%)	7 (9%)	9	32
7	G	152/152 (100%)	140 (92%)	12 (8%)	11	36
8	H	119/128 (93%)	113 (95%)	6 (5%)	22	47
9	I	110/116 (95%)	97 (88%)	13 (12%)	5	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
10	J	60/65 (92%)	55 (92%)	5 (8%)	10	34
11	K	99/102 (97%)	89 (90%)	10 (10%)	7	27
12	L	40/57 (70%)	37 (92%)	3 (8%)	12	38
All	All	3458/4009 (86%)	3134 (91%)	324 (9%)	8	30

5 of 324 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	214	ASN
7	G	171	ILE
4	D	13	ARG
4	D	206	GLU
9	I	78	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 106 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	821	GLN
3	C	112	ASN
9	I	60	GLN
2	B	842	ASN
2	B	1076	HIS

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
13	P	9/11 (81%)	1 (11%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
13	P	10	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
14	BRU	T	22	13,14	18,21,22	0.62	0	25,30,33	1.24	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	BRU	T	22	13,14	-	0/7/21/22	0/2/2/2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	T	22	BRU	C6-C5-C4	-3.27	117.36	120.67
14	T	22	BRU	BR-C5-C4	2.72	121.15	118.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
14	T	22	BRU	3	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	2
3	C	1
6	F	1
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	2:SER	C	3:GLU	N	4.08
1	B	18:PHE	C	19:GLU	N	3.80
1	F	69:LEU	C	70:LYS	N	3.44
1	A	1175:SER	C	1176:LEU	N	3.41
1	B	337:ARG	C	338:GLY	N	2.62

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1421/1733 (81%)	0.10	40 (2%) 55 38	12, 73, 146, 189	0
2	B	1115/1224 (91%)	0.33	63 (5%) 29 23	12, 83, 154, 191	0
3	C	267/318 (83%)	-0.02	4 (1%) 72 50	30, 69, 125, 150	0
4	D	177/221 (80%)	0.41	9 (5%) 33 25	53, 104, 144, 162	0
5	E	214/215 (99%)	0.38	3 (1%) 73 51	45, 126, 174, 178	0
6	F	87/155 (56%)	-0.25	2 (2%) 61 42	17, 49, 91, 119	0
7	G	171/171 (100%)	0.20	7 (4%) 41 30	53, 77, 115, 126	0
8	H	135/146 (92%)	0.86	15 (11%) 10 13	87, 128, 162, 171	0
9	I	116/122 (95%)	0.61	9 (7%) 19 17	68, 122, 151, 173	0
10	J	65/70 (92%)	-0.09	1 (1%) 72 50	35, 66, 109, 118	0
11	K	114/120 (95%)	0.01	4 (3%) 47 33	33, 73, 102, 140	0
12	L	46/70 (65%)	1.21	6 (13%) 7 11	69, 141, 161, 169	0
13	P	10/11 (90%)	0.86	2 (20%) 3 5	71, 85, 160, 163	0
14	T	10/25 (40%)	1.09	2 (20%) 3 5	72, 96, 137, 158	0
All	All	3948/4601 (85%)	0.24	167 (4%) 40 29	12, 82, 154, 191	0

The worst 5 of 167 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	882	THR	8.5
11	K	113	THR	7.5
4	D	25	ALA	6.4
1	A	1081	LEU	6.1
8	H	90	ALA	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
14	BRU	T	22	20/21	0.85	0.12	59,61,68,69	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
16	ZN	A	2461	1/1	0.97	0.09	163,163,163,163	0
16	ZN	A	2464	1/1	0.98	0.05	82,82,82,82	0
16	ZN	A	2459	1/1	0.99	0.03	115,115,115,115	0
16	ZN	A	2460	1/1	1.00	0.05	69,69,69,69	0
16	ZN	A	2458	1/1	1.00	0.04	51,51,51,51	0
16	ZN	A	2462	1/1	1.00	0.01	23,23,23,23	0
16	ZN	A	2463	1/1	1.00	0.01	42,42,42,42	0
15	MG	A	2457	1/1	1.00	0.02	21,21,21,21	0
16	ZN	A	2465	1/1	1.00	0.02	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.