



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 13, 2026 – 11:38 PM UTC

PDB ID : 4JA8 / pdb_00004ja8
Title : Complex of Mitochondrial Isocitrate Dehydrogenase R140Q Mutant with AGI-6780 Inhibitor
Authors : Wei, W.; Chen, L.; Wu, M.; Jiang, F.; Travins, J.; Qian, K.; DeLaBarre, B.
Deposited on : 2013-02-18
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	416	
1	B	416	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7629 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Isocitrate dehydrogenase [NADP], mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	416	Total	C	N	O	S	0	13	0
			3428	2175	602	633	18			
1	B	406	Total	C	N	O	S	0	6	0
			3269	2081	562	607	19			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	140	GLN	ARG	engineered mutation	UNP P48735
A	453	HIS	-	expression tag	UNP P48735
A	454	HIS	-	expression tag	UNP P48735
A	455	HIS	-	expression tag	UNP P48735
A	456	HIS	-	expression tag	UNP P48735
B	140	GLN	ARG	engineered mutation	UNP P48735
B	453	HIS	-	expression tag	UNP P48735
B	454	HIS	-	expression tag	UNP P48735
B	455	HIS	-	expression tag	UNP P48735
B	456	HIS	-	expression tag	UNP P48735

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is 1-[5-(cyclopropylsulfamoyl)-2-thiophen-3-yl-phenyl]-3-[3-(trifluoromethyl)phenyl]urea (CCD ID: 1K9) (formula: C₂₁H₁₈F₃N₃O₃S₂).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	S	0	1
			64	42	6	6	6	4		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Ca 1 1	0	0
4	B	1	Total Ca 1 1	0	0

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	469	Total O 469 469	0	0
6	B	277	Total O 277 277	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Isocitrate dehydrogenase [NADP], mitochondrial



- Molecule 1: Isocitrate dehydrogenase [NADP], mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.66Å 119.66Å 125.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.90 – 1.55 31.90 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.5 (31.90-1.55) 99.5 (31.90-1.55)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.182 , 0.218 0.180 , 0.217	Depositor DCC
R_{free} test set	6461 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	14.6	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.011 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7629	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GOL, 1K9, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.62	18/3507 (0.5%)	1.27	4/4733 (0.1%)
1	B	1.50	22/3343 (0.7%)	1.25	3/4516 (0.1%)
All	All	1.56	40/6850 (0.6%)	1.26	7/9249 (0.1%)

All (40) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	ILE	CA-CB	8.42	1.64	1.54
1	A	154	CYS	N-CA	7.63	1.55	1.46
1	A	174	ALA	N-CA	7.52	1.56	1.46
1	A	216	GLY	N-CA	7.15	1.55	1.45
1	B	295	ALA	C-O	6.79	1.32	1.24
1	A	172	ARG	N-CA	6.34	1.53	1.46
1	B	114	ALA	CA-CB	6.31	1.64	1.53
1	B	224	THR	N-CA	6.10	1.53	1.45
1	A	400	LYS	CG-CD	6.03	1.70	1.52
1	B	197	THR	N-CA	5.86	1.53	1.46
1	B	75	VAL	CA-CB	5.83	1.62	1.54
1	A	60	ARG	CD-NE	5.79	1.54	1.46
1	B	271	ASP	C-O	5.75	1.30	1.24
1	B	266	PHE	CA-CB	5.75	1.62	1.53
1	B	364	THR	N-CA	5.74	1.52	1.45
1	B	93	ASP	CA-C	5.74	1.60	1.52
1	B	317	SER	N-CA	5.69	1.53	1.46
1	B	216	GLY	N-CA	5.56	1.53	1.45
1	B	186	ALA	N-CA	5.48	1.52	1.46
1	A	68	GLU	C-O	5.46	1.31	1.24
1	B	195	VAL	N-CA	5.46	1.52	1.46
1	B	197	THR	CA-CB	-5.44	1.46	1.53
1	B	353	ARG	C-O	5.43	1.30	1.24
1	A	406	VAL	CA-CB	5.39	1.61	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	447	ARG	CZ-NH1	5.38	1.40	1.32
1	A	420	HIS	C-O	5.30	1.30	1.23
1	B	97	THR	N-CA	5.29	1.52	1.46
1	A	161	VAL	N-CA	5.24	1.51	1.46
1	B	238	TYR	CA-CB	5.21	1.61	1.53
1	A	140	GLN	CD-OE1	5.19	1.33	1.23
1	A	438	LEU	CA-C	5.16	1.59	1.52
1	B	334	LEU	C-O	5.15	1.30	1.24
1	B	246	LEU	CA-C	-5.13	1.46	1.52
1	A	317	SER	CA-CB	-5.13	1.44	1.53
1	A	168	ILE	CA-CB	-5.09	1.48	1.54
1	A	383	GLY	N-CA	5.08	1.52	1.45
1	A	318	ASP	CA-CB	5.05	1.61	1.53
1	B	62	ILE	N-CA	-5.04	1.40	1.46
1	B	269	ILE	CA-CB	5.04	1.60	1.54
1	B	268	GLU	N-CA	5.00	1.52	1.46

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	384	LYS	N-CA-C	-6.12	104.53	111.14
1	A	120	GLU	N-CA-C	5.71	117.96	111.11
1	A	292	ASP	N-CA-C	-5.52	105.34	111.36
1	A	125	GLU	N-CA-C	5.32	117.49	111.11
1	B	157	ILE	O-C-N	5.16	124.62	121.37
1	B	97	THR	N-CA-C	-5.09	105.64	111.14
1	B	61	ILE	N-CA-C	5.04	115.26	110.42

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3428	0	3382	34	0
1	B	3269	0	3241	31	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	48	0	26	1	0
2	B	48	0	26	1	0
3	A	64	0	36	3	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	24	0	30	4	0
6	A	469	0	0	11	0
6	B	277	0	0	4	0
All	All	7629	0	6741	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330:MET:HE2	6:A:993:HOH:O	1.15	1.25
1:B:288:ARG:HE	1:B:296:GLN:HE22	1.04	0.99
1:B:286:GLU:OE2	1:B:288:ARG:NH1	2.06	0.89
1:A:294:VAL:HB	1:B:319[B]:ILE:HD11	1.69	0.75
1:B:109:VAL:HG22	1:B:381:HIS:HD2	1.50	0.74
5:A:506:GOL:H32	1:B:205:LYS:HE3	1.72	0.71
1:A:105[A]:GLN:HG3	1:A:143:LEU:CD2	2.23	0.68
1:B:75:VAL:HG12	1:B:77:ILE:HG22	1.77	0.66
1:A:322:GLN:HG3	6:A:855:HOH:O	1.96	0.65
1:A:319[A]:ILE:HD11	1:B:294:VAL:HB	1.81	0.63
5:A:506:GOL:H12	1:B:205:LYS:HE3	1.79	0.62
1:B:43:ARG:HD3	1:B:76:ASP:HB2	1.80	0.62
1:B:288:ARG:HE	1:B:296:GLN:NE2	1.89	0.60
1:B:425:VAL:HG22	1:B:431:PHE:HD2	1.66	0.59
1:B:50:VAL:CG2	1:B:382:ARG:HG3	2.34	0.58
1:B:288:ARG:NE	1:B:296:GLN:HE22	1.88	0.57
1:A:273:HIS:HE1	6:A:894:HOH:O	1.86	0.56
6:A:946:HOH:O	1:B:199:LYS:HD2	2.03	0.56
1:A:74:HIS:HE1	6:A:818:HOH:O	1.89	0.56
1:B:327:LEU:HB3	1:B:348:HIS:HB3	1.88	0.56
1:B:77:ILE:HD12	6:B:718:HOH:O	2.07	0.54
1:A:105[A]:GLN:CG	1:A:143:LEU:CD2	2.86	0.54
1:A:327:LEU:HB3	1:A:348:HIS:HB3	1.90	0.53
1:B:340:LYS:HD3	6:B:868:HOH:O	2.10	0.51
1:A:330:MET:HB2	6:A:993:HOH:O	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105[A]:GLN:HG3	1:A:143:LEU:HD21	1.92	0.50
1:B:115:THR:O	2:B:501:NDP:H2N	2.12	0.49
1:A:42:LYS:HD3	1:A:453[B]:HIS:CD2	2.47	0.49
1:A:444:ASN:OD1	1:A:447:ARG:NH2	2.46	0.49
1:A:377[B]:ARG:HD3	1:A:399:GLU:OE2	2.13	0.49
5:A:506:GOL:H32	1:B:205:LYS:CE	2.41	0.48
1:A:74:HIS:HD2	6:A:939:HOH:O	1.97	0.48
1:A:115:THR:O	2:A:501:NDP:H2N	2.14	0.48
1:B:155:LYS:HG2	1:B:407:GLU:OE2	2.14	0.47
1:B:120:GLU:H	1:B:120:GLU:HG2	1.55	0.47
1:B:398:LEU:HA	1:B:401:VAL:HG22	1.97	0.46
1:A:249:SER:HA	1:A:288:ARG:O	2.15	0.46
1:B:50:VAL:HG23	1:B:382:ARG:HG3	1.96	0.46
1:B:169:THR:HG21	1:B:238:TYR:CE2	2.51	0.46
1:A:176:GLY:H	1:A:180:LYS:HZ3	1.62	0.45
3:A:502[A]:1K9:OAA	3:A:502[A]:1K9:H7	2.15	0.45
1:B:296:GLN:CG	6:B:874:HOH:O	2.64	0.45
1:B:75:VAL:HG12	1:B:77:ILE:CG2	2.46	0.45
1:A:332:SER:HB3	1:A:345[A]:GLU:HG2	1.98	0.44
1:A:109:VAL:HG22	1:A:381:HIS:HD2	1.82	0.44
1:A:319[B]:ILE:HG23	1:B:298:LEU:HD11	2.00	0.43
1:A:92:THR:O	1:A:95[B]:GLN:HG3	2.19	0.42
1:A:112:LYS:NZ	1:A:136:ASN:HD21	2.17	0.42
3:A:502[B]:1K9:OAA	3:A:502[B]:1K9:H7	2.19	0.42
1:A:272[B]:LYS:HD2	6:A:1041:HOH:O	2.19	0.42
1:B:405:THR:O	1:B:408:SER:HB2	2.18	0.42
1:A:345[B]:GLU:HG3	6:A:604:HOH:O	2.20	0.42
1:A:175:HIS:CD2	1:A:180:LYS:HE2	2.55	0.41
1:A:256:LYS:NZ	6:A:995:HOH:O	2.48	0.41
1:A:276:THR:OG1	5:A:504:GOL:H12	2.21	0.41
1:A:42:LYS:CD	1:A:453[B]:HIS:CD2	3.04	0.41
1:A:312:ASP:O	1:A:316[B]:GLN:HG2	2.20	0.41
1:A:330:MET:CE	6:A:993:HOH:O	2.04	0.41
3:A:502[B]:1K9:OAA	3:A:502[B]:1K9:H1	2.20	0.41
1:B:397:MET:SD	1:B:448:ALA:HB1	2.60	0.41
1:B:401:VAL:O	1:B:405:THR:OG1	2.34	0.41
1:A:320[B]:LEU:HG	1:A:330:MET:HE1	2.02	0.40
1:A:426:LYS:HB3	1:A:429:GLU:HB3	2.03	0.40
1:B:339:GLY:HA2	6:B:825:HOH:O	2.21	0.40
1:A:50:VAL:HG23	1:A:77:ILE:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	427/416 (103%)	413 (97%)	14 (3%)	0	100	100
1	B	410/416 (99%)	391 (95%)	19 (5%)	0	100	100
All	All	837/832 (101%)	804 (96%)	33 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/357 (104%)	366 (99%)	4 (1%)	65	43
1	B	354/357 (99%)	338 (96%)	16 (4%)	24	3
All	All	724/714 (101%)	704 (97%)	20 (3%)	38	11

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	204	VAL
1	A	340	LYS
1	A	351	VAL
1	A	427	LEU
1	B	45	LYS
1	B	59	THR
1	B	67	LYS
1	B	77	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	78	GLN
1	B	80	LYS
1	B	105	GLN
1	B	120	GLU
1	B	130	LYS
1	B	180	LYS
1	B	340	LYS
1	B	351	VAL
1	B	382	ARG
1	B	384	LYS
1	B	435	THR
1	B	442	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	HIS
1	A	136	ASN
1	A	273	HIS
1	A	420	HIS
1	A	452	GLN
1	B	105	GLN
1	B	241	GLN
1	B	296	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NDP	B	501	-	51,52,52	2.11	15 (29%)	71,80,80	2.01	20 (28%)
3	1K9	A	502[B]	-	35,35,35	2.13	6 (17%)	50,52,52	2.07	8 (16%)
2	NDP	A	501	-	51,52,52	1.70	13 (25%)	71,80,80	1.88	19 (26%)
5	GOL	A	505	-	5,5,5	0.50	0	5,5,5	0.69	0
5	GOL	A	506	-	5,5,5	0.84	0	5,5,5	0.95	0
5	GOL	A	504	-	5,5,5	1.38	1 (20%)	5,5,5	0.74	0
5	GOL	A	507	-	5,5,5	1.15	0	5,5,5	0.86	0
3	1K9	A	502[A]	-	35,35,35	2.53	13 (37%)	50,52,52	1.51	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	B	501	-	-	6/34/77/77	0/5/5/5
3	1K9	A	502[B]	-	-	1/27/31/31	0/4/4/4
2	NDP	A	501	-	-	8/34/77/77	0/5/5/5
5	GOL	A	505	-	-	0/4/4/4	-
5	GOL	A	506	-	-	4/4/4/4	-
5	GOL	A	504	-	-	0/4/4/4	-
5	GOL	A	507	-	-	2/4/4/4	-
3	1K9	A	502[A]	-	-	4/27/31/31	0/4/4/4

All (48) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502[A]	1K9	OAC-SBF	9.30	1.54	1.43
3	A	502[B]	1K9	OAC-SBF	6.92	1.51	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	NDP	PA-O3	-6.77	1.52	1.59
3	A	502[B]	1K9	CBC-CAY	-6.49	1.40	1.48
2	B	501	NDP	C4N-C3N	5.45	1.60	1.50
3	A	502[A]	1K9	CBC-CAY	-4.85	1.42	1.48
3	A	502[A]	1K9	CBE-CAZ	-4.53	1.40	1.49
2	B	501	NDP	P2B-O2B	4.31	1.67	1.59
2	B	501	NDP	O4D-C1D	4.24	1.51	1.42
3	A	502[B]	1K9	CBE-CAZ	-4.08	1.41	1.49
3	A	502[A]	1K9	CBA-SBF	3.90	1.82	1.76
3	A	502[B]	1K9	CAR-CBD	3.67	1.57	1.48
2	B	501	NDP	C4A-N9A	-3.66	1.30	1.37
2	A	501	NDP	C7N-C3N	3.64	1.56	1.48
2	A	501	NDP	PA-O3	3.53	1.63	1.59
2	A	501	NDP	C2A-N3A	3.51	1.40	1.33
2	B	501	NDP	C4A-N3A	3.28	1.40	1.34
2	A	501	NDP	C4N-C5N	-3.12	1.40	1.49
3	A	502[A]	1K9	CAO-CAZ	2.98	1.43	1.39
2	A	501	NDP	PN-O5D	-2.86	1.48	1.59
2	A	501	NDP	C1B-N9A	2.84	1.54	1.46
2	B	501	NDP	C6N-C5N	2.83	1.41	1.33
3	A	502[A]	1K9	CAP-CBA	2.82	1.44	1.39
3	A	502[A]	1K9	CAX-NAS	-2.81	1.36	1.41
2	B	501	NDP	C4N-C5N	-2.80	1.41	1.49
3	A	502[B]	1K9	CAG-CAI	2.75	1.43	1.38
3	A	502[A]	1K9	CAO-CAX	2.71	1.43	1.39
3	A	502[A]	1K9	OAB-SBF	-2.55	1.40	1.43
2	A	501	NDP	O4B-C1B	2.54	1.47	1.42
5	A	504	GOL	C1-C2	-2.54	1.42	1.51
2	A	501	NDP	PA-O2A	-2.52	1.43	1.55
2	A	501	NDP	C6N-C5N	2.48	1.40	1.33
2	A	501	NDP	C4N-C3N	2.47	1.54	1.50
3	A	502[A]	1K9	CBB-NAT	-2.46	1.36	1.41
2	B	501	NDP	PN-O3	2.46	1.62	1.59
3	A	502[A]	1K9	CAG-CAI	2.46	1.43	1.38
2	A	501	NDP	O4D-C4D	2.45	1.50	1.45
2	B	501	NDP	O3B-C3B	2.37	1.48	1.43
2	B	501	NDP	C2A-N1A	-2.29	1.29	1.33
2	B	501	NDP	O4B-C1B	2.29	1.47	1.42
3	A	502[A]	1K9	FAD-CBE	-2.28	1.24	1.33
2	A	501	NDP	PN-O2N	-2.27	1.44	1.55
2	B	501	NDP	C5D-C4D	2.18	1.58	1.51
3	A	502[B]	1K9	CAX-NAS	-2.13	1.37	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	NDP	PN-O2N	-2.09	1.45	1.55
2	A	501	NDP	P2B-O1X	2.07	1.56	1.50
2	B	501	NDP	PA-O1A	2.05	1.57	1.50
3	A	502[A]	1K9	OAA-CAW	-2.05	1.19	1.23

All (57) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502[B]	1K9	CBD-NAU-SBF	-7.84	108.36	122.18
3	A	502[B]	1K9	CAQ-CBD-NAU	-6.81	109.33	118.47
2	B	501	NDP	N9A-C8A-N7A	-5.66	105.91	113.94
2	B	501	NDP	O4D-C1D-N1N	-5.51	97.58	108.08
2	B	501	NDP	C5A-C4A-N3A	-5.27	119.46	126.72
2	A	501	NDP	N3A-C2A-N1A	-4.65	121.54	128.58
2	B	501	NDP	C4A-C5A-N7A	-4.60	105.32	110.58
2	B	501	NDP	C5A-N7A-C8A	4.35	110.28	103.45
2	A	501	NDP	C5A-N7A-C8A	4.32	110.24	103.45
2	A	501	NDP	C4A-C5A-N7A	-4.26	105.72	110.58
3	A	502[A]	1K9	CAQ-CBD-NAU	-4.22	112.81	118.47
2	A	501	NDP	C5A-C4A-N3A	-4.10	121.07	126.72
2	A	501	NDP	N9A-C8A-N7A	-3.99	108.28	113.94
2	A	501	NDP	C1D-N1N-C2N	-3.97	114.60	121.14
3	A	502[B]	1K9	CAY-CAN-SAV	-3.88	104.57	112.33
2	B	501	NDP	C5A-C4A-N9A	3.62	109.76	105.81
3	A	502[B]	1K9	CAR-CAQ-CBD	3.60	62.93	59.80
3	A	502[A]	1K9	CBD-NAU-SBF	-3.56	115.90	122.18
2	A	501	NDP	O4D-C1D-N1N	-3.43	101.54	108.08
2	B	501	NDP	O4D-C4D-C5D	-3.22	99.03	109.33
2	A	501	NDP	O3-PN-O1N	-3.12	101.33	110.70
2	B	501	NDP	O3X-P2B-O2X	3.07	119.32	107.80
2	B	501	NDP	C3D-C2D-C1D	3.06	107.25	101.46
2	B	501	NDP	C6A-C5A-C4A	3.03	121.32	117.18
2	B	501	NDP	C1D-N1N-C2N	-2.85	116.44	121.14
3	A	502[A]	1K9	CAY-CAN-SAV	-2.84	106.65	112.33
2	A	501	NDP	O2A-PA-O1A	2.83	125.59	112.44
2	B	501	NDP	C4A-N9A-C8A	2.82	108.70	105.74
2	A	501	NDP	C1D-N1N-C6N	2.74	126.56	120.77
3	A	502[A]	1K9	OAC-SBF-CBA	-2.71	104.56	107.98
2	B	501	NDP	O4D-C4D-C3D	2.68	110.48	105.15
3	A	502[A]	1K9	CAM-CBC-CAY	-2.60	116.27	119.62
2	A	501	NDP	O2N-PN-O1N	2.60	124.54	112.44
3	A	502[B]	1K9	CAK-CAY-CAN	2.54	114.79	111.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502[A]	1K9	CAM-CAL-CBA	-2.51	117.00	119.44
2	A	501	NDP	O2N-PN-O5D	2.43	118.60	107.57
2	A	501	NDP	O5B-C5B-C4B	2.38	117.11	108.99
2	A	501	NDP	O5D-PN-O1N	-2.37	99.53	108.94
3	A	502[B]	1K9	CAQ-CAR-CBD	-2.37	57.75	59.80
3	A	502[B]	1K9	CBC-CAY-CAN	-2.32	118.60	123.74
3	A	502[A]	1K9	CAK-CAY-CAN	2.29	114.42	111.17
2	A	501	NDP	O2A-PA-O3	2.27	113.42	107.27
2	B	501	NDP	O2N-PN-O3	2.19	113.18	107.27
2	B	501	NDP	O4D-C1D-C2D	-2.17	101.97	106.62
2	B	501	NDP	C3B-C2B-C1B	-2.16	98.66	102.81
2	B	501	NDP	O3D-C3D-C2D	-2.15	104.93	111.82
2	A	501	NDP	C2A-N1A-C6A	2.14	122.24	118.73
3	A	502[A]	1K9	CAI-CAX-NAS	-2.13	113.28	120.41
2	B	501	NDP	C4D-O4D-C1D	-2.12	104.79	109.47
2	A	501	NDP	N3A-C4A-N9A	2.11	130.76	127.17
2	B	501	NDP	N3A-C4A-N9A	2.11	130.75	127.17
2	A	501	NDP	C5A-C4A-N9A	2.08	108.08	105.81
2	A	501	NDP	O2B-C2B-C1B	-2.06	102.80	110.05
3	A	502[B]	1K9	CAK-CAH-SAV	-2.04	100.21	108.45
2	B	501	NDP	C3N-C2N-N1N	2.04	126.20	123.20
3	A	502[A]	1K9	CAR-CBD-NAU	2.03	121.19	118.47
3	A	502[A]	1K9	CBC-CBB-NAT	-2.01	115.09	118.70

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	506	GOL	O1-C1-C2-C3
2	B	501	NDP	C2D-C1D-N1N-C6N
5	A	506	GOL	C1-C2-C3-O3
5	A	507	GOL	C1-C2-C3-O3
5	A	506	GOL	O1-C1-C2-O2
5	A	506	GOL	O2-C2-C3-O3
3	A	502[A]	1K9	CAK-CAY-CBC-CAM
3	A	502[A]	1K9	CAK-CAY-CBC-CBB
5	A	507	GOL	O1-C1-C2-O2
3	A	502[A]	1K9	CAN-CAY-CBC-CAM
2	B	501	NDP	C3B-C4B-C5B-O5B
3	A	502[A]	1K9	CAN-CAY-CBC-CBB
2	A	501	NDP	C5B-O5B-PA-O1A
2	A	501	NDP	C5D-O5D-PN-O3

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	A	501	NDP	C5D-O5D-PN-O2N
2	B	501	NDP	O4D-C1D-N1N-C6N
2	A	501	NDP	C2D-C1D-N1N-C6N
2	B	501	NDP	O4D-C1D-N1N-C2N
2	A	501	NDP	C2D-C1D-N1N-C2N
2	B	501	NDP	C2D-C1D-N1N-C2N
2	A	501	NDP	O4D-C1D-N1N-C2N
2	A	501	NDP	C3B-C4B-C5B-O5B
2	B	501	NDP	O4B-C4B-C5B-O5B
3	A	502[B]	1K9	CAN-CAY-CBC-CBB
2	A	501	NDP	O4D-C1D-N1N-C6N



There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	NDP	1	0
3	A	502[B]	1K9	2	0
2	A	501	NDP	1	0
5	A	506	GOL	3	0
5	A	504	GOL	1	0
3	A	502[A]	1K9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	416/416 (100%)	-0.28	5 (1%) 76 83	5, 13, 28, 36	13 (3%)
1	B	406/416 (97%)	0.89	79 (19%) 3 3	7, 25, 54, 71	6 (1%)
All	All	822/832 (98%)	0.30	84 (10%) 12 14	5, 16, 47, 71	19 (2%)

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	448	ALA	5.3
1	B	75	VAL	4.7
1	B	445	LEU	4.3
1	B	132	TRP	4.2
1	B	394	PHE	4.2
1	B	391	LEU	3.9
1	B	46	VAL	3.8
1	B	44	ILE	3.8
1	B	72	LEU	3.8
1	B	392	ILE	3.7
1	B	47	ALA	3.7
1	B	438	LEU	3.6
1	B	49	PRO	3.5
1	B	395	ALA	3.5
1	B	385	LEU	3.4
1	B	65	PHE	3.4
1	B	397	MET	3.4
1	B	408	SER	3.4
1	B	383	GLY	3.4
1	B	401	VAL	3.3
1	B	444	ASN	3.3
1	B	128	LEU	3.2
1	B	50	VAL	3.2
1	B	446	ASP	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	427	LEU	3.1
1	B	437	PHE	3.1
1	B	441	ILE	3.1
1	B	443	SER	3.1
1	B	440	THR	3.0
1	B	405	THR	3.0
1	B	398	LEU	3.0
1	B	63	TRP	2.9
1	B	406	VAL	2.8
1	B	66	ILE	2.8
1	B	390	ASP	2.8
1	B	126	PHE	2.8
1	B	77	ILE	2.8
1	B	379	LEU	2.8
1	B	93	ASP	2.7
1	B	62	ILE	2.7
1	B	432	LEU	2.6
1	B	92	THR	2.6
1	B	109	VAL	2.6
1	B	121	ALA	2.6
1	B	61	ILE	2.6
1	B	73	PRO	2.6
1	B	410	ALA	2.5
1	A	456	HIS	2.5
1	B	435	THR	2.5
1	B	141	ASN	2.5
1	A	427	LEU	2.5
1	B	70	LEU	2.5
1	B	423	SER	2.5
1	B	419	ILE	2.5
1	B	123	VAL	2.5
1	B	71	ILE	2.4
1	B	409	GLY	2.4
1	B	87	PRO	2.3
1	B	447	ARG	2.3
1	B	442	LYS	2.3
1	B	54	ASP	2.3
1	A	422	LEU	2.3
1	B	108	SER	2.3
1	B	83	ASP	2.3
1	A	91	GLN	2.3
1	B	393	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	340	LYS	2.3
1	B	79	LEU	2.3
1	B	64	GLN	2.2
1	B	373	PHE	2.2
1	B	130	LYS	2.2
1	B	43	ARG	2.2
1	B	436	ASP	2.2
1	B	78	GLN	2.2
1	B	84	LEU	2.1
1	B	86	LEU	2.1
1	B	378	GLY	2.1
1	B	431	PHE	2.1
1	B	353	ARG	2.1
1	B	388	ASN	2.1
1	B	381	HIS	2.1
1	B	82	PHE	2.0
1	A	425	VAL	2.0
1	B	426	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	A	507	6/6	0.84	0.14	21,24,28,33	0
5	GOL	A	505	6/6	0.88	0.13	30,36,41,44	0
4	CA	A	503	1/1	0.91	0.08	21,21,21,21	0
5	GOL	A	504	6/6	0.91	0.12	17,17,32,34	0
5	GOL	A	506	6/6	0.92	0.09	28,30,33,35	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CA	B	502	1/1	0.95	0.07	20,20,20,20	0
2	NDP	B	501	48/48	0.97	0.06	15,19,26,28	0
3	1K9	A	502[A]	32/32	0.97	0.06	4,7,11,12	32
3	1K9	A	502[B]	32/32	0.97	0.06	6,12,15,17	32
2	NDP	A	501	48/48	0.99	0.04	7,10,14,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 1K9 A 502 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 1K9 A 502 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NDP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.