



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 25, 2026 – 02:44 PM UTC

PDB ID : 1JBA / pdb_00001jba
Title : UNMYRISTOYLATED GCAP-2 WITH THREE CALCIUM IONS BOUND
Authors : Ames, J.B.; Dizhoor, A.M.; Ikura, M.; Palczewski, K.; Stryer, L.
Deposited on : 1999-04-03

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	204	

2 Ensemble composition and analysis

This entry contains 22 models. The atoms present in the NMR models are not consistent. Some calculations may have failed as a result. All residues are included in the validation scores. Model 19 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:19-A:131, A:144-A:186 (156)	1.48	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 2 single-model clusters were found.

Cluster number	Models
1	4, 5, 6, 7, 13, 16, 17, 18, 19, 21, 22
2	9, 14, 20
3	3, 12
4	1, 15
5	8, 10
Single-model clusters	2; 11

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3033 atoms, of which 1489 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PROTEIN (GUANYLATE CYCLASE ACTIVATING PROTEIN 2).

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	189	3030	980	1489	254	299	8	0

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	
2	A	3	Total	Ca
			3	3

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: PROTEIN (GUANYLATE CYCLASE ACTIVATING PROTEIN 2)



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 19. Colouring as in section 4.1 above.

- Molecule 1: PROTEIN (GUANYLATE CYCLASE ACTIVATING PROTEIN 2)



N189	P190
SER	SER
TRP	TRP
ILE	ILE
SER	SER
GLN	GLN
ARG	ARG
LYS	LYS
SER	SER
ALA	ALA
MET	MET
PHE	PHE

5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *distance geometry*.

Of the 50 calculated structures, 22 were deposited, based on the following criterion: *LEAST RESTRAINT VIOLATION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.1

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.44±0.01	7±1/1319 (0.5± 0.0%)	1.44±0.01	12±1/1777 (0.7± 0.1%)
All	All	1.44	158/29018 (0.5%)	1.44	267/39094 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	7.4±0.9
All	All	0	162

5 of 11 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	95	HIS	CG-ND1	-12.29	1.24	1.38	21	22
1	A	43	HIS	CG-ND1	-11.95	1.25	1.38	7	22
1	A	51	VAL	CA-CB	8.06	1.59	1.53	7	1
1	A	146	PRO	N-CA	-7.63	1.37	1.47	4	1
1	A	27	TRP	CG-CD2	-7.42	1.30	1.43	15	22

5 of 20 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	43	HIS	CE1-NE2-CD2	-7.64	101.36	109.00	22	22
1	A	95	HIS	CE1-NE2-CD2	-7.62	101.38	109.00	21	22
1	A	27	TRP	NE1-CE2-CZ2	7.61	141.51	130.10	22	22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	99	TRP	NE1-CE2-CZ2	6.94	140.51	130.10	12	22
1	A	27	TRP	CG-CD2-CE3	-6.83	127.07	133.90	22	22

There are no chirality outliers.

5 of 8 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	175	ARG	Sidechain	22
1	A	108	ARG	Sidechain	21
1	A	114	ARG	Sidechain	21
1	A	176	ARG	Sidechain	21
1	A	66	ARG	Sidechain	20

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1292	1269	1269	108±12
All	All	28489	27918	27917	2385

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 42.

5 of 1029 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:117:LEU:HD13	1:A:118:LEU:N	1.06	1.65	12	1
1:A:93:LEU:H	1:A:93:LEU:HD23	0.97	1.19	12	3
1:A:89:LEU:HD12	1:A:90:ARG:N	0.95	1.74	3	3
1:A:79:LEU:HD22	1:A:79:LEU:H	0.92	1.19	9	10
1:A:68:PHE:CZ	1:A:84:ALA:HB2	0.90	2.01	9	9

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/204 (76%)	107±5 (69±3%)	35±5 (22±3%)	14±3 (9±2%)	1	11
All	All	3432/4488 (76%)	2360 (69%)	766 (22%)	306 (9%)	1	11

5 of 65 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	95	HIS	18
1	A	20	ASP	16
1	A	108	ARG	16
1	A	90	ARG	12
1	A	186	MET	12

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/178 (78%)	112±3 (81±2%)	26±3 (19±2%)	3	34
All	All	3058/3916 (78%)	2475 (81%)	583 (19%)	3	34

5 of 113 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	103	ILE	21
1	A	124	ILE	20
1	A	87	LEU	18
1	A	160	ASN	17
1	A	70	THR	15

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers ⓘ

There are no such molecules in this entry.

6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided