



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 10:56 AM UTC

PDB ID : 3JB3 / pdb_00003jb3
EMDB ID : EMD-6377
Title : Atomic model of cytoplasmic polyhedrosis virus with SAM, GTP and ATP
Authors : Yu, X.K.; Jiang, J.S.; Sun, J.C.; Zhou, Z.H.
Deposited on : 2015-07-06
Resolution : 3.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14724 (2.60 - 3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1058	11% 69% 26% 5%
2	B	1333	61% 25% 11%
2	C	1333	63% 26% 6%
3	D	448	46% 18% 35%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	448	 A horizontal bar chart showing the quality of chain E. The bar is divided into three segments: green (47%), yellow (17%), and grey (35%). A red dot is at the start of the green segment, and a black dot is at the end of the grey segment.

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 32368 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Structural protein VP3.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1057	Total	C	N	O	S	0	0
			8434	5345	1457	1587	45		


- Molecule 2 is a protein called Capsid protein VP1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	1191	Total	C	N	O	S	0	0
			9397	5937	1634	1789	37		
2	C	1251	Total	C	N	O	S	0	0
			9857	6222	1713	1884	38		

- Molecule 3 is a protein called Viral structural protein 5.


Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	292	Total	C	N	O	S	0	0
			2281	1449	399	425	8		
3	E	292	Total	C	N	O	S	0	0
			2281	1449	399	425	8		

- Molecule 4 is S-ADENOSYLMETHIONINE (CCD ID: SAM) (formula: C₁₅H₂₂N₆O₅S).



Mol	Chain	Residues	Atoms					AltConf
4	A	1	Total 27	C 15	N 6	O 5	S 1	0
4	A	1	Total 27	C 15	N 6	O 5	S 1	0

- Molecule 5 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $\text{C}_{10}\text{H}_{16}\text{N}_5\text{O}_{13}\text{P}_3$).



Mol	Chain	Residues	Atoms					AltConf
5	A	1	Total 31	C 10	N 5	O 13	P 3	0

- Molecule 6 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					AltConf
6	A	1	Total	C	N	O	P	0
			32	10	5	14	3	


- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		AltConf
7	A	1	Total	Mg	0
			1	1	

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Structural protein VP3







• Molecule 2: Capsid protein VP1


Chain B: 61% 25% 11%






• Molecule 2: Capsid protein VP1







4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	40898	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	Each particle	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	25	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	60535	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	25.770	Depositor
Minimum map value	-14.720	Depositor
Average map value	0.116	Depositor
Map value standard deviation	1.599	Depositor
Recommended contour level	4	Depositor
Map size ($\text{\AA}$)	772.8, 772.8, 772.8	wwPDB
Map dimensions	700, 700, 700	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.104, 1.104, 1.104	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, MG, GTP, SAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/8619	0.83	13/11737 (0.1%)
2	B	0.46	0/9590	0.89	23/13056 (0.2%)
2	C	0.46	0/10058	0.88	23/13695 (0.2%)
3	D	0.41	0/2327	0.84	3/3163 (0.1%)
3	E	0.42	0/2327	0.84	3/3163 (0.1%)
All	All	0.43	0/32921	0.86	65/44814 (0.1%)

There are no bond length outliers.

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	900	GLY	N-CA-C	8.85	122.10	111.93
2	C	159	ASP	CA-C-N	7.99	128.01	119.78
2	C	159	ASP	C-N-CA	7.99	128.01	119.78
2	C	229	GLN	N-CA-C	7.75	119.80	111.36
2	B	1005	LEU	N-CA-C	7.29	119.02	111.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8434	0	8398	180	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	9397	0	9315	205	0
2	C	9857	0	9767	204	0
3	D	2281	0	2282	50	0
3	E	2281	0	2282	36	0
4	A	54	0	44	3	0
5	A	31	0	12	3	0
6	A	32	0	12	3	0
7	A	1	0	0	0	0
All	All	32368	0	32112	655	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 655 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:267:THR:HG22	1:A:271:ARG:HE	1.41	0.83
2:C:873:TYR:HB3	2:C:898:GLN:HB2	1.64	0.79
1:A:305:TYR:O	1:A:317:ARG:NH1	2.16	0.78
2:B:1144:ARG:NH1	2:B:1170:ASP:OD1	2.17	0.78
5:A:1103:ATP:H5'1	5:A:1103:ATP:H8	1.50	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1055/1058 (100%)	1002 (95%)	53 (5%)	0	100	100
2	B	1187/1333 (89%)	1126 (95%)	58 (5%)	3 (0%)	36	67
2	C	1247/1333 (94%)	1191 (96%)	49 (4%)	7 (1%)	21	52

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	D	290/448 (65%)	282 (97%)	6 (2%)	2 (1%)	18	49
3	E	290/448 (65%)	282 (97%)	7 (2%)	1 (0%)	36	67
All	All	4069/4620 (88%)	3883 (95%)	173 (4%)	13 (0%)	37	67

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	769	GLN
2	C	1251	VAL
3	D	61	ASN
2	B	1310	ARG
2	C	340	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	942/943 (100%)	834 (88%)	108 (12%)	5	23
2	B	1038/1153 (90%)	932 (90%)	106 (10%)	7	28
2	C	1090/1153 (94%)	953 (87%)	137 (13%)	4	19
3	D	240/379 (63%)	219 (91%)	21 (9%)	9	33
3	E	240/379 (63%)	214 (89%)	26 (11%)	6	25
All	All	3550/4007 (89%)	3152 (89%)	398 (11%)	8	24

5 of 398 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	221	LEU
2	C	776	ARG
2	C	270	THR
2	C	532	ILE
2	C	971	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	1100	GLN
3	D	14	GLN
2	C	308	ASN
3	D	100	ASN
2	C	1248	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SAM	A	1101	-	27,29,29	1.09	3 (11%)	34,42,42	2.03	9 (26%)
4	SAM	A	1102	-	27,29,29	1.11	4 (14%)	34,42,42	1.97	10 (29%)
6	GTP	A	1104	7	33,34,34	1.13	3 (9%)	50,54,54	1.64	9 (18%)
5	ATP	A	1103	-	32,33,33	1.41	6 (18%)	48,52,52	1.73	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SAM	A	1101	-	-	9/17/33/33	0/3/3/3
4	SAM	A	1102	-	-	5/17/33/33	0/3/3/3
6	GTP	A	1104	7	-	3/22/38/38	0/3/3/3
5	ATP	A	1103	-	-	5/22/38/38	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1103	ATP	C5-C4	4.59	1.47	1.39
6	A	1104	GTP	PA-O3A	3.13	1.62	1.59
6	A	1104	GTP	PB-O3A	2.99	1.62	1.59
4	A	1102	SAM	C2-N1	2.61	1.38	1.33
4	A	1101	SAM	C2-N3	2.60	1.38	1.33

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1101	SAM	N3-C2-N1	-5.66	120.02	128.58
4	A	1102	SAM	N3-C2-N1	-5.52	120.23	128.58
5	A	1103	ATP	C5-C4-N3	-5.48	119.17	126.72
6	A	1104	GTP	C5-C4-N3	-4.86	120.66	128.39
6	A	1104	GTP	C2-N3-C4	4.66	120.32	112.30

There are no chirality outliers.

5 of 22 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	A	1101	SAM	O-C-CA-N
4	A	1101	SAM	C4'-C5'-SD-CE
4	A	1102	SAM	N-CA-CB-CG
4	A	1102	SAM	C-CA-CB-CG
4	A	1102	SAM	CB-CG-SD-C5'


There are no ring outliers.


4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1101	SAM	2	0
4	A	1102	SAM	1	0
6	A	1104	GTP	3	0
5	A	1103	ATP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-6377. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 350



Y Index: 350




Z Index: 350


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 373



Y Index: 397




Z Index: 279


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views

This section was not generated.


6.6 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 10082 nm³; this corresponds to an approximate mass of 9107 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹

8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-6377 and PDB model 3JB3. Per-residue inclusion information can be found in section 3 on page 7.

9.1 Map-model overlay [i](#)

This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 75% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (4) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7500	0.5310
A	0.6710	0.5050
B	0.7700	0.5390
C	0.7850	0.5410
D	0.7880	0.5460
E	0.7760	0.5350

