



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 02:43 AM UTC

PDB ID : 4JB9 / pdb_00004jb9
Title : Crystal structure of antibody VRC06 in complex with HIV-1 gp120 core
Authors : Kwon, Y.D.; Zhou, T.; Srivatsan, S.; Kwong, P.D.
Deposited on : 2013-02-19
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	353	2% 84% 13% ..
2	H	232	7% 78% 19% ..
3	L	209	20% 77% 21% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called clade A/E 93TH057 HIV-1 gp120 core.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	344	Total	C	N	O	S	0	0	0
			2693	1689	469	512	23			

- Molecule 2 is a protein called antibody VRC06 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	225	Total	C	N	O	S	0	0	0
			1746	1113	301	322	10			

- Molecule 3 is a protein called antibody VRC06 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	209	Total	C	N	O	S	0	0	0
			1618	1011	276	326	5			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0
4	G	1	Total C N O 14 8 1 5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	G	11	Total O 11 11	0	0
5	H	17	Total O 17 17	0	0
5	L	12	Total O 12 12	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: clade A/E 93TH057 HIV-1 gp120 core



- Molecule 2: antibody VRC06 heavy chain



- Molecule 3: antibody VRC06 light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.24Å 68.92Å 224.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.72 – 2.60 48.72 – 2.60	Depositor EDS
% Data completeness (in resolution range)	92.0 (48.72-2.60) 92.0 (48.72-2.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.247 , 0.287 0.252 , 0.292	Depositor DCC
R_{free} test set	1545 reflections (4.41%)	wwPDB-VP
Wilson B-factor (Å ²)	62.0	Xtriage
Anisotropy	0.573	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 69.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6237	wwPDB-VP
Average B, all atoms (Å ²)	109.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	G	0.30	0/2749	0.74	3/3730 (0.1%)
2	H	0.32	0/1798	0.73	1/2443 (0.0%)
3	L	0.34	0/1651	0.77	2/2239 (0.1%)
All	All	0.32	0/6198	0.75	6/8412 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	90	PHE	CB-CA-C	-5.64	110.08	116.63
1	G	65	VAL	N-CA-C	5.26	115.88	110.36
3	L	40	GLY	N-CA-C	-5.05	108.68	114.69
1	G	211	ASP	CA-C-N	5.02	124.93	119.76
1	G	211	ASP	C-N-CA	5.02	124.93	119.76
2	H	152	VAL	N-CA-C	5.00	115.17	107.77

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2693	0	2625	29	0
2	H	1746	0	1687	33	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	L	1618	0	1575	40	0
4	G	140	0	129	1	0
5	G	11	0	0	2	0
5	H	17	0	0	8	0
5	L	12	0	0	2	0
All	All	6237	0	6016	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:133:ASN:OD1	3:L:167:THR:HB	1.70	0.90
1:G:95:MET:HE1	1:G:273:ARG:HB3	1.60	0.82
2:H:196:CYS:O	5:H:309:HOH:O	1.97	0.81
2:H:141:LEU:HD12	3:L:128:VAL:HG21	1.62	0.80
3:L:84:VAL:HG11	3:L:86:TYR:CZ	2.16	0.80
1:G:229:ASN:ND2	4:G:502:NAG:O6	2.15	0.79
2:H:198:VAL:N	5:H:308:HOH:O	2.18	0.77
3:L:108:PRO:HB3	3:L:134:PHE:HB2	1.71	0.72
3:L:84:VAL:HG12	3:L:86:TYR:CE1	2.26	0.69
3:L:37:LYS:HD2	3:L:43:PRO:HG3	1.73	0.69
3:L:133:ASN:OD1	3:L:167:THR:CB	2.40	0.68
2:H:141:LEU:CD1	3:L:128:VAL:HG21	2.25	0.66
1:G:466:GLU:OE1	2:H:61:ARG:NH2	2.30	0.65
3:L:37:LYS:NZ	3:L:41:GLN:O	2.30	0.65
2:H:100(A):CYS:HB2	2:H:100(E):HIS:CD2	2.33	0.64
3:L:84:VAL:CG1	3:L:86:TYR:CE1	2.82	0.63
3:L:84:VAL:CG1	3:L:86:TYR:CZ	2.81	0.62
1:G:465:ASN:O	2:H:61:ARG:NH1	2.32	0.62
3:L:84:VAL:HG11	3:L:86:TYR:OH	2.00	0.61
1:G:91:GLU:N	5:G:603:HOH:O	2.23	0.61
1:G:207:LYS:HE2	1:G:439:ILE:HG23	1.83	0.61
2:H:28:ASN:ND2	2:H:31:ASP:OD1	2.34	0.60
2:H:95:ARG:HD3	2:H:100(C):ASP:HA	1.84	0.60
2:H:168:ALA:HA	2:H:178:LEU:HB3	1.84	0.60
1:G:95:MET:HE2	1:G:235:GLY:HA3	1.84	0.59
2:H:193:THR:HG23	2:H:210:LYS:HE3	1.86	0.57
1:G:357:LYS:HD2	1:G:466:GLU:HG2	1.88	0.56
1:G:231:LYS:HE2	1:G:267:GLU:HB3	1.88	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:108:PRO:HD2	3:L:196:LEU:HD13	1.88	0.55
1:G:101:VAL:HG21	1:G:480:ARG:HG2	1.88	0.55
2:H:66:ARG:NH2	2:H:86:ASP:OD2	2.40	0.55
2:H:119:PRO:HB3	2:H:145:TYR:HB3	1.88	0.54
3:L:47:ILE:HG23	3:L:52:ARG:H	1.72	0.54
1:G:336:THR:HG22	1:G:411:ASN:HB2	1.88	0.54
3:L:10:THR:HG22	3:L:98:THR:HB	1.89	0.54
3:L:142:GLN:HB3	3:L:190:GLU:HB3	1.88	0.54
1:G:65:VAL:HG21	1:G:208:ILE:HD12	1.92	0.52
3:L:13:LEU:HD12	3:L:99:LEU:HG	1.90	0.52
1:G:198:GLY:H	2:H:76(A):LEU:HD13	1.75	0.52
2:H:141:LEU:CD1	3:L:128:VAL:CG2	2.88	0.51
2:H:153:SER:O	5:H:307:HOH:O	2.19	0.51
3:L:133:ASN:HA	3:L:168:TYR:HB2	1.92	0.51
2:H:41:ARG:NH1	5:H:304:HOH:O	2.43	0.51
3:L:134:PHE:CZ	3:L:139:ALA:HB2	2.45	0.51
3:L:129:CYS:SG	3:L:172:SER:HB3	2.51	0.50
3:L:46:LEU:HA	3:L:57:ILE:HD13	1.94	0.50
2:H:52(A):PRO:O	2:H:71:ARG:NH1	2.45	0.49
2:H:45:PHE:O	5:H:301:HOH:O	2.18	0.49
1:G:265:LEU:HD21	1:G:291:SER:HB3	1.94	0.49
3:L:103:ARG:HA	3:L:104:THR:O	2.13	0.49
3:L:158:VAL:O	5:L:304:HOH:O	2.20	0.48
1:G:453:LEU:HD13	1:G:472:GLY:HA3	1.95	0.48
2:H:197:ASN:N	5:H:307:HOH:O	2.46	0.48
2:H:152:VAL:HG11	2:H:180:SER:OG	2.14	0.48
3:L:36:GLN:HB2	3:L:46:LEU:HD11	1.95	0.47
1:G:292:VAL:HG21	1:G:338:TRP:HE3	1.80	0.47
3:L:13:LEU:HD12	3:L:99:LEU:CG	2.44	0.47
3:L:49:ASP:O	3:L:51:SER:N	2.43	0.46
3:L:62:VAL:O	3:L:73:THR:N	2.45	0.46
1:G:333:ILE:HD12	1:G:390:LEU:HD21	1.98	0.46
1:G:439:ILE:HG13	1:G:443:ILE:HD11	1.97	0.46
1:G:122:LEU:HD11	1:G:431:GLY:HA3	1.98	0.45
3:L:108:PRO:HB3	3:L:134:PHE:CB	2.44	0.45
2:H:39:PHE:N	2:H:89:ASP:O	2.48	0.45
1:G:477:ASP:OD1	1:G:480:ARG:NH1	2.49	0.45
1:G:438:PRO:O	5:G:602:HOH:O	2.20	0.45
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.53	0.43
3:L:10:THR:O	5:L:312:HOH:O	2.20	0.43
2:H:47:TRP:HB2	3:L:93:PHE:CE1	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:134:PHE:CE2	3:L:139:ALA:HB2	2.54	0.43
1:G:273:ARG:HB2	1:G:285:ILE:HB	2.00	0.43
3:L:48:TYR:CE1	3:L:52:ARG:HD2	2.54	0.42
3:L:193:HIS:CE1	3:L:195:GLY:H	2.36	0.42
2:H:83:THR:OG1	2:H:84:SER:N	2.51	0.42
1:G:407:MET:HG3	1:G:408:LYS:HG3	2.01	0.42
3:L:84:VAL:HG12	3:L:85:TYR:N	2.34	0.41
1:G:219:THR:HG23	1:G:225:ILE:HG13	2.02	0.41
1:G:343:LYS:HD3	1:G:396:ILE:O	2.21	0.41
2:H:154:TRP:NE1	5:H:316:HOH:O	2.35	0.41
2:H:6:GLU:H	2:H:105:GLN:HE22	1.68	0.41
3:L:18:ARG:HH11	3:L:73:THR:HG21	1.85	0.41
2:H:167:PRO:HD2	3:L:157:SER:OG	2.21	0.41
2:H:5:VAL:O	2:H:23:ALA:N	2.52	0.41
3:L:3:VAL:HB	3:L:26:SER:HB3	2.02	0.41
3:L:196:LEU:HD12	3:L:196:LEU:HA	1.85	0.41
3:L:33:ASN:O	3:L:87:CYS:HA	2.21	0.41
2:H:35:HIS:HB2	2:H:93:VAL:HG23	2.03	0.40
1:G:95:MET:HE3	1:G:96:TRP:CE2	2.57	0.40
2:H:152:VAL:HG13	5:H:312:HOH:O	2.21	0.40
1:G:211:ASP:HA	1:G:212:PRO:HD2	1.92	0.40
1:G:416:LEU:HA	1:G:417:PRO:HD3	1.98	0.40
2:H:146:PHE:HA	2:H:147:PRO:HA	1.81	0.40
1:G:121:LYS:HE3	1:G:426:MET:HE1	2.02	0.40
2:H:214:LYS:HB2	2:H:214:LYS:HE3	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	338/353 (96%)	314 (93%)	24 (7%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	221/232 (95%)	199 (90%)	21 (10%)	1 (0%)	24	46
3	L	207/209 (99%)	182 (88%)	18 (9%)	7 (3%)	3	4
All	All	766/794 (96%)	695 (91%)	63 (8%)	8 (1%)	12	28

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	L	134	PHE
2	H	99	PRO
3	L	103	ARG
3	L	104	THR
3	L	106	ALA
3	L	15	PRO
3	L	105	VAL
3	L	158	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	307/311 (99%)	304 (99%)	3 (1%)	68	86
2	H	193/197 (98%)	192 (100%)	1 (0%)	81	92
3	L	183/183 (100%)	178 (97%)	5 (3%)	39	67
All	All	683/691 (99%)	674 (99%)	9 (1%)	61	82

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	120	VAL
1	G	255	VAL
1	G	328	LYS
2	H	71	ARG
3	L	99	LEU
3	L	104	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	L	134	PHE
3	L	158	VAL
3	L	196	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	389	GLN
2	H	164	HIS
3	L	88	GLN
3	L	184	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	G	501	1	14,14,15	0.31	0	17,19,21	0.40	0
4	NAG	G	504	1	14,14,15	0.60	0	17,19,21	0.46	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	G	506	1	14,14,15	0.31	0	17,19,21	0.37	0
4	NAG	G	507	1	14,14,15	0.43	0	17,19,21	0.39	0
4	NAG	G	510	1	14,14,15	0.21	0	17,19,21	0.48	0
4	NAG	G	502	1	14,14,15	1.08	1 (7%)	17,19,21	0.70	0
4	NAG	G	503	1	14,14,15	0.31	0	17,19,21	0.57	0
4	NAG	G	508	1	14,14,15	0.39	0	17,19,21	0.33	0
4	NAG	G	505	1	14,14,15	0.28	0	17,19,21	0.41	0
4	NAG	G	509	1	14,14,15	0.21	0	17,19,21	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	501	1	-	0/6/23/26	0/1/1/1
4	NAG	G	504	1	-	0/6/23/26	0/1/1/1
4	NAG	G	506	1	-	0/6/23/26	0/1/1/1
4	NAG	G	507	1	-	0/6/23/26	0/1/1/1
4	NAG	G	510	1	-	2/6/23/26	0/1/1/1
4	NAG	G	502	1	-	2/6/23/26	0/1/1/1
4	NAG	G	503	1	-	1/6/23/26	0/1/1/1
4	NAG	G	508	1	-	4/6/23/26	0/1/1/1
4	NAG	G	505	1	-	0/6/23/26	0/1/1/1
4	NAG	G	509	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	502	NAG	C1-C2	3.76	1.57	1.52

There are no bond angle outliers.

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	G	508	NAG	O5-C5-C6-O6
4	G	502	NAG	O5-C5-C6-O6
4	G	502	NAG	C4-C5-C6-O6
4	G	508	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	G	508	NAG	C8-C7-N2-C2
4	G	508	NAG	O7-C7-N2-C2
4	G	509	NAG	C8-C7-N2-C2
4	G	509	NAG	O7-C7-N2-C2
4	G	510	NAG	C8-C7-N2-C2
4	G	510	NAG	O7-C7-N2-C2
4	G	503	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	502	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	344/353 (97%)	0.21	6 (1%) 69 64	37, 78, 149, 191	0
2	H	225/232 (96%)	0.62	17 (7%) 20 16	51, 107, 180, 209	0
3	L	209/209 (100%)	1.13	42 (20%) 3 2	72, 150, 187, 205	0
All	All	778/794 (97%)	0.58	65 (8%) 17 13	37, 104, 179, 209	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	159	LEU	4.3
3	L	88	GLN	4.2
3	L	13	LEU	4.1
3	L	133	ASN	4.0
3	L	134	PHE	3.8
2	H	182	VAL	3.8
3	L	102	ASN	3.6
3	L	168	TYR	3.6
1	G	472	GLY	3.5
3	L	2	ILE	3.5
3	L	94	GLY	3.5
3	L	11	LEU	3.4
3	L	72	LEU	3.3
2	H	158	ALA	3.3
2	H	157	GLY	3.2
1	G	123	THR	3.0
3	L	132	ASN	3.0
3	L	21	LEU	2.9
3	L	99	LEU	2.9
3	L	104	THR	2.9
2	H	164	HIS	2.9
3	L	47	ILE	2.9
3	L	34	TRP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	L	135	TYR	2.8
3	L	105	VAL	2.8
2	H	141	LEU	2.8
3	L	93	PHE	2.7
1	G	360	ILE	2.7
3	L	74	ILE	2.7
3	L	159	THR	2.7
1	G	124	GLY	2.7
3	L	8	PRO	2.6
2	H	100(C)	ASP	2.5
3	L	101	ILE	2.5
3	L	103	ARG	2.5
3	L	35	TYR	2.5
1	G	122	LEU	2.5
3	L	107	ALA	2.5
1	G	396	ILE	2.4
3	L	32	LEU	2.4
3	L	23	CYS	2.4
3	L	169	SER	2.4
2	H	181	VAL	2.4
3	L	61	PHE	2.4
3	L	106	ALA	2.3
2	H	100(E)	HIS	2.3
2	H	162	GLY	2.3
2	H	33	SER	2.3
2	H	121	VAL	2.3
3	L	71	SER	2.3
2	H	195	ILE	2.2
3	L	158	VAL	2.2
3	L	75	THR	2.2
2	H	194	TYR	2.2
3	L	161	GLN	2.2
3	L	163	SER	2.2
3	L	137	ARG	2.2
3	L	128	VAL	2.2
3	L	4	LEU	2.1
2	H	139	GLY	2.1
2	H	184	VAL	2.1
3	L	70	PHE	2.1
2	H	137	ALA	2.1
3	L	48	TYR	2.1
3	L	67	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	G	502	14/15	0.27	0.25	150,162,176,184	0
4	NAG	G	510	14/15	0.38	0.14	141,150,165,169	0
4	NAG	G	509	14/15	0.44	0.12	135,146,150,151	0
4	NAG	G	504	14/15	0.57	0.17	130,149,193,200	0
4	NAG	G	501	14/15	0.69	0.11	66,112,139,139	0
4	NAG	G	505	14/15	0.78	0.10	76,104,116,122	0
4	NAG	G	508	14/15	0.81	0.11	80,91,98,108	0
4	NAG	G	507	14/15	0.82	0.14	111,132,143,143	0
4	NAG	G	506	14/15	0.84	0.15	83,107,130,134	0
4	NAG	G	503	14/15	0.93	0.08	47,64,69,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.