



wwPDB EM Validation Summary Report ⓘ

Mar 6, 2026 – 02:48 PM UTC

PDB ID : 8JB0 / pdb_00008jb0
EMDB ID : EMD-36139
Title : Cryo-EM structure of Holo form of ScBfr in C1 symmetry
Authors : Jobichen, C.; Sivaraman, J.
Deposited on : 2023-05-07
Resolution : 4.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	5410 (3.70 - 4.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	 66% 31% .
1	B	167	 64% 32% .
1	C	167	 69% 28% .
1	D	167	 7% 67% 29% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	167	
1	F	167	
1	G	167	
1	H	167	
1	I	167	
1	J	167	
1	K	167	
1	L	167	
1	M	167	
1	N	167	
1	O	167	
1	P	167	
1	Q	167	
1	R	167	
1	S	167	
1	T	167	
1	U	167	
1	V	167	
1	W	167	
1	X	167	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 31387 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacterioferritin.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	162	Total	C	N	O	S	0	0
			1316	828	227	257	4		
1	B	160	Total	C	N	O	S	0	0
			1293	814	224	251	4		
1	C	162	Total	C	N	O	S	0	0
			1318	832	226	256	4		
1	D	161	Total	C	N	O	S	0	0
			1305	823	225	253	4		
1	F	161	Total	C	N	O	S	0	0
			1310	826	225	255	4		
1	E	161	Total	C	N	O	S	0	0
			1304	823	225	252	4		
1	G	161	Total	C	N	O	S	0	0
			1304	823	225	252	4		
1	H	161	Total	C	N	O	S	0	0
			1304	823	225	252	4		
1	I	160	Total	C	N	O	S	0	0
			1299	820	224	251	4		
1	J	162	Total	C	N	O	S	0	0
			1318	831	227	256	4		
1	K	157	Total	C	N	O	S	0	0
			1285	811	221	249	4		
1	L	162	Total	C	N	O	S	0	0
			1318	832	226	256	4		
1	M	162	Total	C	N	O	S	0	0
			1310	826	226	254	4		
1	N	161	Total	C	N	O	S	0	0
			1310	826	225	255	4		
1	O	162	Total	C	N	O	S	0	0
			1315	828	227	256	4		
1	P	161	Total	C	N	O	S	0	0
			1305	823	225	253	4		
1	Q	162	Total	C	N	O	S	0	0
			1318	832	226	256	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	157	Total	C	N	O	S	0	0
			1285	811	221	249	4		
1	S	161	Total	C	N	O	S	0	0
			1310	826	225	255	4		
1	T	162	Total	C	N	O	S	0	0
			1315	828	227	256	4		
1	U	157	Total	C	N	O	S	0	0
			1285	811	221	249	4		
1	V	162	Total	C	N	O	S	0	0
			1318	832	226	256	4		
1	W	160	Total	C	N	O	S	0	0
			1296	818	223	251	4		
1	X	161	Total	C	N	O	S	0	0
			1310	826	225	255	4		

- Molecule 2 is FE (II) ION (CCD ID: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		AltConf
2	A	2	Total	Fe	0
			2	2	
2	B	1	Total	Fe	0
			1	1	
2	C	2	Total	Fe	0
			2	2	
2	D	1	Total	Fe	0
			1	1	
2	F	1	Total	Fe	0
			1	1	
2	E	2	Total	Fe	0
			2	2	
2	G	2	Total	Fe	0
			2	2	
2	H	2	Total	Fe	0
			2	2	
2	I	1	Total	Fe	0
			1	1	
2	J	1	Total	Fe	0
			1	1	
2	K	2	Total	Fe	0
			2	2	
2	L	1	Total	Fe	0
			1	1	

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
2	M	2	Total 2	Fe 2	0
2	N	1	Total 1	Fe 1	0
2	O	2	Total 2	Fe 2	0
2	P	1	Total 1	Fe 1	0
2	Q	1	Total 1	Fe 1	0
2	R	2	Total 2	Fe 2	0
2	S	2	Total 2	Fe 2	0
2	T	2	Total 2	Fe 2	0
2	U	1	Total 1	Fe 1	0
2	V	1	Total 1	Fe 1	0
2	W	2	Total 2	Fe 2	0
2	X	1	Total 1	Fe 1	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Bacterioferritin



• Molecule 1: Bacterioferritin




• Molecule 1: Bacterioferritin



• Molecule 1: Bacterioferritin








• Molecule 1: Bacterioferritin




• Molecule 1: Bacterioferritin



• Molecule 1: Bacterioferritin




• Molecule 1: Bacterioferritin






- Molecule 1: Bacterioferritin





- Molecule 1: Bacterioferritin



- Molecule 1: Bacterioferritin




- Molecule 1: Bacterioferritin




- Molecule 1: Bacterioferritin








• Molecule 1: Bacterioferritin




• Molecule 1: Bacterioferritin



• Molecule 1: Bacterioferritin



• Molecule 1: Bacterioferritin






• Molecule 1: Bacterioferritin



• Molecule 1: Bacterioferritin



• Molecule 1: Bacterioferritin



• Molecule 1: Bacterioferritin



• Molecule 1: Bacterioferritin



L123	L124	A125	H129	H130	L134	Q137	L138	D139	L140	T141	K143	L144	L148	Y149	L150	V153	L154	E155	Q156	T157	Q158	P159	D160	P161	S162	GLY	PRO	GLY	SER	LEU	M1	Q2	G3	D4	P5	E6	V7	I8	E9	Q14	L15	Y25	K30	K39	L40	A41	K42	L58	R61	I62	L65	P69	N70	Y71	Q72	V83	T84	E85	M86	F87	Q88	A89	D90	R91	E92	E96	G104	V107	M108	R109	T115	S116	A117	F120	E121	A122
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	11517	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	1.8	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.031	Depositor
Minimum map value	-0.010	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.0079	Depositor
Map size (Å)	219.648, 219.648, 219.648	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.858, 0.858, 0.858	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/1337	0.38	0/1806
1	B	0.17	0/1313	0.36	0/1774
1	C	0.15	0/1341	0.35	0/1814
1	D	0.19	0/1326	0.44	0/1792
1	E	0.17	0/1325	0.37	0/1790
1	F	0.16	0/1332	0.42	0/1801
1	G	0.15	0/1325	0.32	0/1790
1	H	0.21	0/1325	0.45	0/1790
1	I	0.15	0/1320	0.35	0/1784
1	J	0.16	0/1340	0.33	0/1812
1	K	0.16	0/1306	0.34	0/1764
1	L	0.18	0/1341	0.40	0/1814
1	M	0.17	0/1331	0.38	0/1799
1	N	0.18	0/1332	0.44	0/1801
1	O	0.16	0/1336	0.36	0/1805
1	P	0.16	0/1326	0.40	0/1792
1	Q	0.15	0/1341	0.31	0/1814
1	R	0.18	0/1306	0.38	0/1764
1	S	0.15	0/1332	0.34	0/1801
1	T	0.17	0/1336	0.37	0/1805
1	U	0.18	0/1306	0.41	0/1764
1	V	0.15	0/1341	0.34	0/1814
1	W	0.15	0/1317	0.32	0/1780
1	X	0.18	0/1332	0.38	0/1801
All	All	0.17	0/31867	0.38	0/43071

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1316	0	1283	38	0
1	B	1293	0	1260	41	0
1	C	1318	0	1289	35	0
1	D	1305	0	1268	39	0
1	E	1304	0	1268	31	0
1	F	1310	0	1276	51	0
1	G	1304	0	1268	30	0
1	H	1304	0	1268	47	0
1	I	1299	0	1267	32	0
1	J	1318	0	1287	42	0
1	K	1285	0	1262	36	0
1	L	1318	0	1289	49	0
1	M	1310	0	1270	48	0
1	N	1310	0	1276	41	0
1	O	1315	0	1280	34	0
1	P	1305	0	1268	45	0
1	Q	1318	0	1289	31	0
1	R	1285	0	1262	41	0
1	S	1310	0	1276	31	0
1	T	1315	0	1280	46	0
1	U	1285	0	1262	48	0
1	V	1318	0	1289	47	0
1	W	1296	0	1261	31	0
1	X	1310	0	1276	53	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
2	C	2	0	0	0	0
2	D	1	0	0	0	0
2	E	2	0	0	0	0
2	F	1	0	0	0	0
2	G	2	0	0	0	0
2	H	2	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
2	K	2	0	0	0	0
2	L	1	0	0	0	0
2	M	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	N	1	0	0	0	0
2	O	2	0	0	0	0
2	P	1	0	0	0	0
2	Q	1	0	0	0	0
2	R	2	0	0	0	0
2	S	2	0	0	0	0
2	T	2	0	0	0	0
2	U	1	0	0	0	0
2	V	1	0	0	0	0
2	W	2	0	0	0	0
2	X	1	0	0	0	0
All	All	31387	0	30574	851	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 851 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:108:MET:HE1	1:L:116:SER:HB2	1.37	1.01
1:L:151:SER:OG	1:N:148:LEU:HD21	1.67	0.93
1:M:32:GLN:NE2	1:M:40:LEU:HD23	1.85	0.92
1:V:134:LEU:O	1:V:137:GLN:HG2	1.71	0.91
1:V:134:LEU:HD23	1:V:137:GLN:NE2	1.87	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	160/167 (96%)	157 (98%)	3 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	158/167 (95%)	155 (98%)	3 (2%)	0	100	100
1	C	160/167 (96%)	156 (98%)	4 (2%)	0	100	100
1	D	159/167 (95%)	155 (98%)	3 (2%)	1 (1%)	21	58
1	E	159/167 (95%)	154 (97%)	5 (3%)	0	100	100
1	F	159/167 (95%)	154 (97%)	5 (3%)	0	100	100
1	G	159/167 (95%)	153 (96%)	5 (3%)	1 (1%)	21	58
1	H	159/167 (95%)	152 (96%)	6 (4%)	1 (1%)	21	58
1	I	158/167 (95%)	152 (96%)	5 (3%)	1 (1%)	21	58
1	J	160/167 (96%)	154 (96%)	6 (4%)	0	100	100
1	K	155/167 (93%)	153 (99%)	2 (1%)	0	100	100
1	L	160/167 (96%)	153 (96%)	7 (4%)	0	100	100
1	M	160/167 (96%)	155 (97%)	5 (3%)	0	100	100
1	N	159/167 (95%)	149 (94%)	9 (6%)	1 (1%)	21	58
1	O	160/167 (96%)	157 (98%)	3 (2%)	0	100	100
1	P	159/167 (95%)	154 (97%)	5 (3%)	0	100	100
1	Q	160/167 (96%)	159 (99%)	1 (1%)	0	100	100
1	R	155/167 (93%)	151 (97%)	4 (3%)	0	100	100
1	S	159/167 (95%)	156 (98%)	3 (2%)	0	100	100
1	T	160/167 (96%)	154 (96%)	5 (3%)	1 (1%)	21	58
1	U	155/167 (93%)	150 (97%)	5 (3%)	0	100	100
1	V	160/167 (96%)	155 (97%)	5 (3%)	0	100	100
1	W	158/167 (95%)	152 (96%)	6 (4%)	0	100	100
1	X	159/167 (95%)	157 (99%)	2 (1%)	0	100	100
All	All	3810/4008 (95%)	3697 (97%)	107 (3%)	6 (0%)	44	77

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	159	PRO
1	G	159	PRO
1	H	159	PRO
1	I	159	PRO
1	T	161	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/147 (96%)	141 (100%)	0	100	100
1	B	137/147 (93%)	137 (100%)	0	100	100
1	C	142/147 (97%)	142 (100%)	0	100	100
1	D	138/147 (94%)	138 (100%)	0	100	100
1	E	138/147 (94%)	138 (100%)	0	100	100
1	F	140/147 (95%)	140 (100%)	0	100	100
1	G	138/147 (94%)	138 (100%)	0	100	100
1	H	138/147 (94%)	138 (100%)	0	100	100
1	I	138/147 (94%)	138 (100%)	0	100	100
1	J	141/147 (96%)	141 (100%)	0	100	100
1	K	138/147 (94%)	138 (100%)	0	100	100
1	L	142/147 (97%)	141 (99%)	1 (1%)	76	78
1	M	138/147 (94%)	137 (99%)	1 (1%)	76	78
1	N	140/147 (95%)	140 (100%)	0	100	100
1	O	140/147 (95%)	140 (100%)	0	100	100
1	P	138/147 (94%)	138 (100%)	0	100	100
1	Q	142/147 (97%)	141 (99%)	1 (1%)	76	78
1	R	138/147 (94%)	138 (100%)	0	100	100
1	S	140/147 (95%)	140 (100%)	0	100	100
1	T	140/147 (95%)	140 (100%)	0	100	100
1	U	138/147 (94%)	138 (100%)	0	100	100
1	V	142/147 (97%)	142 (100%)	0	100	100
1	W	137/147 (93%)	137 (100%)	0	100	100
1	X	140/147 (95%)	140 (100%)	0	100	100
All	All	3344/3528 (95%)	3341 (100%)	3 (0%)	87	88

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	160	ASP
1	M	129	HIS
1	Q	160	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	U	76	HIS
1	V	28	HIS
1	W	14	GLN
1	K	12	ASN
1	J	81	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 36 ligands modelled in this entry, 36 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-36139. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.3.2 Raw map



X Index: 129



Y Index: 128




Z Index: 121


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0079. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 273 nm³; this corresponds to an approximate mass of 247 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.238 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.238 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.20	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	6.08	8.05	6.54

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.08 differs from the reported value 4.2 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-36139 and PDB model 8JB0. Per-residue inclusion information can be found in section [3](#) on page [7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0079 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0079).

9.4 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0079) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7420	 0.3630
A	 0.7310	 0.3630
B	 0.7420	 0.3680
C	 0.7480	 0.3410
D	 0.7270	 0.3520
E	 0.7830	 0.3570
F	 0.7320	 0.3450
G	 0.7550	 0.3730
H	 0.7080	 0.3510
I	 0.7350	 0.3740
J	 0.7430	 0.3430
K	 0.7490	 0.3660
L	 0.7670	 0.3640
M	 0.7460	 0.3790
N	 0.7320	 0.3570
O	 0.7370	 0.3660
P	 0.7340	 0.3680
Q	 0.7460	 0.3520
R	 0.7380	 0.3530
S	 0.7120	 0.3820
T	 0.7480	 0.3680
U	 0.7470	 0.3610
V	 0.7520	 0.3730
W	 0.7680	 0.3800
X	 0.7240	 0.3680

