



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 08:43 AM UTC

PDB ID : 5JCF / pdb_00005jcf
Title : Crystal structure of chicken MDA5 with 5'p 10-mer dsRNA and ADP-Mg²⁺ at 2.6 Å resolution (orthorhombic form).
Authors : Cusack, S.; Uchikawa, E.
Deposited on : 2016-04-15
Resolution : 2.60 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)
RNA backbone	3983	1014 (2.84-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	701	12% 91% 6%
1	B	701	13% 90% 6%
2	C	10	100%
2	D	10	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	X	10	 100%
3	Y	11	 9%  100%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 11870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Melanoma differentiation associated protein-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	660	Total	C	N	O	S	0	0	0
			5386	3394	964	996	32			
1	B	660	Total	C	N	O	S	0	0	0
			5385	3393	964	996	32			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	294	GLY	-	expression tag	UNP D9N195
A	295	ALA	-	expression tag	UNP D9N195
A	296	MET	-	expression tag	UNP D9N195
A	297	GLY	-	expression tag	UNP D9N195
A	436	GLN	GLU	engineered mutation	UNP D9N195
B	294	GLY	-	expression tag	UNP D9N195
B	295	ALA	-	expression tag	UNP D9N195
B	296	MET	-	expression tag	UNP D9N195
B	297	GLY	-	expression tag	UNP D9N195
B	436	GLN	GLU	engineered mutation	UNP D9N195


- Molecule 2 is a RNA chain called RNA (5'-R(P*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	10	Total	C	N	O	P	0	0	0
			214	95	38	71	10			
2	C	10	Total	C	N	O	P	0	0	0
			214	95	38	71	10			
2	D	10	Total	C	N	O	P	0	0	0
			214	95	38	71	10			

- Molecule 3 is a RNA chain called RNA (5'-R(P*AP*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Y	11	Total	C	N	O	P	0	0	0
			231	105	43	73	10			

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	B	1	Total	Mg	0	0
			1	1		

- Molecule 6 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		
6	B	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	74	Total 74	O 74	0	0
7	X	5	Total 5	O 5	0	0
7	Y	7	Total 7	O 7	0	0
7	B	70	Total 70	O 70	0	0
7	C	5	Total 5	O 5	0	0
7	D	7	Total 7	O 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Melanoma differentiation associated protein-5



- Molecule 1: Melanoma differentiation associated protein-5



- Molecule 2: RNA (5'-R(P*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3')

Chain X:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: RNA (5'-R(P*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3')

Chain C:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: RNA (5'-R(P*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3')

Chain D:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: RNA (5'-R(P*AP*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3')

Chain Y:  9% 100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	101.92Å 132.47Å 139.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.60 50.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.2 (50.00-2.60) 98.2 (50.00-2.60)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.277 , 0.312 0.278 , 0.310	Depositor DCC
R_{free} test set	2901 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	51.3	Xtriage
Anisotropy	0.783	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 28.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11870	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4199e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/5471	0.78	1/7338 (0.0%)
1	B	0.55	1/5470 (0.0%)	0.77	1/7337 (0.0%)
2	C	0.35	0/238	0.66	0/367
2	D	0.35	0/238	0.66	0/367
2	X	0.36	0/238	0.66	0/367
3	Y	0.35	0/258	0.64	0/401
All	All	0.53	1/11913 (0.0%)	0.77	2/16177 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	785	THR	CA-C	5.02	1.54	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	485	VAL	CB-CA-C	5.51	116.16	110.70
1	B	485	VAL	CB-CA-C	5.46	116.11	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5386	0	5463	5	0
1	B	5385	0	5461	5	0
2	C	214	0	109	0	0
2	D	214	0	109	0	0
2	X	214	0	109	0	0
3	Y	231	0	118	0	0
4	A	27	0	12	0	0
4	B	27	0	12	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	74	0	0	0	0
7	B	70	0	0	0	0
7	C	5	0	0	0	0
7	D	7	0	0	0	0
7	X	5	0	0	0	0
7	Y	7	0	0	0	0
All	All	11870	0	11393	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:538:ARG:NH1	1:A:672:ASP:O	2.30	0.64
1:B:538:ARG:NH1	1:B:672:ASP:O	2.30	0.64
1:B:434:ILE:HD11	1:B:450:MET:HE3	1.85	0.59
1:A:434:ILE:HD11	1:A:450:MET:HE3	1.85	0.58
1:B:308:VAL:HG21	1:B:322:LEU:HD21	1.99	0.45
1:A:308:VAL:HG21	1:A:322:LEU:HD21	1.99	0.45
1:B:686:GLU:HG2	1:B:777:ILE:HB	2.00	0.44
1:A:686:GLU:HG2	1:A:777:ILE:HB	2.00	0.42
1:A:416:ASN:O	1:A:417:ALA:HB3	2.20	0.42
1:B:366:LEU:HD11	1:B:382:GLY:HA3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	646/701 (92%)	619 (96%)	25 (4%)	2 (0%)	36	58
1	B	646/701 (92%)	619 (96%)	25 (4%)	2 (0%)	36	58
All	All	1292/1402 (92%)	1238 (96%)	50 (4%)	4 (0%)	36	58

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	416	ASN
1	B	416	ASN
1	A	609	ILE
1	B	609	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	593/630 (94%)	579 (98%)	14 (2%)	43	70
1	B	593/630 (94%)	578 (98%)	15 (2%)	42	69
All	All	1186/1260 (94%)	1157 (98%)	29 (2%)	43	70

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	308	VAL
1	A	328	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	419	GLU
1	A	423	SER
1	A	485	VAL
1	A	609	ILE
1	A	611	MET
1	A	626	LYS
1	A	694	ARG
1	A	715	ASN
1	A	721	VAL
1	A	885	HIS
1	A	898	GLU
1	A	905	VAL
1	B	308	VAL
1	B	328	LYS
1	B	422	GLU
1	B	423	SER
1	B	485	VAL
1	B	609	ILE
1	B	611	MET
1	B	626	LYS
1	B	694	ARG
1	B	715	ASN
1	B	721	VAL
1	B	885	HIS
1	B	898	GLU
1	B	899	ASN
1	B	905	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	448	ASN
1	A	460	ASN
1	A	463	GLN
1	A	491	ASN
1	A	522	ASN
1	A	554	GLN
1	A	742	GLN
1	A	758	ASN
1	A	945	ASN
1	B	448	ASN
1	B	460	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	463	GLN
1	B	491	ASN
1	B	522	ASN
1	B	554	GLN
1	B	742	GLN
1	B	758	ASN
1	B	899	ASN
1	B	945	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	C	9/10 (90%)	0	0
2	D	9/10 (90%)	0	0
2	X	9/10 (90%)	0	0
3	Y	10/11 (90%)	0	0
All	All	37/41 (90%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ADP	B	1001	5	28,29,29	1.50	5 (17%)	43,45,45	1.90	11 (25%)
4	ADP	A	1001	5	28,29,29	1.51	5 (17%)	43,45,45	1.89	11 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	B	1001	5	-	2/16/32/32	0/3/3/3
4	ADP	A	1001	5	-	2/16/32/32	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1001	ADP	C5-C4	4.91	1.47	1.39
4	B	1001	ADP	C5-C4	4.89	1.47	1.39
4	A	1001	ADP	C5-C6	2.95	1.49	1.41
4	B	1001	ADP	C5-C6	2.91	1.49	1.41
4	B	1001	ADP	C8-N7	2.43	1.36	1.31
4	A	1001	ADP	C8-N7	2.37	1.36	1.31
4	B	1001	ADP	C5-N7	-2.29	1.34	1.39
4	A	1001	ADP	C5-N7	-2.28	1.34	1.39
4	A	1001	ADP	PA-O3A	2.23	1.61	1.59
4	B	1001	ADP	PA-O3A	2.13	1.61	1.59

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1001	ADP	C5-C4-N3	-5.71	118.85	126.72
4	B	1001	ADP	C5-C4-N3	-5.67	118.91	126.72
4	A	1001	ADP	N3-C4-N9	4.61	135.00	127.17
4	B	1001	ADP	N3-C4-N9	4.51	134.84	127.17
4	B	1001	ADP	N3-C2-N1	-3.88	122.70	128.58
4	B	1001	ADP	C2-N3-C4	3.88	121.30	111.83
4	A	1001	ADP	C2-N3-C4	3.86	121.27	111.83
4	A	1001	ADP	N3-C2-N1	-3.80	122.82	128.58
4	B	1001	ADP	C4-C5-N7	-3.58	106.49	110.58
4	A	1001	ADP	C4-C5-N7	-3.56	106.51	110.58
4	A	1001	ADP	C4-N9-C8	2.88	108.76	105.74
4	B	1001	ADP	C4-N9-C8	2.79	108.67	105.74

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1001	ADP	C5-N7-C8	2.71	107.70	103.45
4	B	1001	ADP	C5-N7-C8	2.64	107.61	103.45
4	B	1001	ADP	C3'-C2'-C1'	2.59	106.37	101.46
4	A	1001	ADP	C3'-C2'-C1'	2.39	105.98	101.46
4	B	1001	ADP	C6-C5-N7	2.30	136.52	132.09
4	A	1001	ADP	C6-C5-N7	2.24	136.41	132.09
4	B	1001	ADP	C2-N1-C6	2.22	122.38	118.73
4	A	1001	ADP	N9-C8-N7	-2.16	110.88	113.94
4	A	1001	ADP	C2-N1-C6	2.13	122.24	118.73
4	B	1001	ADP	N9-C8-N7	-2.08	110.99	113.94

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	1001	ADP	O4'-C4'-C5'-O5'
4	A	1001	ADP	O4'-C4'-C5'-O5'
4	B	1001	ADP	C3'-C4'-C5'-O5'
4	A	1001	ADP	C3'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	660/701 (94%)	0.91	87 (13%) 7 5	41, 63, 102, 142	0
1	B	660/701 (94%)	0.91	91 (13%) 6 5	41, 63, 103, 128	0
2	C	10/10 (100%)	0.01	0 100 100	41, 51, 70, 75	0
2	D	10/10 (100%)	0.28	0 100 100	42, 46, 71, 84	0
2	X	10/10 (100%)	-0.08	0 100 100	43, 51, 67, 74	0
3	Y	11/11 (100%)	0.68	1 (9%) 15 11	41, 46, 86, 106	0
All	All	1361/1443 (94%)	0.89	179 (13%) 7 5	41, 63, 102, 142	0

All (179) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	Y	0	A	5.5
1	A	417	ALA	4.9
1	B	601	ASP	3.9
1	A	416	ASN	3.9
1	A	531	THR	3.6
1	B	417	ALA	3.6
1	A	899	ASN	3.6
1	A	905	VAL	3.6
1	A	636	ASP	3.5
1	B	536	ASP	3.5
1	B	888	ILE	3.5
1	A	536	ASP	3.5
1	A	972	THR	3.4
1	A	673	ASN	3.4
1	B	667	ARG	3.4
1	B	391	ILE	3.3
1	B	736	GLU	3.3
1	B	668	LYS	3.3
1	B	418	THR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	666	ALA	3.2
1	B	905	VAL	3.2
1	B	673	ASN	3.2
1	B	918	ASN	3.2
1	B	463	GLN	3.2
1	A	918	ASN	3.2
1	A	884	CYS	3.1
1	A	391	ILE	3.1
1	B	563	SER	3.1
1	A	466	GLU	3.1
1	A	930	THR	3.1
1	B	952	LEU	3.0
1	A	885	HIS	3.0
1	A	988	ASP	3.0
1	A	463	GLN	3.0
1	A	459	LYS	3.0
1	B	885	HIS	3.0
1	A	977	LYS	3.0
1	B	652	MET	2.9
1	B	423	SER	2.9
1	A	892	GLU	2.8
1	B	867	ARG	2.8
1	B	464	ALA	2.8
1	B	677	MET	2.8
1	A	660	LYS	2.7
1	A	934	ILE	2.7
1	B	712	ILE	2.7
1	B	564	GLU	2.7
1	B	972	THR	2.7
1	B	471	ILE	2.7
1	A	957	LEU	2.7
1	A	677	MET	2.7
1	A	867	ARG	2.7
1	A	738	LYS	2.7
1	B	935	ILE	2.7
1	B	962	PHE	2.7
1	B	738	LYS	2.7
1	A	470	LEU	2.7
1	A	679	LEU	2.7
1	B	931	ASN	2.7
1	B	636	ASP	2.7
1	B	470	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	884	CYS	2.6
1	A	667	ARG	2.6
1	A	960	ARG	2.6
1	B	456	GLU	2.6
1	B	459	LYS	2.6
1	A	962	PHE	2.6
1	A	729	ILE	2.6
1	A	733	HIS	2.6
1	A	716	PRO	2.6
1	B	716	PRO	2.6
1	B	713	MET	2.6
1	B	620	ASN	2.6
1	B	462	LYS	2.6
1	B	690	THR	2.5
1	B	957	LEU	2.5
1	B	660	LYS	2.5
1	B	664	GLU	2.5
1	B	624	GLU	2.5
1	A	929	GLN	2.5
1	A	652	MET	2.5
1	A	931	ASN	2.5
1	A	465	LYS	2.5
1	B	960	ARG	2.5
1	A	471	ILE	2.5
1	A	864	ARG	2.5
1	B	545	ARG	2.5
1	A	901	HIS	2.5
1	B	531	THR	2.4
1	B	634	ASP	2.4
1	B	901	HIS	2.4
1	B	674	GLU	2.4
1	A	626	LYS	2.4
1	B	973	LYS	2.4
1	B	642	SER	2.4
1	A	415	LEU	2.4
1	A	422	GLU	2.4
1	B	967	GLU	2.4
1	B	879	PHE	2.4
1	B	976	PHE	2.4
1	B	412	ASN	2.3
1	B	460	ASN	2.3
1	B	685	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	668	LYS	2.3
1	B	861	LYS	2.3
1	A	986	PHE	2.3
1	A	614	ALA	2.3
1	A	642	SER	2.3
1	A	712	ILE	2.3
1	A	462	LYS	2.3
1	A	608	THR	2.3
1	B	349	GLN	2.3
1	B	982	LEU	2.3
1	B	766	ALA	2.2
1	B	929	GLN	2.2
1	A	932	VAL	2.2
1	B	759	LEU	2.2
1	A	888	ILE	2.2
1	B	678	LYS	2.2
1	B	863	LYS	2.2
1	A	684	MET	2.2
1	A	419	GLU	2.2
1	B	608	THR	2.2
1	B	715	ASN	2.2
1	A	297	GLY	2.2
1	B	297	GLY	2.2
1	A	713	MET	2.2
1	B	657	ALA	2.2
1	A	418	THR	2.2
1	A	674	GLU	2.2
1	A	538	ARG	2.2
1	A	671	TYR	2.2
1	A	349	GLN	2.2
1	A	634	ASP	2.2
1	B	421	ASP	2.2
1	B	968	ASP	2.2
1	B	963	VAL	2.2
1	A	967	GLU	2.2
1	A	913	TYR	2.2
1	B	465	LYS	2.2
1	A	968	ASP	2.2
1	B	684	MET	2.2
1	A	423	SER	2.2
1	B	934	ILE	2.2
1	B	866	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	714	ASP	2.1
1	B	567	SER	2.1
1	B	986	PHE	2.1
1	B	653	ARG	2.1
1	A	601	ASP	2.1
1	B	413	SER	2.1
1	A	862	ALA	2.1
1	A	539	ARG	2.1
1	A	717	LYS	2.1
1	B	937	LYS	2.1
1	B	899	ASN	2.1
1	A	609	ILE	2.1
1	A	973	LYS	2.1
1	A	741	THR	2.1
1	A	763	THR	2.1
1	B	930	THR	2.1
1	A	661	GLN	2.1
1	B	416	ASN	2.1
1	B	750	ASP	2.0
1	B	666	ALA	2.0
1	B	979	TRP	2.0
1	A	880	LEU	2.0
1	A	458	ILE	2.0
1	A	894	ILE	2.0
1	A	980	GLY	2.0
1	A	540	ASP	2.0
1	A	631	ALA	2.0
1	A	645	ASP	2.0
1	B	392	SER	2.0
1	A	715	ASN	2.0
1	A	935	ILE	2.0
1	A	985	ILE	2.0
1	B	749	ILE	2.0
1	B	894	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ADP	A	1001	27/27	0.91	0.09	64,68,71,71	0
4	ADP	B	1001	27/27	0.91	0.09	58,62,65,66	0
5	MG	B	1002	1/1	0.91	0.14	41,41,41,41	0
5	MG	A	1002	1/1	0.94	0.13	42,42,42,42	0
6	ZN	A	1003	1/1	0.98	0.04	63,63,63,63	0
6	ZN	B	1003	1/1	0.99	0.03	65,65,65,65	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.