



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:58 AM UTC

PDB ID : 5JCH / pdb_00005jch
Title : Crystal structure of chicken MDA5 with 5'p 10-mer dsRNA and ADP-Mg2+
at 2.95 Å resolution (untwinned).
Authors : Cusack, S.; Uchikawa, E.
Deposited on : 2016-04-15
Resolution : 2.95 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1130 (2.98-2.94)
Clashscore	190562	1157 (2.98-2.94)
Ramachandran outliers	187476	1101 (2.98-2.94)
Sidechain outliers	187428	1101 (2.98-2.94)
RSRZ outliers	180081	1130 (2.98-2.94)
RNA backbone	3983	1046 (3.16-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	701	4% 91% 6%
1	B	701	5% 90% 7%
2	U	10	100%
2	V	10	90% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	X	10	 90%10%
2	Y	10	 90%10%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11661 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Melanoma differentiation associated protein-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	661	Total	C	N	O	S	0	0	0
			5396	3403	967	994	32			
1	B	655	Total	C	N	O	S	0	0	0
			5353	3376	959	986	32			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	294	GLY	-	expression tag	UNP D9N195
A	295	ALA	-	expression tag	UNP D9N195
A	296	MET	-	expression tag	UNP D9N195
A	297	GLY	-	expression tag	UNP D9N195
A	436	GLN	GLU	engineered mutation	UNP D9N195
B	294	GLY	-	expression tag	UNP D9N195
B	295	ALA	-	expression tag	UNP D9N195
B	296	MET	-	expression tag	UNP D9N195
B	297	GLY	-	expression tag	UNP D9N195
B	436	GLN	GLU	engineered mutation	UNP D9N195


- Molecule 2 is a RNA chain called RNA (5'-R(P*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	10	Total	C	N	O	P	0	0	0
			214	95	38	71	10			
2	Y	10	Total	C	N	O	P	0	0	0
			213	95	38	70	10			
2	U	10	Total	C	N	O	P	0	0	0
			214	95	38	71	10			
2	V	10	Total	C	N	O	P	0	0	0
			213	95	38	70	10			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
4	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	B	1	Total	Mg	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Melanoma differentiation associated protein-5



- Molecule 1: Melanoma differentiation associated protein-5




- Molecule 2: RNA (5'-R(P*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3')

Chain X: 



- Molecule 2: RNA (5'-R(P*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3')

Chain Y:  90% 10%




- Molecule 2: RNA (5'-R(P*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3')

Chain U:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: RNA (5'-R(P*GP*GP*UP*AP*CP*GP*UP*AP*CP*C)-3')

Chain V:  90% 10%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.08Å 139.73Å 103.19Å 90.00° 110.14° 90.00°	Depositor
Resolution (Å)	50.00 – 2.95 50.00 – 2.95	Depositor EDS
% Data completeness (in resolution range)	94.2 (50.00-2.95) 94.2 (50.00-2.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 2.96Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.256 , 0.270 (Not available) , 0.250	Depositor DCC
R_{free} test set	1907 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	63.5	Xtriage
Anisotropy	0.863	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11661	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.60 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.7162e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/5482	0.78	0/7353
1	B	0.55	0/5438	0.78	0/7293
2	U	0.35	0/238	0.67	0/367
2	V	0.35	0/237	0.65	0/367
2	X	0.35	0/238	0.65	0/367
2	Y	0.36	0/237	0.66	0/367
All	All	0.53	0/11870	0.77	0/16114

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5396	0	5482	7	0
1	B	5353	0	5438	6	0
2	U	214	0	109	0	0
2	V	213	0	109	0	0
2	X	214	0	109	0	0
2	Y	213	0	109	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
4	A	27	0	12	0	0
4	B	27	0	12	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
All	All	11661	0	11380	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:308:VAL:HG11	1:B:320:ILE:HG23	1.87	0.57
1:A:308:VAL:HG11	1:A:320:ILE:HG23	1.87	0.56
1:B:383:LEU:HD11	1:B:396:VAL:HG11	1.93	0.51
1:A:383:LEU:HD11	1:A:396:VAL:HG11	1.93	0.51
1:A:778:VAL:CG1	1:A:797:ALA:HB2	2.45	0.47
1:B:778:VAL:CG1	1:B:797:ALA:HB2	2.45	0.47
1:A:609:ILE:HD11	1:A:704:SER:CB	2.46	0.45
1:B:609:ILE:HD11	1:B:704:SER:CB	2.46	0.45
1:A:436:GLN:HA	1:A:436:GLN:OE1	2.19	0.43
1:B:686:GLU:HG2	1:B:777:ILE:HB	2.02	0.41
1:A:686:GLU:HG2	1:A:777:ILE:HB	2.02	0.41
1:A:434:ILE:HD11	1:A:450:MET:HE3	2.03	0.41
1:B:434:ILE:HD11	1:B:450:MET:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	647/701 (92%)	619 (96%)	28 (4%)	0	100	100
1	B	641/701 (91%)	615 (96%)	26 (4%)	0	100	100
All	All	1288/1402 (92%)	1234 (96%)	54 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	595/630 (94%)	585 (98%)	10 (2%)	53	73
1	B	590/630 (94%)	579 (98%)	11 (2%)	50	71
All	All	1185/1260 (94%)	1164 (98%)	21 (2%)	51	72

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	328	LYS
1	A	529	LYS
1	A	598	LYS
1	A	609	ILE
1	A	694	ARG
1	A	715	ASN
1	A	778	VAL
1	A	852	LEU
1	A	885	HIS
1	A	945	ASN
1	B	328	LYS
1	B	529	LYS
1	B	598	LYS
1	B	609	ILE
1	B	653	ARG
1	B	694	ARG
1	B	715	ASN
1	B	778	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	852	LEU
1	B	885	HIS
1	B	945	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	441	GLN
1	A	497	HIS
1	A	522	ASN
1	A	568	GLN
1	A	617	HIS
1	A	644	GLN
1	B	441	GLN
1	B	497	HIS
1	B	522	ASN
1	B	568	GLN
1	B	617	HIS
1	B	644	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	U	9/10 (90%)	0	0
2	V	9/10 (90%)	1 (11%)	0
2	X	9/10 (90%)	1 (11%)	0
2	Y	9/10 (90%)	1 (11%)	0
All	All	36/40 (90%)	3 (8%)	0

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	X	10	C
2	Y	4	A
2	V	4	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ADP	B	1002	5	28,29,29	1.49	5 (17%)	43,45,45	1.88	10 (23%)
4	ADP	A	1002	5	28,29,29	1.49	5 (17%)	43,45,45	1.88	11 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	B	1002	5	-	2/16/32/32	0/3/3/3
4	ADP	A	1002	5	-	2/16/32/32	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1002	ADP	C5-C4	4.95	1.47	1.39
4	B	1002	ADP	C5-C4	4.90	1.47	1.39
4	B	1002	ADP	C5-C6	2.98	1.49	1.41
4	A	1002	ADP	C5-C6	2.86	1.49	1.41
4	B	1002	ADP	C8-N7	2.43	1.36	1.31
4	A	1002	ADP	C8-N7	2.37	1.36	1.31

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1002	ADP	C5-N7	-2.34	1.34	1.39
4	B	1002	ADP	C5-N7	-2.27	1.34	1.39
4	B	1002	ADP	PA-O3A	2.08	1.61	1.59
4	A	1002	ADP	PA-O3A	2.02	1.61	1.59

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1002	ADP	C5-C4-N3	-5.91	118.58	126.72
4	A	1002	ADP	C5-C4-N3	-5.80	118.73	126.72
4	B	1002	ADP	N3-C4-N9	4.66	135.10	127.17
4	A	1002	ADP	N3-C4-N9	4.66	135.09	127.17
4	B	1002	ADP	C2-N3-C4	3.98	121.56	111.83
4	A	1002	ADP	C2-N3-C4	3.92	121.42	111.83
4	A	1002	ADP	N3-C2-N1	-3.87	122.73	128.58
4	B	1002	ADP	N3-C2-N1	-3.82	122.80	128.58
4	B	1002	ADP	C4-C5-N7	-3.67	106.38	110.58
4	A	1002	ADP	C4-C5-N7	-3.55	106.53	110.58
4	B	1002	ADP	C5-N7-C8	2.84	107.92	103.45
4	A	1002	ADP	C4-N9-C8	2.78	108.66	105.74
4	A	1002	ADP	C5-N7-C8	2.72	107.73	103.45
4	B	1002	ADP	C4-N9-C8	2.68	108.55	105.74
4	B	1002	ADP	C6-C5-N7	2.28	136.49	132.09
4	A	1002	ADP	C3'-C2'-C1'	2.26	105.75	101.46
4	A	1002	ADP	C6-C5-N7	2.24	136.41	132.09
4	B	1002	ADP	N9-C8-N7	-2.20	110.82	113.94
4	A	1002	ADP	N9-C8-N7	-2.13	110.91	113.94
4	A	1002	ADP	C2-N1-C6	2.12	122.21	118.73
4	B	1002	ADP	C2-N1-C6	2.02	122.05	118.73

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	1002	ADP	O4'-C4'-C5'-O5'
4	B	1002	ADP	C3'-C4'-C5'-O5'
4	A	1002	ADP	O4'-C4'-C5'-O5'
4	A	1002	ADP	C3'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	661/701 (94%)	0.59	31 (4%) 36 30	38, 74, 114, 134	0
1	B	655/701 (93%)	0.68	33 (5%) 34 28	39, 82, 132, 155	0
2	U	10/10 (100%)	-0.37	0 100 100	48, 58, 70, 80	0
2	V	10/10 (100%)	-0.32	0 100 100	53, 58, 66, 87	0
2	X	10/10 (100%)	-0.35	0 100 100	44, 54, 67, 69	0
2	Y	10/10 (100%)	-0.21	0 100 100	50, 54, 59, 73	0
All	All	1356/1442 (94%)	0.61	64 (4%) 36 30	38, 78, 125, 155	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	536	ASP	5.5
1	B	536	ASP	4.5
1	B	690	THR	4.4
1	A	967	GLU	3.7
1	A	482	SER	3.7
1	B	685	GLU	3.5
1	A	566	GLY	3.4
1	B	952	LEU	3.3
1	A	563	SER	3.2
1	B	912	LEU	3.1
1	B	563	SER	3.0
1	A	918	ASN	2.9
1	B	464	ALA	2.9
1	B	565	PHE	2.9
1	A	565	PHE	2.8
1	B	967	GLU	2.7
1	A	782	GLY	2.7
1	B	972	THR	2.7
1	B	948	VAL	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	673	ASN	2.6
1	A	960	ARG	2.6
1	B	655	PHE	2.6
1	A	867	ARG	2.5
1	B	625	LEU	2.5
1	A	601	ASP	2.5
1	B	601	ASP	2.5
1	B	957	LEU	2.5
1	A	876	LEU	2.5
1	A	691	GLU	2.4
1	B	977	LYS	2.4
1	A	564	GLU	2.4
1	A	608	THR	2.4
1	A	736	GLU	2.3
1	A	441	GLN	2.3
1	B	879	PHE	2.3
1	A	885	HIS	2.3
1	A	935	ILE	2.3
1	A	480	THR	2.3
1	B	631	ALA	2.3
1	B	766	ALA	2.3
1	A	673	ASN	2.2
1	A	977	LYS	2.2
1	A	613	ASP	2.2
1	B	482	SER	2.2
1	B	986	PHE	2.2
1	A	602	ALA	2.2
1	A	631	ALA	2.2
1	B	929	GLN	2.2
1	A	707	ALA	2.2
1	A	985	ILE	2.1
1	A	973	LYS	2.1
1	B	973	LYS	2.1
1	B	894	ILE	2.1
1	B	463	GLN	2.1
1	B	884	CYS	2.1
1	A	733	HIS	2.1
1	A	412	ASN	2.1
1	B	618	LEU	2.1
1	B	965	ALA	2.0
1	B	968	ASP	2.0
1	B	530	LYS	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	885	HIS	2.0
1	A	464	ALA	2.0
1	B	947	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ADP	B	1002	27/27	0.80	0.12	101,104,110,111	0
4	ADP	A	1002	27/27	0.87	0.09	81,87,100,101	0
5	MG	A	1003	1/1	0.95	0.05	53,53,53,53	0
5	MG	B	1003	1/1	0.97	0.05	46,46,46,46	0
3	ZN	A	1001	1/1	0.99	0.03	74,74,74,74	0
3	ZN	B	1001	1/1	0.99	0.02	100,100,100,100	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ADP B 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ADP A 1002:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.