



Full wwPDB EM Validation Report ⓘ

Mar 8, 2026 – 08:08 AM UTC

PDB ID : 6JCR / pdb_00006jcr
EMDB ID : EMD-9795
Title : AAV1 in neutral condition at 3.07 Ang
Authors : Lou, Z.; Zhang, R.
Deposited on : 2019-01-30
Resolution : 3.07 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13977 (2.57 - 3.57)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	518	10% 86% 14%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 4112 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	518	4112	2601	708	787	16	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Capsid protein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	2642	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TECNAI ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	1.6	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON II (4k x 4k)	Depositor
Maximum map value	0.214	Depositor
Minimum map value	-0.113	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.017	Depositor
Recommended contour level	0.05	Depositor
Map size ($\text{\AA}$)	353.4, 353.4, 353.4	wwPDB
Map dimensions	380, 380, 380	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	0.93, 0.93, 0.93	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/4238	0.70	5/5778 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	500	ASN	CA-C-N	9.71	139.17	121.70
1	A	500	ASN	C-N-CA	9.71	139.17	121.70
1	A	465	ARG	CA-C-N	-7.56	114.28	121.46
1	A	465	ARG	C-N-CA	-7.56	114.28	121.46
1	A	466	GLY	N-CA-C	6.55	120.92	110.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	665	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4112	0	3876	64	0
All	All	4112	0	3876	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:571:PRO:HG3	1:A:577:PHE:CZ	1.37	1.53
1:A:571:PRO:CG	1:A:577:PHE:CZ	1.90	1.53
1:A:571:PRO:HG2	1:A:577:PHE:CE2	1.46	1.46
1:A:571:PRO:HG3	1:A:577:PHE:CE1	1.62	1.35
1:A:481:GLY:O	1:A:577:PHE:CE2	1.85	1.28
1:A:571:PRO:CG	1:A:577:PHE:CE2	2.17	1.15
1:A:483:CYS:HB2	1:A:577:PHE:CD2	2.04	0.92
1:A:481:GLY:O	1:A:577:PHE:CZ	2.27	0.88
1:A:481:GLY:O	1:A:577:PHE:HE2	1.57	0.84
1:A:571:PRO:CD	1:A:577:PHE:CZ	2.61	0.84
1:A:571:PRO:HG2	1:A:577:PHE:CD2	2.13	0.83
1:A:483:CYS:HB2	1:A:577:PHE:HD2	1.46	0.80
1:A:286:ASN:HD21	1:A:619:TRP:H	1.34	0.73
1:A:480:PRO:HB2	1:A:577:PHE:CE1	2.31	0.66
1:A:480:PRO:HG2	1:A:577:PHE:CE1	2.31	0.66
1:A:483:CYS:CB	1:A:577:PHE:HD2	2.12	0.63
1:A:316:LEU:HB3	1:A:410:PHE:HB3	1.83	0.60
1:A:448:ARG:HB2	1:A:462:LEU:HB2	1.84	0.60
1:A:404:LEU:HD11	1:A:410:PHE:HB2	1.86	0.56
1:A:528:LYS:NZ	1:A:531:GLU:OE1	2.38	0.56
1:A:480:PRO:CG	1:A:577:PHE:HE1	2.19	0.56
1:A:571:PRO:HD3	1:A:577:PHE:CZ	2.40	0.56
1:A:559:MET:HB3	1:A:726:PRO:HD3	1.88	0.55
1:A:361:GLY:HA3	1:A:374:PRO:HG3	1.89	0.55
1:A:628:HIS:HD2	1:A:631:PRO:HG3	1.71	0.55
1:A:428:SER:HB3	1:A:733:THR:HB	1.87	0.55
1:A:480:PRO:CG	1:A:577:PHE:CE1	2.90	0.54
1:A:250:PRO:HG3	1:A:372:MET:HE3	1.90	0.54
1:A:298:GLN:NE2	1:A:701:TYR:O	2.41	0.54
1:A:273:TYR:HA	1:A:382:ASN:HD22	1.72	0.54
1:A:252:TYR:OH	1:A:373:ILE:O	2.23	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:480:PRO:HB2	1:A:577:PHE:CZ	2.43	0.53
1:A:521:GLY:HA2	1:A:634:LEU:HD13	1.89	0.53
1:A:571:PRO:HD3	1:A:577:PHE:HZ	1.73	0.53
1:A:571:PRO:CG	1:A:577:PHE:CE1	2.51	0.51
1:A:440:ASP:HB3	1:A:468:PRO:HD3	1.94	0.50
1:A:480:PRO:HB2	1:A:577:PHE:HE1	1.75	0.50
1:A:414:TYR:OH	1:A:642:ASN:O	2.29	0.49
1:A:342:GLN:HG2	1:A:403:MET:HG2	1.96	0.48
1:A:436:ASN:HD22	1:A:437:PRO:HD2	1.78	0.48
1:A:572:VAL:HG12	1:A:574:THR:H	1.78	0.48
1:A:509:TYR:HD1	1:A:518:ILE:HD13	1.79	0.47
1:A:302:ASN:HD21	1:A:701:TYR:H	1.62	0.47
1:A:243:SER:OG	1:A:245:ARG:NH1	2.48	0.47
1:A:312:LEU:HG	1:A:414:TYR:HB3	1.97	0.46
1:A:480:PRO:CB	1:A:577:PHE:HE1	2.29	0.45
1:A:280:TRP:CE2	1:A:650:LYS:HD2	2.52	0.45
1:A:650:LYS:NZ	1:A:651:ASN:O	2.50	0.45
1:A:315:LYS:HG3	1:A:411:THR:HG22	1.98	0.44
1:A:354:VAL:H	1:A:646:GLN:HE22	1.65	0.44
1:A:451:ASN:HD22	1:A:458:ASN:HB3	1.83	0.44
1:A:487:GLN:HB3	1:A:537:MET:HE3	1.99	0.43
1:A:480:PRO:CB	1:A:577:PHE:CE1	2.98	0.43
1:A:585:GLN:HB3	1:A:591:PRO:HB3	2.01	0.43
1:A:453:SER:OG	1:A:458:ASN:ND2	2.51	0.43
1:A:249:LEU:HB2	1:A:373:ILE:HD12	2.01	0.42
1:A:440:ASP:OD1	1:A:440:ASP:N	2.53	0.41
1:A:451:ASN:ND2	1:A:460:ASP:OD2	2.39	0.41
1:A:655:PRO:HA	1:A:670:PHE:HA	2.02	0.41
1:A:323:GLU:HB3	1:A:336:ASN:ND2	2.36	0.41
1:A:289:HIS:ND1	1:A:615:GLN:HA	2.36	0.41
1:A:382:ASN:HD21	1:A:514:ARG:HH12	1.68	0.41
1:A:628:HIS:CD2	1:A:631:PRO:HG3	2.53	0.40
1:A:382:ASN:OD1	1:A:382:ASN:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	516/518 (100%)	483 (94%)	33 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	451/451 (100%)	449 (100%)	2 (0%)	84	84

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	528	LYS
1	A	577	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	ASN
1	A	229	HIS
1	A	286	ASN
1	A	302	ASN
1	A	304	ASN
1	A	327	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	336	ASN
1	A	342	GLN
1	A	408	ASN
1	A	422	HIS
1	A	436	ASN
1	A	458	ASN
1	A	474	GLN
1	A	497	ASN
1	A	557	ASN
1	A	615	GLN
1	A	628	HIS
1	A	646	GLN
1	A	651	ASN
1	A	704	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-9795. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 190



Y Index: 190




Z Index: 190


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 174



Y Index: 206




Z Index: 206


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views

This section was not generated.


6.6 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 11111 nm³; this corresponds to an approximate mass of 1004 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.326 Å⁻¹

8 Fourier-Shell correlation ⓘ

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-9795 and PDB model 6JCR. Per-residue inclusion information can be found in section 3 on page 4.

9.1 Map-model overlay [i](#)

This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

9.4 Atom inclusion [i](#)



At the recommended contour level, 77% of all backbone atoms, 68% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.05) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6780	0.4520
A	0.6780	0.4520

