



Full wwPDB EM Validation Report ⓘ

Mar 7, 2026 – 01:28 AM UTC

PDB ID : 3JD6 / pdb_00003jd6
EMDB ID : EMD-6425
Title : Double octamer structure of retinoschisin, a cell-cell adhesion protein of the retina
Authors : Tolun, G.; Vijayasarathy, C.; Huang, R.; Zeng, Y.; Li, Y.; Steven, A.C.; Sieving, P.A.; Heymann, J.B.
Deposited on : 2016-04-12
Resolution : 4.10 Å (reported)
Based on initial model : 1SDD

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	6458 (3.60 - 4.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	O	207	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2693 atoms, of which 1326 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinoschisin.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
1	O	167	2693	869	1326	237	252	9	0	0


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	225	HIS	-	expression tag	UNP O15537
O	226	HIS	-	expression tag	UNP O15537
O	227	HIS	-	expression tag	UNP O15537
O	228	HIS	-	expression tag	UNP O15537
O	229	HIS	-	expression tag	UNP O15537
O	230	HIS	-	expression tag	UNP O15537

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Retinoschisin



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, D8	Depositor
Number of particles used	9096	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	Included in iterative refinement	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	30	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	39000	Depositor
Image detector	GATAN K2 (4k x 4k)	Depositor
Maximum map value	6.216	Depositor
Minimum map value	-3.080	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	3	Depositor
Map size ($\text{\AA}$)	164.79999, 164.79999, 164.79999	wwPDB
Map dimensions	160, 160, 160	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.03, 1.03, 1.03	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	O	1.94	52/1403 (3.7%)	1.94	64/1905 (3.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	O	0	3

All (52) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	O	201	LEU	N-CA	-9.19	1.34	1.46
1	O	115	LYS	CA-C	-8.72	1.42	1.52
1	O	60	ILE	CA-C	-8.38	1.43	1.52
1	O	82	THR	CA-C	-7.55	1.43	1.52
1	O	178	GLY	CA-C	-7.44	1.43	1.51
1	O	215	GLU	CA-C	-7.03	1.43	1.52
1	O	83	CYS	N-CA	-7.01	1.37	1.45
1	O	201	LEU	C-N	6.98	1.39	1.33
1	O	99	ASN	N-CA	-6.82	1.37	1.46
1	O	112	TRP	CA-C	-6.78	1.44	1.52
1	O	69	LEU	CA-C	-6.53	1.44	1.52
1	O	180	SER	N-CA	-6.26	1.39	1.46
1	O	179	ASN	CA-C	-6.20	1.45	1.52
1	O	147	TRP	C-N	6.13	1.41	1.33
1	O	124	GLN	CA-C	-6.12	1.45	1.52
1	O	74	GLY	CA-C	-6.10	1.43	1.51
1	O	74	GLY	C-N	6.09	1.42	1.33
1	O	185	THR	CA-C	6.01	1.59	1.52
1	O	213	ARG	CA-C	-6.00	1.45	1.52
1	O	207	HIS	CA-C	-5.99	1.45	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	O	172	ASN	CA-C	-5.99	1.45	1.53
1	O	87	GLU	C-N	5.93	1.42	1.33
1	O	112	TRP	CD2-CE3	-5.92	1.30	1.40
1	O	150	LYS	N-CA	-5.86	1.38	1.45
1	O	100	LYS	CA-C	-5.79	1.44	1.52
1	O	188	ASN	C-O	-5.77	1.17	1.23
1	O	75	GLU	CA-C	-5.77	1.45	1.52
1	O	152	SER	CA-C	-5.66	1.45	1.52
1	O	204	LEU	C-N	5.56	1.41	1.33
1	O	148	MET	N-CA	-5.54	1.39	1.46
1	O	152	SER	N-CA	-5.52	1.39	1.46
1	O	106	GLN	N-CA	-5.50	1.39	1.46
1	O	191	ARG	CA-C	-5.49	1.46	1.52
1	O	110	CYS	CA-C	-5.48	1.46	1.52
1	O	218	GLU	CA-C	-5.46	1.47	1.53
1	O	181	ASP	CA-C	-5.45	1.46	1.52
1	O	76	VAL	N-CA	-5.42	1.39	1.46
1	O	184	SER	N-CA	5.40	1.52	1.46
1	O	139	GLN	N-CA	-5.40	1.39	1.46
1	O	102	ARG	C-N	5.33	1.41	1.33
1	O	89	TYR	N-CA	-5.29	1.40	1.46
1	O	72	GLU	C-N	5.28	1.41	1.34
1	O	134	SER	CA-C	5.27	1.59	1.52
1	O	96	TRP	CA-C	-5.24	1.46	1.52
1	O	152	SER	C-N	5.23	1.40	1.33
1	O	183	THR	CA-C	-5.23	1.47	1.53
1	O	70	GLY	C-O	-5.22	1.16	1.23
1	O	164	ILE	N-CA	-5.17	1.40	1.46
1	O	150	LYS	C-N	5.15	1.40	1.33
1	O	145	ASP	C-N	5.15	1.40	1.33
1	O	197	ARG	C-N	5.09	1.40	1.33
1	O	124	GLN	N-CA	5.01	1.52	1.45

All (64) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	O	60	ILE	CB-CA-C	9.94	120.62	110.88
1	O	220	VAL	N-CA-C	-9.90	93.86	108.12
1	O	72	GLU	N-CA-C	-9.65	99.63	111.40
1	O	78	PRO	CA-C-N	8.69	134.50	120.60
1	O	78	PRO	C-N-CA	8.69	134.50	120.60
1	O	63	CYS	O-C-N	-7.94	112.19	121.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	O	71	PHE	N-CA-C	-7.75	103.66	113.43
1	O	182	ARG	CA-C-N	7.45	133.36	122.63
1	O	182	ARG	C-N-CA	7.45	133.36	122.63
1	O	221	SER	N-CA-C	7.18	121.68	112.34
1	O	176	PHE	CA-C-N	6.92	131.32	121.42
1	O	176	PHE	C-N-CA	6.92	131.32	121.42
1	O	171	GLY	N-CA-C	-6.87	106.22	113.58
1	O	106	GLN	CA-C-O	6.80	128.01	120.94
1	O	178	GLY	O-C-N	6.73	129.95	122.82
1	O	77	THR	CA-C-O	-6.61	113.48	120.88
1	O	85	ASN	CA-CB-CG	-6.55	106.05	112.60
1	O	121	GLN	N-CA-C	-6.51	99.14	109.23
1	O	120	SER	N-CA-C	-6.28	103.30	111.71
1	O	93	TYR	N-CA-C	-6.18	105.68	113.72
1	O	97	THR	CA-C-N	6.17	129.16	120.28
1	O	97	THR	C-N-CA	6.17	129.16	120.28
1	O	85	ASN	CA-C-N	6.09	125.72	119.56
1	O	85	ASN	C-N-CA	6.09	125.72	119.56
1	O	189	LEU	N-CA-C	-6.06	99.78	109.96
1	O	172	ASN	N-CA-C	6.01	119.72	111.54
1	O	212	ILE	O-C-N	5.95	129.34	122.97
1	O	90	VAL	N-CA-C	-5.89	107.17	112.12
1	O	217	LEU	N-CA-C	-5.83	99.90	109.40
1	O	147	TRP	CA-C-N	5.81	129.09	120.90
1	O	147	TRP	C-N-CA	5.81	129.09	120.90
1	O	175	VAL	N-CA-C	-5.80	99.92	108.85
1	O	159	GLU	N-CA-C	-5.79	105.52	112.58
1	O	112	TRP	CA-CB-CG	5.72	124.47	113.60
1	O	203	PRO	O-C-N	5.72	128.48	122.93
1	O	65	TYR	CA-C-N	5.70	130.85	121.39
1	O	65	TYR	C-N-CA	5.70	130.85	121.39
1	O	110	CYS	N-CA-C	-5.68	100.14	109.40
1	O	205	GLY	O-C-N	-5.68	116.08	122.78
1	O	89	TYR	CA-C-O	5.66	124.91	119.08
1	O	112	TRP	CD2-CE2-CZ2	-5.57	116.83	122.40
1	O	169	GLN	CA-C-N	5.55	128.65	120.82
1	O	169	GLN	C-N-CA	5.55	128.65	120.82
1	O	210	ILE	O-C-N	-5.47	116.46	122.69
1	O	205	GLY	N-CA-C	-5.46	105.80	112.68
1	O	177	TYR	O-C-N	-5.41	116.63	123.01
1	O	160	ARG	CA-C-O	5.36	127.46	121.40
1	O	171	GLY	CA-C-N	5.32	129.67	122.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	O	171	GLY	C-N-CA	5.32	129.67	122.07
1	O	95	SER	N-CA-C	-5.29	99.54	110.80
1	O	81	ILE	CA-C-O	5.26	126.81	120.66
1	O	111	ALA	N-CA-CB	5.25	119.37	110.49
1	O	87	GLU	CA-C-N	5.25	130.00	122.40
1	O	87	GLU	C-N-CA	5.25	130.00	122.40
1	O	120	SER	O-C-N	5.22	128.68	122.21
1	O	201	LEU	N-CA-C	-5.18	100.08	108.52
1	O	83	CYS	CA-C-N	5.16	130.37	120.97
1	O	83	CYS	C-N-CA	5.16	130.37	120.97
1	O	118	ASP	CA-C-N	5.13	129.42	122.19
1	O	118	ASP	C-N-CA	5.13	129.42	122.19
1	O	70	GLY	O-C-N	5.12	129.36	122.70
1	O	68	PRO	CA-C-O	-5.08	115.55	121.34
1	O	81	ILE	N-CA-C	-5.05	99.90	107.37
1	O	213	ARG	O-C-N	-5.02	116.70	123.23

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	O	221	SER	Peptide
1	O	62	GLU	Peptide
1	O	95	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	O	1367	1326	1327	7	0
All	All	1367	1326	1327	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:222:LYS:O	1:O:223:CYS:HB2	2.12	0.49
1:O:207:HIS:O	1:O:208:VAL:HB	2.16	0.46
1:O:91:GLY:O	1:O:92:TRP:HB2	2.17	0.44
1:O:108:PHE:CG	1:O:108:PHE:O	2.71	0.44
1:O:222:LYS:O	1:O:223:CYS:CB	2.68	0.42
1:O:96:TRP:HA	1:O:96:TRP:CE3	2.55	0.42
1:O:71:PHE:CD1	1:O:215:GLU:HA	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	O	165/207 (80%)	142 (86%)	22 (13%)	1 (1%)	21 58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	O	61	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	O	152/184 (83%)	152 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-6425. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 80



Y Index: 80




Z Index: 80


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 86



Y Index: 74




Z Index: 57


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 146 nm³; this corresponds to an approximate mass of 131 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.244 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.244 Å⁻¹

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.10	-	-
Author-provided FSC curve	4.24	4.88	4.36
Unmasked-calculated*	-	-	-


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-6425 and PDB model 3JD6. Per-residue inclusion information can be found in section 3 on page 4.

9.1 Map-model overlays


9.1.1 Map-model overlay [i](#)



X




Y




Z


9.1.2 Map-model assembly overlay [i](#)



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 3.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3).

9.4 Atom inclusion [i](#)



At the recommended contour level, 49% of all backbone atoms, 50% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.4980	0.2300
O	0.5190	0.2300

