



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:53 PM UTC

PDB ID : 2JEA / pdb_00002jea
Title : Structure of a 9-subunit archaeal exosome bound to RNA
Authors : Lorentzen, E.; Conti, E.
Deposited on : 2007-01-16
Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)
RNA backbone	3983	1004 (2.64-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	18% 85% 12% • •
2	B	250	13% 77% 16% • 6%
3	C	35	20% 11% • 6% 80%
4	I	251	26% 68% 12% • 18%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5627 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EXOSOME COMPLEX EXONUCLEASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	S	0	9	0
			2091	1340	338	406	7			

- Molecule 2 is a protein called EXOSOME COMPLEX EXONUCLEASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	234	Total	C	N	O	S	0	7	0
			1809	1152	309	337	11			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	182	ALA	ASP	engineered mutation	UNP Q9UXC2

- Molecule 3 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	7	Total	C	N	O	P	0	0	2
			112	50	25	32	5			

- Molecule 4 is a protein called EXOSOME COMPLEX RNA-BINDING PROTEIN 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	I	207	Total	C	N	O	S	0	0	0
			1477	963	238	274	2			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	8	GLU	LYS	conflict	UNP Q9UXC4

- Molecule 5 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: C₁₀H₂₂O₆).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	C	O	0
			16	10	6	


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	62	Total	O	0	0
			62	62		
6	B	51	Total	O	0	0
			51	51		
6	C	3	Total	O	0	0
			3	3		
6	I	6	Total	O	0	0
			6	6		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: EXOSOME COMPLEX EXONUCLEASE 2



• Molecule 2: EXOSOME COMPLEX EXONUCLEASE 1




• Molecule 3: RNA



• Molecule 4: EXOSOME COMPLEX RNA-BINDING PROTEIN 1





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	135.85Å 135.85Å 135.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.95 – 2.33 42.95 – 2.33	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.95-2.33) 99.7 (42.95-2.33)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.46 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.205 , 0.243 0.270 , 0.285	Depositor DCC
R_{free} test set	1788 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	43.4	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.038 for l,-k,h	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5627	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/2147	0.89	1/2922 (0.0%)
2	B	0.76	0/1858	0.88	0/2514
3	C	0.94	1/125 (0.8%)	1.50	3/193 (1.6%)
4	I	0.68	0/1505	0.85	1/2064 (0.0%)
All	All	0.75	1/5635 (0.0%)	0.90	5/7693 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	400	A	O3'-P	-8.42	1.48	1.61

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	404	A	C2'-C3'-O3'	-7.48	102.47	113.70
3	C	403	A	P-O3'-C3'	-6.28	110.78	120.20
1	A	7	ASN	N-CA-C	5.69	119.78	112.26
3	C	404	A	N9-C1'-C2'	5.52	120.28	112.00
4	I	126	VAL	CB-CA-C	-5.36	102.50	110.33

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	6	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2091	0	2128	22	0
2	B	1809	0	1859	32	0
3	C	112	0	57	0	0
4	I	1477	0	1373	18	0
5	A	16	0	22	0	0
6	A	62	0	0	0	0
6	B	51	0	0	5	0
6	C	3	0	0	0	0
6	I	6	0	0	0	0
All	All	5627	0	5439	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:I:167:VAL:HG22	4:I:214:ILE:HD11	1.49	0.92
1:A:53:SER:H	1:A:167:ASN:HD22	1.27	0.82
2:B:81:TYR:HA	2:B:129:THR:HG22	1.65	0.78
4:I:208:ILE:HA	4:I:211:ILE:HG22	1.67	0.74
4:I:76:VAL:HG22	4:I:191:ASN:CG	2.13	0.73
1:A:14:ILE:HD12	1:A:14:ILE:H	1.54	0.71
2:B:206:ASN:ND2	6:B:2039:HOH:O	2.27	0.68
2:B:129:THR:HG21	2:B:142:SER:OG	1.94	0.68
2:B:168:LYS:NZ	6:B:2028:HOH:O	2.27	0.68
1:A:8:GLN:O	1:A:10:ILE:N	2.30	0.64
2:B:235:ALA:O	2:B:238:SER:OG	2.15	0.63
2:B:118:GLU:HG3	6:B:2017:HOH:O	1.99	0.61
1:A:91:LEU:HD12	1:A:147:LEU:HD23	1.85	0.59
2:B:49:MET:HE1	2:B:144:MET:HE1	1.85	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:198:ASN:O	2:B:198:ASN:ND2	2.36	0.59
1:A:116:ARG:NH2	2:B:102:GLU:OE1	2.35	0.58
4:I:13:PRO:HD2	4:I:15:SER:OG	2.04	0.58
1:A:246[B]:MET:HE2	1:A:250:ASP:HB3	1.86	0.56
2:B:49:MET:HE3	6:B:2022:HOH:O	2.04	0.56
4:I:206:ILE:O	4:I:206:ILE:HG23	2.04	0.56
2:B:198:ASN:O	2:B:198:ASN:CG	2.49	0.55
1:A:11:ILE:HG21	1:A:219[A]:SER:OG	2.07	0.53
1:A:91:LEU:HD11	1:A:145:TYR:HD2	1.74	0.53
1:A:116:ARG:HH22	2:B:102:GLU:CD	2.17	0.52
4:I:16:ILE:HD13	4:I:50:LEU:HD12	1.92	0.51
2:B:164[A]:VAL:CG1	2:B:228:ILE:CD1	2.88	0.51
2:B:135:ASP:O	2:B:136:ALA:HB3	2.09	0.51
1:A:116:ARG:HD2	6:B:2037:HOH:O	2.11	0.50
1:A:1[A]:MET:HE2	2:B:78:ARG:HB3	1.93	0.50
4:I:167:VAL:CG2	4:I:214:ILE:HD11	2.33	0.50
2:B:37:VAL:HG11	2:B:46:ILE:HD11	1.93	0.49
1:A:28:ARG:CZ	1:A:210[B]:VAL:HG22	2.43	0.49
1:A:20:VAL:HG13	1:A:23:PHE:CD2	2.49	0.48
2:B:64:MET:HE1	2:B:76:VAL:CG2	2.44	0.47
4:I:194:ILE:HG13	4:I:194:ILE:O	2.13	0.47
4:I:16:ILE:HD13	4:I:50:LEU:CD1	2.45	0.47
4:I:62:PRO:HG2	4:I:162:VAL:HG22	1.96	0.47
1:A:28:ARG:CZ	1:A:210[A]:VAL:HG13	2.44	0.47
1:A:28:ARG:HD2	1:A:34:LEU:HA	1.97	0.47
4:I:189:ALA:C	4:I:191:ASN:H	2.21	0.47
2:B:108:ARG:O	2:B:112:GLU:HG2	2.15	0.46
2:B:48:GLU:HG2	2:B:53:LYS:HG3	1.98	0.46
4:I:36:ILE:HD11	4:I:59:GLU:HA	1.97	0.46
4:I:96:TYR:HB3	4:I:138:ASP:OD1	2.17	0.45
2:B:64:MET:CE	2:B:70:SER:HA	2.46	0.45
2:B:115:VAL:HA	2:B:157:MET:HG2	2.00	0.44
2:B:135:ASP:O	2:B:136:ALA:CB	2.65	0.44
4:I:62:PRO:O	4:I:162:VAL:HG13	2.18	0.44
1:A:163:ALA:HA	1:A:273:LEU:HD21	2.00	0.44
4:I:191:ASN:HD22	4:I:191:ASN:HA	1.57	0.44
2:B:65:HIS:N	2:B:66:PRO:CD	2.81	0.44
2:B:131:ILE:HD12	2:B:138:SER:HB2	2.00	0.43
1:A:230:THR:HB	1:A:231:PRO:CD	2.49	0.43
2:B:164[A]:VAL:HG13	2:B:228:ILE:CD1	2.49	0.42
2:B:190:PRO:HG2	2:B:204:GLN:HB2	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:I:77:ILE:HD11	4:I:193:ARG:HB3	2.00	0.42
1:A:7:ASN:O	1:A:9:ASN:N	2.54	0.41
1:A:88:VAL:HG23	1:A:111:ALA:HA	2.02	0.41
2:B:64:MET:HE1	2:B:70:SER:HA	2.02	0.41
2:B:144:MET:HG2	2:B:161:ILE:O	2.21	0.41
2:B:164[A]:VAL:HG22	2:B:225:ILE:HG13	2.03	0.41
4:I:92:ILE:O	4:I:191:ASN:ND2	2.51	0.41
1:A:203:ALA:HB3	1:A:210[B]:VAL:HG12	2.03	0.41
4:I:111:ASN:H	4:I:114:GLU:HG3	1.86	0.40
1:A:11:ILE:HD11	1:A:16:LYS:HG3	2.03	0.40
1:A:233:LEU:HG	1:A:259:ARG:HG3	2.03	0.40
2:B:76:VAL:O	2:B:124:ALA:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/277 (101%)	269 (96%)	8 (3%)	2 (1%)	18	18
2	B	239/250 (96%)	232 (97%)	6 (2%)	1 (0%)	30	32
4	I	201/251 (80%)	183 (91%)	17 (8%)	1 (0%)	24	26
All	All	719/778 (92%)	684 (95%)	31 (4%)	4 (1%)	21	22

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	GLN
1	A	9	ASN
4	I	13	PRO
2	B	136	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/243 (95%)	219 (95%)	12 (5%)	21	25
2	B	193/208 (93%)	187 (97%)	6 (3%)	35	45
4	I	140/223 (63%)	121 (86%)	19 (14%)	3	3
All	All	564/674 (84%)	527 (93%)	37 (7%)	15	17

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	HIS
1	A	6	SER
1	A	11	ILE
1	A	15	LYS
1	A	22	LEU
1	A	32	ARG
1	A	72	ASP
1	A	210[A]	VAL
1	A	210[B]	VAL
1	A	224	LYS
1	A	233	LEU
1	A	236	VAL
2	B	11	LYS
2	B	119	LEU
2	B	129	THR
2	B	177	LEU
2	B	198	ASN
2	B	236	LEU
4	I	17	VAL
4	I	18	VAL
4	I	22	LEU
4	I	66	SER
4	I	76	VAL
4	I	77	ILE
4	I	82	ASP
4	I	83	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	I	97	LYS
4	I	106	LEU
4	I	126	VAL
4	I	152	SER
4	I	162	VAL
4	I	179	THR
4	I	191	ASN
4	I	194	ILE
4	I	207	LEU
4	I	214	ILE
4	I	220	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	83	ASN
1	A	167	ASN
2	B	198	ASN
2	B	199	GLN
4	I	191	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	C	3/35 (8%)	2 (66%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	C	403	A
3	C	404	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	1PE	A	1276	-	15,15,15	0.56	0	14,14,14	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	1PE	A	1276	-	-	5/13/13/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1276	1PE	OH4-C13-C23-OH3
5	A	1276	1PE	OH6-C15-C25-OH5
5	A	1276	1PE	C16-C26-OH6-C15
5	A	1276	1PE	C14-C24-OH4-C13
5	A	1276	1PE	C12-C22-OH3-C23

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/277 (98%)	1.26	50 (18%) 3 4	28, 54, 68, 73	10 (3%)
2	B	234/250 (93%)	1.17	33 (14%) 6 7	31, 54, 63, 72	7 (2%)
3	C	7/35 (20%)	11.96	7 (100%) 0 0	10, 12, 13, 14	7 (100%)
4	I	207/251 (82%)	1.66	65 (31%) 1 1	46, 53, 60, 62	0
All	All	722/813 (88%)	1.45	155 (21%) 2 2	10, 53, 65, 73	24 (3%)

All (155) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	395	A	36.1
3	C	403	A	10.1
3	C	404	A	9.6
3	C	396	A	7.6
3	C	401	A	7.3
3	C	400	A	6.5
3	C	402	A	6.5
2	B	198	ASN	4.5
1	A	8	GLN	4.4
1	A	9	ASN	4.2
1	A	94	ALA	4.2
4	I	110	ILE	4.1
4	I	13	PRO	3.9
4	I	186	ILE	3.9
1	A	220	ILE	3.8
1	A	75	TYR	3.8
2	B	8	GLU	3.7
4	I	206	ILE	3.7
4	I	196	ALA	3.7
1	A	-1	GLY	3.6
2	B	99	ARG	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	I	116	LEU	3.5
1	A	10	ILE	3.5
2	B	219	ASP	3.5
1	A	93	LEU	3.3
1	A	19	ILE	3.3
4	I	220	ILE	3.3
1	A	100	PRO	3.3
1	A	103	PRO	3.3
4	I	155	ILE	3.3
4	I	182	SER	3.3
2	B	30	SER	3.2
2	B	236	LEU	3.2
1	A	24	GLU	3.1
1	A	7	ASN	3.1
2	B	240	TYR	3.0
4	I	121	ASP	3.0
1	A	76	GLU	3.0
4	I	56	THR	3.0
2	B	9	ARG	2.9
4	I	10	VAL	2.9
4	I	90	VAL	2.9
4	I	41	SER	2.9
4	I	168	ILE	2.9
4	I	8	GLU	2.9
4	I	171	ASN	2.9
4	I	221	LYS	2.8
1	A	208	TYR	2.8
2	B	103	LEU	2.8
2	B	200	VAL	2.8
1	A	80	ASN	2.7
1	A	209	LEU	2.7
4	I	58	PHE	2.7
4	I	143	VAL	2.7
4	I	195	TRP	2.7
4	I	147	ASP	2.7
4	I	88	TRP	2.7
1	A	20	VAL	2.6
4	I	202	PHE	2.6
1	A	114	VAL	2.6
4	I	181	LYS	2.6
1	A	102	PRO	2.6
4	I	113	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	21	SER	2.6
4	I	204	GLU	2.6
4	I	24	ALA	2.6
1	A	23	PHE	2.6
1	A	113	VAL	2.5
4	I	66	SER	2.5
1	A	159	LEU	2.5
2	B	183	MET	2.5
2	B	216	GLN	2.5
1	A	14	ILE	2.5
4	I	9	ILE	2.5
4	I	75	ILE	2.5
1	A	22	LEU	2.5
4	I	43	TYR	2.4
4	I	106	LEU	2.4
2	B	110	ALA	2.4
1	A	78	THR	2.4
2	B	241	VAL	2.4
2	B	14	LEU	2.4
2	B	50	GLY	2.4
2	B	207	GLY	2.4
1	A	197	VAL	2.4
4	I	47	VAL	2.4
1	A	149	TYR	2.4
4	I	205	GLU	2.4
4	I	26	GLY	2.4
1	A	12	PRO	2.3
1	A	97	THR	2.3
4	I	179	THR	2.3
1	A	0	HIS	2.3
2	B	65	HIS	2.3
2	B	69	LEU	2.3
4	I	55	ASP	2.3
1	A	74	PRO	2.3
4	I	175	TYR	2.3
1	A	26	GLY	2.3
2	B	120	PHE	2.3
1	A	154	LEU	2.3
4	I	65	GLY	2.3
4	I	145	GLY	2.3
1	A	176	HIS	2.3
4	I	30	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	I	112	VAL	2.3
4	I	140	VAL	2.3
1	A	87[A]	ASN	2.3
1	A	65	ALA	2.3
4	I	180	SER	2.3
1	A	268	GLU	2.3
4	I	71	LYS	2.3
4	I	126	VAL	2.3
4	I	178	LEU	2.2
1	A	95	TYR	2.2
4	I	32	TRP	2.2
4	I	170	LYS	2.2
2	B	127	VAL	2.2
2	B	177	LEU	2.2
4	I	83	VAL	2.2
1	A	82	GLY	2.2
2	B	17	GLY	2.2
1	A	25	LYS	2.2
4	I	203	SER	2.2
2	B	13	ILE	2.2
4	I	77	ILE	2.2
4	I	214	ILE	2.2
1	A	117	SER	2.2
4	I	50	LEU	2.2
1	A	77	ASP	2.2
2	B	226	ASN	2.2
4	I	115	ASP	2.2
4	I	109	SER	2.2
2	B	203	PHE	2.1
4	I	7	GLN	2.1
2	B	181	GLU	2.1
2	B	128	PHE	2.1
4	I	29	GLN	2.1
4	I	169	GLY	2.1
2	B	196	SER	2.1
4	I	209	GLU	2.1
4	I	85	ILE	2.1
2	B	78	ARG	2.1
1	A	29	GLN	2.1
2	B	131	ILE	2.1
1	A	188	VAL	2.1
4	I	177	THR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	273	LEU	2.0
1	A	71	ILE	2.0
1	A	79	PRO	2.0
4	I	119	TYR	2.0
1	A	144	VAL	2.0
2	B	117	VAL	2.0
2	B	45	ALA	2.0
4	I	210	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	1PE	A	1276	16/16	0.89	0.12	56,62,77,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.