



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:48 AM UTC

PDB ID : 4JEU / pdb_00004jeu
Title : Crystal Structure of Munc18a and Syntaxin1 with native N-terminus complex
Authors : Colbert, K.N.; Hattendorf, D.A.; Weiss, T.M.; Burkhardt, P.; Fasshauer, D.; Weis, W.I.
Deposited on : 2013-02-27
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	590	
2	B	242	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6303 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Syntaxin-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	558	Total	C	N	O	S	0	0	0
			4472	2836	759	852	25			


- Molecule 2 is a protein called Syntaxin-1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	225	Total	C	N	O	S	0	0	0
			1831	1121	323	376	11			


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Syntaxin-binding protein 1



• Molecule 2: Syntaxin-1A



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	156.78Å 156.78Å 78.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	110.86 – 3.20 110.86 – 3.20	Depositor EDS
% Data completeness (in resolution range)	(Not available) (110.86-3.20) 99.9 (110.86-3.20)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 3.19Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.8.0, BUSTER 2.8.0	Depositor
R, R_{free}	0.199 , 0.263 0.208 , 0.278	Depositor DCC
R_{free} test set	1342 reflections (3.96%)	wwPDB-VP
Wilson B-factor (Å ²)	81.2	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 112.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6303	wwPDB-VP
Average B, all atoms (Å ²)	97.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.89	0/4551	1.45	32/6143 (0.5%)
2	B	0.83	0/1848	1.55	16/2468 (0.6%)
All	All	0.88	0/6399	1.48	48/8611 (0.6%)

There are no bond length outliers.

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	363	ASP	CA-C-N	7.82	131.09	120.38
1	A	363	ASP	C-N-CA	7.82	131.09	120.38
1	A	255	ASP	CA-CB-CG	7.24	119.84	112.60
1	A	34	ASP	CA-CB-CG	7.05	119.65	112.60
1	A	98	LYS	N-CA-C	6.82	118.50	111.14
1	A	54	GLY	N-CA-C	6.73	124.61	115.32
1	A	47	MET	CA-C-N	6.50	129.28	120.38
1	A	47	MET	C-N-CA	6.50	129.28	120.38
1	A	52	THR	N-CA-C	-6.27	105.46	113.23
1	A	282	ASP	CA-CB-CG	5.90	118.50	112.60
1	A	17	ASP	CA-CB-CG	5.86	118.46	112.60
2	B	204	LYS	CA-C-N	5.86	128.41	120.44
2	B	204	LYS	C-N-CA	5.86	128.41	120.44
2	B	128	VAL	CA-C-N	5.72	127.95	120.28
2	B	128	VAL	C-N-CA	5.72	127.95	120.28
2	B	43	PHE	CA-CB-CG	5.57	119.37	113.80
2	B	72	LYS	CA-C-N	5.53	127.68	120.28
2	B	72	LYS	C-N-CA	5.53	127.68	120.28
1	A	407	ILE	CA-C-N	5.41	127.86	120.46
1	A	407	ILE	C-N-CA	5.41	127.86	120.46
2	B	145	CYS	CA-C-N	5.40	127.46	120.44
2	B	145	CYS	C-N-CA	5.40	127.46	120.44
2	B	102	GLN	CA-C-N	5.40	127.52	120.28
2	B	102	GLN	C-N-CA	5.40	127.52	120.28
1	A	390	ILE	CA-C-N	5.37	125.58	120.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	390	ILE	C-N-CA	5.37	125.58	120.43
1	A	471	GLN	CA-C-N	5.35	128.94	121.24
1	A	471	GLN	C-N-CA	5.35	128.94	121.24
1	A	429	HIS	CA-C-N	5.29	128.10	120.38
1	A	429	HIS	C-N-CA	5.29	128.10	120.38
1	A	150	ALA	CA-C-N	5.25	128.29	120.31
1	A	150	ALA	C-N-CA	5.25	128.29	120.31
1	A	339	LYS	CA-C-N	5.25	127.58	120.65
1	A	339	LYS	C-N-CA	5.25	127.58	120.65
1	A	151	ASP	CA-C-N	5.22	128.00	120.38
1	A	151	ASP	C-N-CA	5.22	128.00	120.38
2	B	97	GLU	CA-C-N	5.21	127.52	120.44
2	B	97	GLU	C-N-CA	5.21	127.52	120.44
2	B	35	GLU	N-CA-C	-5.20	105.62	111.28
1	A	119	VAL	N-CA-CB	5.15	117.16	110.57
1	A	434	PRO	CA-C-N	5.14	127.42	120.38
1	A	434	PRO	C-N-CA	5.14	127.42	120.38
2	B	136	ALA	CA-C-N	5.09	127.10	120.28
2	B	136	ALA	C-N-CA	5.09	127.10	120.28
1	A	66	GLU	CB-CG-CD	5.03	121.16	112.60
1	A	586	ASN	CA-CB-CG	5.03	117.63	112.60
1	A	227	ARG	CA-C-N	5.01	127.83	120.71
1	A	227	ARG	C-N-CA	5.01	127.83	120.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4472	0	4563	52	0
2	B	1831	0	1807	20	0
All	All	6303	0	6370	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:LEU:HD11	1:A:338:GLN:HG2	1.74	0.68
1:A:188:ALA:HB3	1:A:229:GLN:HG2	1.76	0.67
1:A:480:PRO:HG2	1:A:552:CYS:SG	2.34	0.67
1:A:39:ARG:NH1	1:A:148:ASP:OD1	2.28	0.66
2:B:181:ILE:HD11	2:B:192:LEU:HD12	1.85	0.59
1:A:256:LEU:HD22	1:A:357:HIS:HB3	1.85	0.58
2:B:131:MET:HB2	2:B:219:MET:HE1	1.87	0.56
1:A:242:PRO:HB2	1:A:578:LEU:HD22	1.87	0.56
1:A:169:LEU:HD22	1:A:201:LEU:HB2	1.88	0.55
1:A:119:VAL:HG12	2:B:8:LEU:HD23	1.88	0.55
1:A:47:MET:O	1:A:51:MET:HG3	2.07	0.54
1:A:3:PRO:HB3	1:A:532:ARG:HH22	1.73	0.54
1:A:152:SER:HB3	1:A:156:PHE:CE2	2.42	0.54
2:B:214:ASP:O	2:B:218:ASP:HB2	2.07	0.54
1:A:39:ARG:HH21	1:A:261:ASN:HD21	1.55	0.54
1:A:64:ARG:HB3	2:B:242:ASP:HB3	1.90	0.53
1:A:61:ILE:HG12	1:A:86:SER:HB3	1.89	0.53
1:A:362:VAL:HA	1:A:365:LEU:HD12	1.91	0.52
1:A:133:ILE:HD12	1:A:135:ILE:HD12	1.92	0.51
1:A:368:VAL:HG21	1:A:393:ILE:HD11	1.93	0.50
2:B:82:ILE:HD13	2:B:134:TYR:CE1	2.46	0.50
1:A:573:LEU:HD22	1:A:578:LEU:HB2	1.94	0.50
1:A:212:TYR:HB3	1:A:219:MET:HE1	1.94	0.49
1:A:256:LEU:HD21	1:A:362:VAL:HG21	1.93	0.49
2:B:160:THR:HG21	2:B:168:MET:HE2	1.93	0.49
1:A:239:PRO:HB2	1:A:413:PHE:CZ	2.48	0.48
1:A:399:VAL:HG12	1:A:404:LYS:HG3	1.95	0.48
2:B:148:ARG:HD2	2:B:198:ARG:HE	1.79	0.48
1:A:39:ARG:O	1:A:43:SER:HB2	2.13	0.48
2:B:173:ASN:HB3	2:B:176:ILE:HB	1.96	0.48
1:A:393:ILE:HG23	1:A:399:VAL:HG11	1.95	0.48
1:A:403:ASP:O	1:A:407:ILE:HG12	2.14	0.47
1:A:295:HIS:O	1:A:299:VAL:HG23	2.14	0.47
1:A:42:SER:OG	2:B:234:GLU:HB2	2.14	0.47
1:A:84:VAL:HB	1:A:114:LEU:HD22	1.97	0.47
1:A:47:MET:HE1	1:A:58:VAL:HG23	1.97	0.47
1:A:246:GLU:HG3	1:A:548:ASN:HB3	1.98	0.46
1:A:338:GLN:HG3	2:B:220:ALA:HB1	1.98	0.46
2:B:37:VAL:HG13	2:B:123:LEU:HD11	1.98	0.45
2:B:220:ALA:HA	2:B:223:VAL:HG22	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:199:HIS:CE1	2:B:203:ILE:HD11	2.52	0.45
1:A:174:GLU:O	1:A:178:THR:HG23	2.17	0.45
2:B:213:HIS:HE1	2:B:217:MET:HE2	1.81	0.45
1:A:261:ASN:O	1:A:263:VAL:HG23	2.18	0.44
1:A:361:THR:HG23	1:A:399:VAL:HG22	1.98	0.44
2:B:108:ARG:O	2:B:113:LEU:HD22	2.18	0.44
2:B:241:VAL:HG22	2:B:242:ASP:H	1.83	0.44
1:A:234:ASP:O	1:A:237:PHE:HD2	2.00	0.44
1:A:192:ARG:HB3	1:A:198:ASN:ND2	2.34	0.43
2:B:127:PHE:O	2:B:130:VAL:HG12	2.19	0.43
1:A:540:PHE:HA	1:A:568:GLY:O	2.18	0.43
1:A:584:LYS:HA	1:A:587:LYS:HG3	2.01	0.43
1:A:405:ILE:HD13	1:A:439:ILE:HD12	2.01	0.43
1:A:532:ARG:NH1	1:A:533:SER:HB2	2.34	0.43
1:A:444:ALA:HA	1:A:448:VAL:O	2.19	0.42
1:A:328:SER:O	1:A:330:MET:N	2.50	0.42
1:A:30:VAL:HG13	1:A:57:ILE:HG23	2.02	0.42
1:A:103:HIS:ND1	1:A:129:THR:HG22	2.34	0.42
1:A:43:SER:HB3	1:A:139:PRO:HD2	2.01	0.42
1:A:292:ARG:C	1:A:294:LYS:H	2.28	0.41
1:A:119:VAL:HG12	2:B:8:LEU:HA	2.03	0.41
1:A:77:ILE:O	1:A:106:PHE:HA	2.21	0.41
1:A:545:VAL:HG21	1:A:567:ILE:HG12	2.03	0.41
2:B:103:GLU:HA	2:B:106:LEU:HD12	2.01	0.41
1:A:279:VAL:HG11	1:A:346:THR:CG2	2.51	0.41
1:A:150:ALA:HA	1:A:571:HIS:ND1	2.35	0.40
1:A:574:THR:O	1:A:575:PRO:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	552/590 (94%)	497 (90%)	49 (9%)	6 (1%)	11	43
2	B	221/242 (91%)	210 (95%)	10 (4%)	1 (0%)	24	59
All	All	773/832 (93%)	707 (92%)	59 (8%)	7 (1%)	14	47

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	329	GLN
1	A	4	ILE
1	A	326	ASP
1	A	162	ALA
1	A	359	GLN
1	A	385	ASP
2	B	174	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	505/530 (95%)	484 (96%)	21 (4%)	26	60
2	B	207/223 (93%)	191 (92%)	16 (8%)	12	41
All	All	712/753 (95%)	675 (95%)	37 (5%)	21	54

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	MET
1	A	48	THR
1	A	86	SER
1	A	131	THR
1	A	138	LEU
1	A	168	ILE
1	A	204	LEU
1	A	230	LEU
1	A	279	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	316	MET
1	A	348	LEU
1	A	382	LYS
1	A	400	SER
1	A	401	THR
1	A	418	ILE
1	A	423	LEU
1	A	482	ILE
1	A	496	THR
1	A	539	ILE
1	A	565	VAL
1	A	584	LYS
2	B	7	GLU
2	B	9	ARG
2	B	38	GLU
2	B	58	HIS
2	B	66	ASN
2	B	91	SER
2	B	96	ILE
2	B	132	SER
2	B	165	LEU
2	B	176	ILE
2	B	181	ILE
2	B	189	LYS
2	B	192	LEU
2	B	205	LEU
2	B	209	ILE
2	B	222	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	206	GLN
1	A	261	ASN
1	A	329	GLN
1	A	558	GLN
1	A	560	ASN
1	A	586	ASN
2	B	36	GLN
2	B	50	ASN
2	B	58	HIS
2	B	66	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	135	ASN
2	B	138	GLN
2	B	150	GLN
2	B	152	GLN
2	B	213	HIS
2	B	226	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	558/590 (94%)	-0.15	1 (0%) 91 85	50, 84, 159, 201	0
2	B	225/242 (92%)	-0.19	0 100 100	61, 96, 201, 244	0
All	All	783/832 (94%)	-0.17	1 (0%) 92 89	50, 89, 168, 244	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	324	MET	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.