



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 05:26 AM UTC

PDB ID : 8JE7 / pdb_00008je7
Title : Crystal Structure of Anopheles culicifacies Prolyl-tRNA Synthetase (AcPRS)
in complex with two inhibitors (Halofuginone and L95)
Authors : Goswami, R.; Manickam, Y.; Chhibber-Goel, J.; Harlos, K.; Sharma, A.
Deposited on : 2023-05-15
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3689 (2.16-2.12)
Clashscore	190562	3812 (2.16-2.12)
Ramachandran outliers	187476	3773 (2.16-2.12)
Sidechain outliers	187428	3772 (2.16-2.12)
RSRZ outliers	180081	3691 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	
1	B	505	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 8279 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROLYL-TRNA SYNTHETASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	485	Total	C	N	O	S	0	1	0
			3851	2462	661	708	20			
1	B	491	Total	C	N	O	S	0	4	0
			3941	2524	676	722	19			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1218	MET	-	initiating methionine	UNP A0A182M1F8
B	1218	MET	-	initiating methionine	UNP A0A182M1F8

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is 7-bromo-6-chloro-3-{3-[(2R,3S)-3-hydroxypiperidin-2-yl]-2-oxopropyl}quinazolin-4(3H)-one (CCD ID: HFG) (formula: C₁₆H₁₇BrClN₃O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	Br	C	Cl	N	O	0	0
			24	1	16	1	3	3		
3	B	1	Total	Br	C	Cl	N	O	0	0
			24	1	16	1	3	3		

- Molecule 4 is {N}-[4-[(3 {S})-3-cyano-3-cyclopropyl-2-oxidanylidene-pyrrolidin-1-yl]-6-methyl-pyridin-2-yl]-2-phenyl-ethanamide (CCD ID: JE6) (formula: C₂₂H₂₂N₄O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			28	22	4	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			28	22	4	2		

- Molecule 5 is BROMIDE ION (CCD ID: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Br	0	0
			2	2		
5	B	2	Total	Br	0	0
			2	2		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total 1	Cl 1	0	0
7	B	2	Total 3	Cl 3	0	1


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	136	Total 139	O 139	0	3
8	B	208	Total 210	O 210	0	3


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROLYL-TRNA SYNTHETASE



• Molecule 1: PROLYL-TRNA SYNTHETASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	83.67Å 113.71Å 124.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	83.81 – 2.14 83.81 – 2.14	Depositor EDS
% Data completeness (in resolution range)	99.2 (83.81-2.14) 99.3 (83.81-2.14)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 2.14Å)	Xtriage
Refinement program	PHENIX (1.15rc1_3423: ???)	Depositor
R, R_{free}	0.180 , 0.228 0.186 , 0.228	Depositor DCC
R_{free} test set	3246 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	46.4	Xtriage
Anisotropy	0.282	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 42.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8279	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: JE6, BR, CL, GOL, ZN, HFG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	7/3953 (0.2%)	0.66	3/5372 (0.1%)
1	B	0.57	4/4056 (0.1%)	0.70	8/5507 (0.1%)
All	All	0.59	11/8009 (0.1%)	0.68	11/10879 (0.1%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1376	TRP	C-O	-6.48	1.16	1.23
1	B	1684	LYS	C-O	-6.40	1.15	1.23
1	A	1369	LEU	C-O	-5.90	1.17	1.24
1	B	1365	PRO	C-O	-5.82	1.16	1.23
1	A	1370	ARG	C-O	-5.73	1.17	1.24
1	A	1438	ALA	C-O	-5.70	1.17	1.24
1	A	1646	LYS	C-O	-5.48	1.16	1.23
1	B	1368	PHE	C-O	-5.26	1.17	1.24
1	A	1353	GLN	C-O	-5.21	1.17	1.24
1	A	1714	PHE	C-O	-5.12	1.17	1.23
1	A	1441	ARG	C-O	-5.10	1.17	1.23

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1258[A]	HIS	CA-C-O	8.53	129.75	120.20
1	B	1258[B]	HIS	CA-C-O	8.53	129.75	120.20
1	B	1258[A]	HIS	N-CA-C	7.59	121.47	111.75
1	B	1258[B]	HIS	N-CA-C	7.59	121.47	111.75
1	A	1369	LEU	N-CA-C	6.91	119.84	111.40
1	B	1521	THR	CA-C-N	6.50	133.96	121.54
1	B	1521	THR	C-N-CA	6.50	133.96	121.54
1	B	1369	LEU	N-CA-C	6.08	117.99	111.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1521	THR	O-C-N	5.89	128.13	122.07
1	A	1297	GLU	N-CA-C	5.29	117.73	111.33
1	A	1439	SER	CB-CA-C	-5.03	102.29	110.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3851	0	3670	44	0
1	B	3941	0	3793	30	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	24	0	17	2	0
3	B	24	0	17	0	0
4	A	28	0	0	0	0
4	B	28	0	0	1	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	12	0	16	2	0
6	B	12	0	15	0	0
7	A	1	0	0	0	0
7	B	3	0	0	2	0
8	A	139	0	0	1	0
8	B	210	0	0	3	0
All	All	8279	0	7528	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1439:SER:HB3	1:A:1441:ARG:HG3	1.55	0.88

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1575:LEU:HD12	1:A:1579:ASP:HB2	1.64	0.79
1:A:1507:CYS:SG	8:A:2017:HOH:O	2.48	0.71
1:A:1575:LEU:HD12	1:A:1579:ASP:CB	2.30	0.62
1:B:1697:ASP:HB3	1:B:1700:VAL:HG22	1.87	0.56
1:A:1632:LEU:HD11	1:A:1715:TYR:CE1	2.40	0.56
1:A:1643:LEU:HD11	1:A:1720:ARG:HG3	1.87	0.55
1:B:1464:GLU:OE2	1:B:1469:LYS:HD2	2.07	0.54
1:A:1686:LEU:HD12	1:A:1717:LEU:HG	1.90	0.54
1:A:1410:ILE:HD13	1:A:1488:VAL:HG12	1.90	0.54
1:B:1331:THR:HB	8:B:1973:HOH:O	2.07	0.53
1:A:1650:MET:HE1	1:A:1689:PRO:HG2	1.90	0.53
1:A:1287:ILE:HD12	1:B:1356:ASN:HD21	1.75	0.52
1:A:1638:GLU:HG2	1:A:1642:PHE:CE2	2.46	0.51
1:A:1381:THR:HG21	1:A:1392:GLU:OE1	2.11	0.51
1:A:1393:VAL:HG21	1:A:1450:HIS:HB2	1.92	0.51
1:A:1354:TRP:CE3	1:A:1377:GLN:HB3	2.45	0.51
1:A:1533:CYS:SG	1:A:1575:LEU:HD23	2.52	0.50
1:B:1370:ARG:HH12	1:B:1722:TYR:C	2.19	0.50
1:A:1435:TYR:HE2	1:A:1629:HIS:ND1	2.10	0.49
1:B:1296[A]:ARG:HD2	1:B:1459:PHE:CE1	2.48	0.49
1:A:1376:TRP:CD1	1:A:1376:TRP:C	2.91	0.48
1:A:1646:LYS:HE3	1:A:1646:LYS:HB2	1.54	0.48
1:A:1223:LEU:HD13	1:A:1566:LEU:CD2	2.43	0.48
1:B:1714:PHE:CE1	1:B:1716:THR:OG1	2.67	0.48
1:A:1410:ILE:HD11	1:A:1489:MET:HA	1.96	0.48
1:A:1592:ASN:HB3	1:A:1594:ALA:H	1.79	0.48
1:B:1227:LYS:HE3	1:B:1234:TRP:CH2	2.48	0.48
1:B:1416:ARG:HD2	8:B:1915:HOH:O	2.12	0.47
1:B:1354:TRP:CE3	1:B:1377:GLN:HB3	2.49	0.47
1:A:1284:TYR:CE1	6:A:1806:GOL:H12	2.50	0.46
1:A:1650:MET:CE	1:A:1689:PRO:HG2	2.45	0.46
1:B:1686:LEU:HD12	1:B:1717:LEU:HG	1.97	0.46
1:B:1370:ARG:HD3	4:B:1802:JE6:C10	2.46	0.46
1:A:1513:VAL:HG13	1:A:1560:LYS:HD3	1.97	0.45
1:B:1296[A]:ARG:HH11	1:B:1459:PHE:HE1	1.64	0.45
1:B:1643:LEU:HD22	1:B:1683:ALA:HB2	1.98	0.45
1:A:1645:THR:HB	1:A:1647:ASN:HD22	1.81	0.45
1:A:1591:ASP:OD1	1:A:1592:ASN:N	2.50	0.44
1:A:1310:TRP:HB2	1:B:1313:LYS:HG3	1.99	0.44
1:A:1632:LEU:HD11	1:A:1715:TYR:HE1	1.83	0.43
1:B:1344[A]:TYR:CG	1:B:1466:PRO:HG3	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1465:HIS:CE1	1:B:1467:GLU:HB3	2.54	0.43
1:A:1550:ASP:CG	1:A:1560:LYS:HE2	2.42	0.43
1:B:1381:THR:HG21	1:B:1392:GLU:OE1	2.18	0.43
1:A:1350:ARG:HA	1:A:1381:THR:HG22	2.01	0.43
1:A:1435:TYR:HE2	1:A:1629:HIS:HD1	1.66	0.42
1:B:1441:ARG:NH1	1:B:1720:ARG:HB3	2.33	0.42
1:A:1356:ASN:OD1	6:A:1806:GOL:O1	2.37	0.42
1:A:1636:TRP:NE1	1:A:1640:LEU:HD11	2.34	0.42
1:A:1532:SER:O	1:A:1600:ARG:NH2	2.51	0.42
1:B:1258[B]:HIS:ND1	7:B:1808:CL:CL	2.83	0.42
1:A:1580:LEU:HD12	1:A:1580:LEU:HA	1.77	0.42
1:A:1695:LYS:HE2	1:A:1695:LYS:HB3	1.93	0.42
1:A:1423:PHE:CE1	3:A:1801:HFG:H15	2.55	0.42
1:A:1511:VAL:HA	1:A:1548:GLU:O	2.20	0.42
1:B:1714:PHE:HE1	1:B:1716:THR:OG1	2.03	0.42
1:A:1330:GLU:OE1	3:A:1801:HFG:N1'	2.53	0.41
1:B:1411:PRO:HG3	1:B:1623:GLU:HA	2.02	0.41
1:B:1471:LYS:HE3	1:B:1471:LYS:HB3	1.74	0.41
7:B:1807[B]:CL:CL	8:B:2090:HOH:O	2.59	0.41
1:A:1650:MET:HB3	1:A:1650:MET:HE2	1.69	0.41
1:B:1304:PHE:C	1:B:1306:PRO:HD3	2.46	0.41
1:B:1394:LEU:HD23	1:B:1394:LEU:HA	1.88	0.41
1:A:1431:THR:OG1	1:A:1444:GLN:HG3	2.20	0.41
1:A:1554:ASN:ND2	1:B:1345:ARG:HD3	2.35	0.41
1:A:1441:ARG:HH11	1:A:1441:ARG:HD3	1.73	0.40
1:B:1423:PHE:HB2	1:B:1449:HIS:CE1	2.56	0.40
1:B:1532:SER:OG	1:B:1600:ARG:NH1	2.52	0.40
1:A:1549:GLY:HA3	1:A:1551:TYR:CZ	2.56	0.40
1:B:1379:GLY:O	1:B:1478:SER:HA	2.20	0.40
1:B:1410:ILE:HD13	1:B:1488:VAL:HG12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	480/505 (95%)	468 (98%)	12 (2%)	0	100	100
1	B	491/505 (97%)	478 (97%)	12 (2%)	1 (0%)	43	42
All	All	971/1010 (96%)	946 (97%)	24 (2%)	1 (0%)	48	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1522	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/433 (91%)	392 (100%)	1 (0%)	86	90
1	B	406/433 (94%)	400 (98%)	6 (2%)	57	63
All	All	799/866 (92%)	792 (99%)	7 (1%)	73	77

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1377	GLN
1	B	1258[A]	HIS
1	B	1258[B]	HIS
1	B	1376	TRP
1	B	1377	GLN
1	B	1523	ASP
1	B	1525	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1476	GLN
1	A	1562	ASN
1	A	1602	GLN
1	A	1647	ASN
1	B	1386	GLN
1	B	1497	GLN
1	B	1562	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 18 ligands modelled in this entry, 10 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	B	1806	-	5,5,5	0.99	0	5,5,5	1.23	1 (20%)
6	GOL	A	1806	-	5,5,5	1.00	0	5,5,5	1.11	0
3	HFG	A	1801	-	26,26,26	0.39	0	29,37,37	0.73	1 (3%)
4	JE6	A	1802	-	29,31,31	0.76	0	42,45,45	0.75	2 (4%)
6	GOL	A	1805	-	5,5,5	0.51	0	5,5,5	1.03	0
4	JE6	B	1802	-	29,31,31	0.18	0	42,45,45	0.71	1 (2%)
3	HFG	B	1801	-	26,26,26	0.38	0	29,37,37	0.92	1 (3%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	B	1805	-	5,5,5	1.34	1 (20%)	5,5,5	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	B	1806	-	-	0/4/4/4	-
6	GOL	A	1806	-	-	0/4/4/4	-
3	HFG	A	1801	-	-	0/8/19/19	0/3/3/3
4	JE6	A	1802	-	-	0/16/39/39	0/4/4/4
6	GOL	A	1805	-	-	0/4/4/4	-
4	JE6	B	1802	-	-	0/16/39/39	0/4/4/4
3	HFG	B	1801	-	-	0/8/19/19	0/3/3/3
6	GOL	B	1805	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	1805	GOL	O2-C2	-2.40	1.36	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1801	HFG	C6'-N1'-C2'	-3.91	109.18	111.62
4	B	1802	JE6	C15-C16-C17	3.32	116.63	111.33
4	A	1802	JE6	C15-C16-C21	3.21	105.94	102.35
3	A	1801	HFG	C6'-N1'-C2'	-2.83	109.86	111.62
6	B	1806	GOL	C3-C2-C1	-2.28	103.42	111.80
4	A	1802	JE6	C15-C16-C17	2.13	114.73	111.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1806	GOL	2	0
3	A	1801	HFG	2	0

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1802	JE6	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand JE6 A 1802



Ligand JE6 B 1802





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	485/505 (96%)	0.12	13 (2%) 56 60	30, 58, 94, 128	1 (0%)
1	B	491/505 (97%)	-0.11	17 (3%) 47 51	23, 49, 88, 124	4 (0%)
All	All	976/1010 (96%)	0.00	30 (3%) 51 55	23, 53, 93, 128	5 (0%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1669	ARG	5.6
1	A	1722	TYR	4.1
1	A	1298	LYS	3.8
1	B	1518	THR	3.7
1	A	1305	ALA	3.4
1	A	1524	ASP	3.3
1	A	1712	ALA	3.2
1	B	1524	ASP	3.1
1	A	1344	TYR	2.9
1	B	1521	THR	2.8
1	A	1468	THR	2.8
1	B	1722	TYR	2.6
1	A	1222	GLY	2.6
1	A	1636	TRP	2.6
1	B	1520	THR	2.5
1	B	1577	PHE	2.5
1	A	1632	LEU	2.3
1	B	1680	ALA	2.3
1	B	1229	ASP	2.3
1	B	1714	PHE	2.2
1	B	1522	THR	2.2
1	A	1592	ASN	2.2
1	B	1681	MET	2.2
1	A	1520	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1578	LYS	2.2
1	B	1301	ILE	2.1
1	B	1517	ILE	2.1
1	B	1467	GLU	2.1
1	A	1639	PHE	2.1
1	B	1528	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	A	1806	6/6	0.80	0.21	83,88,92,92	0
5	BR	B	1804	1/1	0.88	0.13	102,102,102,102	1
7	CL	B	1808	1/1	0.90	0.25	46,46,46,46	1
7	CL	A	1807	1/1	0.92	0.10	70,70,70,70	0
3	HFG	B	1801	24/24	0.93	0.08	27,37,45,128	24
5	BR	A	1804	1/1	0.94	0.09	108,108,108,108	1
3	HFG	A	1801	24/24	0.95	0.07	39,47,59,145	0
6	GOL	B	1806	6/6	0.95	0.07	52,71,73,80	0
6	GOL	A	1805	6/6	0.96	0.06	41,43,44,46	0
4	JE6	A	1802	28/28	0.96	0.07	32,43,49,55	0
6	GOL	B	1805	6/6	0.96	0.08	35,39,46,56	0
4	JE6	B	1802	28/28	0.97	0.06	23,35,40,43	28
2	ZN	A	1800	1/1	0.99	0.03	80,80,80,80	0
5	BR	B	1803	1/1	0.99	0.12	59,59,59,59	1
7	CL	B	1807[A]	1/1	0.99	0.06	53,53,53,53	1
7	CL	B	1807[B]	1/1	0.99	0.06	62,62,62,62	1
5	BR	A	1803	1/1	0.99	0.08	48,48,48,48	1

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	B	1800	1/1	1.00	0.03	41,41,41,41	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around HFG A 1801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around JE6 A 1802:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.