



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:17 PM UTC

PDB ID : 8JE9 / pdb_00008je9
Title : Crystal structure of CGL1 from Crassostrea gigas, mannobiose-bound form (CGL1/Man(alpha)1-2Man)
Authors : Unno, H.; Hatakeyama, T.
Deposited on : 2023-05-15
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	Y	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACY	C	201	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Natterin-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	142	Total	C	N	O	S	0	4	0
			1108	713	184	207	4			
1	B	142	Total	C	N	O	S	0	3	0
			1104	710	185	205	4			
1	C	142	Total	C	N	O	S	0	2	0
			1101	709	183	205	4			
1	D	142	Total	C	N	O	S	0	2	0
			1097	705	183	205	4			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	X	2	Total	C	O	0	0	0
			23	12	11			
2	Y	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is ACETIC ACID (CCD ID: ACY) (formula: C₂H₄O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is alpha-D-mannopyranose (CCD ID: MAN) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			12	6	6		

- Molecule 5 is CACODYLATE ION (CCD ID: CAC) (formula: $C_2H_6AsO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	As	C	O	0	0
			5	1	2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	275	Total	O	0	0
			275	275		
6	B	80	Total	O	0	0
			80	80		
6	C	438	Total	O	0	0
			438	438		
6	D	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Natterin-3





- Molecule 1: Natterin-3



- Molecule 1: Natterin-3



- Molecule 1: Natterin-3



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose

Chain Y: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.75Å 58.70Å 108.41Å 90.00° 93.70° 90.00°	Depositor
Resolution (Å)	39.78 – 1.00 39.78 – 1.00	Depositor EDS
% Data completeness (in resolution range)	85.8 (39.78-1.00) 85.8 (39.78-1.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 1.00Å)	Xtriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.174 , 0.187 0.183 , 0.195	Depositor DCC
R_{free} test set	11557 reflections (4.32%)	wwPDB-VP
Wilson B-factor (Å ²)	9.7	Xtriage
Anisotropy	0.333	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 30.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5287	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.25 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.2641e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, ACY, AYA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/1139	0.96	3/1544 (0.2%)
1	B	0.80	0/1132	1.01	6/1533 (0.4%)
1	C	0.92	3/1126 (0.3%)	1.16	10/1526 (0.7%)
1	D	0.83	0/1122	1.10	3/1521 (0.2%)
All	All	0.83	3/4519 (0.1%)	1.06	22/6124 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	85	ASN	C-O	-7.93	1.13	1.23
1	C	85	ASN	CG-OD1	6.61	1.36	1.23
1	C	84	SER	C-N	6.20	1.42	1.33

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	18	ARG	NE-CZ-NH1	-11.97	109.53	121.50
1	C	85	ASN	CA-CB-CG	-11.04	101.56	112.60
1	C	18	ARG	NE-CZ-NH2	10.91	129.02	119.20
1	D	18	ARG	NE-CZ-NH1	-8.57	112.93	121.50
1	B	18	ARG	NE-CZ-NH2	-7.99	112.01	119.20
1	A	73	ASN	CA-CB-CG	7.94	120.54	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	60	HIS	CB-CG-CD2	-7.42	121.56	131.20
1	C	85	ASN	CA-C-O	7.34	130.05	121.54
1	A	18	ARG	NE-CZ-NH1	-6.72	114.78	121.50
1	D	18	ARG	NE-CZ-NH2	6.70	125.23	119.20
1	C	85	ASN	O-C-N	-6.49	113.90	122.40
1	B	60	HIS	CB-CG-ND1	6.39	132.28	122.70
1	C	84	SER	O-C-N	-6.29	115.86	123.10
1	B	79	ASP	CA-CB-CG	-6.12	106.48	112.60
1	C	85	ASN	CA-C-N	6.07	133.71	121.93
1	C	85	ASN	C-N-CA	6.07	133.71	121.93
1	D	69	VAL	CA-CB-CG1	5.96	120.52	110.40
1	B	85	ASN	CA-CB-CG	-5.53	107.07	112.60
1	B	87	ASP	CA-CB-CG	5.25	117.85	112.60
1	A	18	ARG	NE-CZ-NH2	5.24	123.91	119.20
1	C	87	ASP	O-C-N	5.06	129.06	122.89
1	C	52	HIS	CB-CG-CD2	-5.00	124.69	131.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	84	SER	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1108	0	1099	7	0
1	B	1104	0	1097	1	0
1	C	1101	0	1090	5	0
1	D	1097	0	1085	4	0
2	X	23	0	21	0	0
2	Y	23	0	21	0	0
3	A	4	0	3	0	0
3	B	4	0	3	0	0
3	C	4	0	3	3	0
4	D	12	0	11	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	5	0	0	0	0
6	A	275	0	0	2	0
6	B	80	0	0	0	0
6	C	438	0	0	4	0
6	D	9	0	0	0	0
All	All	5287	0	4433	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:49:GLU:HG3	6:C:375:HOH:O	1.62	0.99
1:C:85:ASN:OD1	1:C:134:LYS:HB3	1.89	0.71
1:A:121:VAL:HG12	1:A:134:LYS:HD2	1.73	0.68
1:C:23[A]:ILE:HG12	6:C:434:HOH:O	1.96	0.64
3:C:201:ACY:H2	6:C:447:HOH:O	2.01	0.61
1:D:8[A]:THR:HG22	1:D:64:ASN:OD1	2.09	0.53
1:C:93:ILE:HD12	3:C:201:ACY:H3	1.91	0.52
1:A:13:PRO:HD2	1:A:67[A]:VAL:HG11	1.94	0.50
1:C:113:CYS:HB2	1:C:123:TYR:O	2.17	0.45
1:D:8[A]:THR:HA	1:D:63:GLN:O	2.17	0.45
1:A:85[B]:ASN:OD1	6:A:2101:HOH:O	2.21	0.43
3:C:201:ACY:CH3	6:C:447:HOH:O	2.64	0.43
1:A:13:PRO:CD	1:A:67[A]:VAL:HG11	2.50	0.42
1:B:113:CYS:HB2	1:B:123:TYR:O	2.19	0.41
1:A:121:VAL:HG12	1:A:134:LYS:CD	2.45	0.41
1:A:5:VAL:HG12	6:A:2205:HOH:O	2.20	0.41
1:D:113:CYS:HB2	1:D:123:TYR:O	2.21	0.41
1:D:8[B]:THR:HA	1:D:63:GLN:O	2.21	0.40
1:A:49[A]:GLU:OE2	1:A:61:ILE:HD13	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/143 (101%)	144 (100%)	0	0	100	100
1	B	143/143 (100%)	142 (99%)	1 (1%)	0	100	100
1	C	142/143 (99%)	141 (99%)	1 (1%)	0	100	100
1	D	142/143 (99%)	141 (99%)	1 (1%)	0	100	100
All	All	571/572 (100%)	568 (100%)	3 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/112 (103%)	115 (100%)	0	100	100
1	B	114/112 (102%)	113 (99%)	1 (1%)	70	36
1	C	113/112 (101%)	111 (98%)	2 (2%)	51	15
1	D	113/112 (101%)	112 (99%)	1 (1%)	70	36
All	All	455/448 (102%)	451 (99%)	4 (1%)	76	36

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	134	LYS
1	C	23[A]	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	23[B]	ILE
1	D	134	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	81	GLN
1	C	52	HIS
1	D	60	HIS
1	D	85	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	AYA	A	2	1	6,7,8	1.26	1 (16%)	6,8,10	0.86	0
1	AYA	D	2	1	6,7,8	1.05	1 (16%)	6,8,10	0.80	0
1	AYA	B	2	1	6,7,8	1.10	1 (16%)	6,8,10	0.52	0
1	AYA	C	2	1	6,7,8	0.76	0	6,8,10	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	AYA	A	2	1	-	0/5/6/8	-
1	AYA	D	2	1	-	1/5/6/8	-
1	AYA	B	2	1	-	0/5/6/8	-
1	AYA	C	2	1	-	1/5/6/8	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2	AYA	CA-N	-2.93	1.43	1.46
1	B	2	AYA	CA-N	-2.53	1.43	1.46
1	D	2	AYA	CA-N	-2.31	1.44	1.46

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	2	AYA	O-C-CA-CB
1	D	2	AYA	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MAN	X	1	2	12,12,12	0.75	0	17,17,17	0.69	0
2	MAN	X	2	2	11,11,12	0.47	0	15,15,17	0.73	0
2	MAN	Y	1	2	12,12,12	0.88	0	17,17,17	1.30	3 (17%)
2	MAN	Y	2	2	11,11,12	0.84	0	15,15,17	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	X	1	2	-	1/2/22/22	0/1/1/1
2	MAN	X	2	2	-	0/2/19/22	0/1/1/1
2	MAN	Y	1	2	-	0/2/22/22	0/1/1/1
2	MAN	Y	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Y	1	MAN	C3-C4-C5	2.84	115.38	110.23
2	Y	1	MAN	O3-C3-C4	2.56	116.42	110.38
2	Y	1	MAN	C1-O5-C5	-2.08	109.63	113.65

There are no chirality outliers.


All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	X	1	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	ACY	C	201	-	3,3,3	1.61	0	3,3,3	1.25	0
3	ACY	B	201	-	3,3,3	1.42	1 (33%)	3,3,3	1.03	0
5	CAC	D	1002	-	2,4,4	2.03	1 (50%)	4,6,6	4.12	2 (50%)
3	ACY	A	2001	-	3,3,3	0.81	0	3,3,3	0.60	0
4	MAN	D	1001	-	12,12,12	0.55	0	17,17,17	0.92	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	D	1001	-	-	0/2/22/22	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	1002	CAC	AS-C1	2.87	1.97	1.90
3	B	201	ACY	O-C	2.18	1.31	1.22

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	1002	CAC	O1-AS-C2	7.69	121.06	111.50
5	D	1002	CAC	O2-AS-C2	2.34	111.50	105.84
4	D	1001	MAN	O1-C1-O5	2.18	116.88	110.41
4	D	1001	MAN	O1-C1-C2	-2.06	103.01	108.98

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	201	ACY	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	141/143 (98%)	0.03	1 (0%) 84 88	5, 10, 16, 20	4 (2%)
1	B	141/143 (98%)	0.78	12 (8%) 16 17	6, 10, 17, 22	3 (2%)
1	C	141/143 (98%)	0.39	9 (6%) 25 27	6, 11, 20, 32	2 (1%)
1	D	141/143 (98%)	1.42	41 (29%) 1 1	7, 13, 22, 26	2 (1%)
All	All	564/572 (98%)	0.65	63 (11%) 10 9	5, 11, 19, 32	11 (1%)

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	85	ASN	7.2
1	C	86	GLY	6.2
1	D	89	PRO	4.4
1	C	118	GLY	4.3
1	D	56	ALA	4.1
1	D	100	TYR	4.0
1	B	106	TYR	4.0
1	C	87	ASP	3.8
1	B	127	ALA	3.7
1	D	5	VAL	3.7
1	D	92	ALA	3.6
1	C	49	GLU	3.5
1	D	88	VAL	3.5
1	C	97	SER	3.4
1	D	34	VAL	3.3
1	B	23	ILE	3.2
1	D	99	ILE	3.2
1	D	55	PHE	3.2
1	B	129	LYS	3.1
1	D	35	VAL	3.1
1	D	37	GLY	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	90	GLY	3.0
1	D	117	THR	3.0
1	D	140	TYR	2.9
1	B	128	GLY	2.9
1	C	88	VAL	2.9
1	D	138	ALA	2.8
1	D	72	ILE	2.7
1	D	143	ILE	2.7
1	D	97	SER	2.7
1	B	126	PHE	2.7
1	D	71	PRO	2.7
1	D	101	ILE	2.7
1	D	142	VAL	2.6
1	D	36	SER	2.6
1	D	127	ALA	2.5
1	C	84	SER	2.5
1	D	70	TYR	2.5
1	D	139	LEU	2.5
1	D	58	LYS	2.5
1	B	63	GLN	2.4
1	D	38	GLU	2.4
1	D	134	LYS	2.4
1	D	6	SER	2.4
1	D	115	ILE	2.4
1	B	79	ASP	2.4
1	D	123	TYR	2.4
1	D	98	GLY	2.4
1	A	73	ASN	2.4
1	D	74	ALA	2.3
1	C	23[A]	ILE	2.2
1	B	5	VAL	2.2
1	D	67	VAL	2.2
1	D	83	ALA	2.2
1	D	4	TRP	2.2
1	B	107[A]	SER	2.2
1	D	85	ASN	2.1
1	D	95	THR	2.1
1	D	60	HIS	2.1
1	B	57	GLY	2.0
1	D	121	VAL	2.0
1	B	97	SER	2.0
1	D	122	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	AYA	C	2	8/9	0.96	0.06	11,12,14,15	0
1	AYA	D	2	8/9	0.96	0.08	13,14,15,15	0
1	AYA	A	2	8/9	0.97	0.06	9,10,12,16	0
1	AYA	B	2	8/9	0.97	0.06	8,9,10,13	0

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	X	1	12/12	-	-	12,16,21,32	0
2	MAN	X	2	11/12	-	-	10,11,15,18	0
2	MAN	Y	1	12/12	-	-	12,16,21,24	0
2	MAN	Y	2	11/12	-	-	9,9,12,17	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain X:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ACY	B	201	4/4	0.84	0.13	22,22,26,26	0
3	ACY	A	2001	4/4	0.86	0.11	17,18,18,22	0
3	ACY	C	201	4/4	0.88	0.12	18,18,19,27	0
4	MAN	D	1001	12/12	0.94	0.09	10,12,17,19	0
5	CAC	D	1002	5/5	0.98	0.10	9,10,13,13	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around MAN D 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.