



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 08:48 AM UTC

PDB ID : 9JEJ / pdb_00009jej
Title : Crystal structure of human EP300 KIX domain (L644C mutant)
Authors : Lee, B.I.; Cho, S.E.
Deposited on : 2024-09-03
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	101	57% 10% 33%
1	B	101	53% 14% 33%
1	C	101	55% 11% 33%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1749 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone acetyltransferase p300.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	68	Total	C	N	O	S	0	0	0
			581	365	106	107	3			
1	B	68	Total	C	N	O	S	0	0	0
			581	365	106	107	3			
1	C	68	Total	C	N	O	S	0	0	0
			581	365	106	107	3			

There are 45 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	552	MET	-	initiating methionine	UNP Q09472
A	553	ARG	-	expression tag	UNP Q09472
A	554	GLY	-	expression tag	UNP Q09472
A	555	SER	-	expression tag	UNP Q09472
A	556	HIS	-	expression tag	UNP Q09472
A	557	HIS	-	expression tag	UNP Q09472
A	558	HIS	-	expression tag	UNP Q09472
A	559	HIS	-	expression tag	UNP Q09472
A	560	HIS	-	expression tag	UNP Q09472
A	561	HIS	-	expression tag	UNP Q09472
A	562	GLY	-	expression tag	UNP Q09472
A	563	MET	-	expression tag	UNP Q09472
A	564	ALA	-	expression tag	UNP Q09472
A	565	SER	-	expression tag	UNP Q09472
A	644	CYS	LEU	engineered mutation	UNP Q09472
B	552	MET	-	initiating methionine	UNP Q09472
B	553	ARG	-	expression tag	UNP Q09472
B	554	GLY	-	expression tag	UNP Q09472
B	555	SER	-	expression tag	UNP Q09472
B	556	HIS	-	expression tag	UNP Q09472
B	557	HIS	-	expression tag	UNP Q09472
B	558	HIS	-	expression tag	UNP Q09472
B	559	HIS	-	expression tag	UNP Q09472

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	560	HIS	-	expression tag	UNP Q09472
B	561	HIS	-	expression tag	UNP Q09472
B	562	GLY	-	expression tag	UNP Q09472
B	563	MET	-	expression tag	UNP Q09472
B	564	ALA	-	expression tag	UNP Q09472
B	565	SER	-	expression tag	UNP Q09472
B	644	CYS	LEU	engineered mutation	UNP Q09472
C	552	MET	-	initiating methionine	UNP Q09472
C	553	ARG	-	expression tag	UNP Q09472
C	554	GLY	-	expression tag	UNP Q09472
C	555	SER	-	expression tag	UNP Q09472
C	556	HIS	-	expression tag	UNP Q09472
C	557	HIS	-	expression tag	UNP Q09472
C	558	HIS	-	expression tag	UNP Q09472
C	559	HIS	-	expression tag	UNP Q09472
C	560	HIS	-	expression tag	UNP Q09472
C	561	HIS	-	expression tag	UNP Q09472
C	562	GLY	-	expression tag	UNP Q09472
C	563	MET	-	expression tag	UNP Q09472
C	564	ALA	-	expression tag	UNP Q09472
C	565	SER	-	expression tag	UNP Q09472
C	644	CYS	LEU	engineered mutation	UNP Q09472

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total Zn 1 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total O 2 2	0	0
3	B	1	Total O 1 1	0	0
3	C	2	Total O 2 2	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone acetyltransferase p300

Chain A: 




- Molecule 1: Histone acetyltransferase p300

Chain B: 



- Molecule 1: Histone acetyltransferase p300

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	96.34Å 96.34Å 136.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.26 – 2.90 31.26 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (31.26-2.90) 99.1 (31.26-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 2.90Å)	Xtriage
Refinement program	PHENIX (1.20.1_4487: ???)	Depositor
R, R_{free}	0.251 , 0.281 0.251 , 0.278	Depositor DCC
R_{free} test set	734 reflections (9.87%)	wwPDB-VP
Wilson B-factor (Å ²)	51.1	Xtriage
Anisotropy	0.080	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.479 for -1/2*h-1/2*k-1/2*l,-1/2*h-1/2*k+1/2*l,-h+k 0.479 for -1/2*h+1/2*k-1/2*l,1/2*h-1/2*k-1/2*l,-h-k	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	1749	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/590	0.28	0/790
1	B	0.12	0/590	0.26	0/790
1	C	0.13	0/590	0.30	0/790
All	All	0.13	0/1770	0.28	0/2370

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	581	0	571	5	0
1	B	581	0	571	8	0
1	C	581	0	571	8	0
2	C	1	0	0	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	C	2	0	0	0	0
All	All	1749	0	1713	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 19 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:575:ILE:HD11	1:B:626:ARG:HG3	1.82	0.61
1:A:575:ILE:HD11	1:A:626:ARG:HG3	1.82	0.61
1:C:619:MET:HE3	1:C:633:LEU:HD22	1.85	0.58
1:B:619:MET:HE1	1:B:633:LEU:HD13	1.85	0.57
1:C:575:ILE:HD11	1:C:626:ARG:HG3	1.87	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	64/101 (63%)	62 (97%)	2 (3%)	0	100	100
1	B	64/101 (63%)	64 (100%)	0	0	100	100
1	C	64/101 (63%)	64 (100%)	0	0	100	100
All	All	192/303 (63%)	190 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	61/88 (69%)	58 (95%)	3 (5%)	22	54

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	61/88 (69%)	59 (97%)	2 (3%)	33	67
1	C	61/88 (69%)	58 (95%)	3 (5%)	22	54
All	All	183/264 (69%)	175 (96%)	8 (4%)	25	59

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	646	GLU
1	C	640	ILE
1	B	646	GLU
1	B	640	ILE
1	C	591	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	585	HIS
1	C	585	HIS
1	C	589	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	68/101 (67%)	-1.01	0 100 100	38, 48, 92, 103	0
1	B	68/101 (67%)	-1.08	0 100 100	40, 48, 93, 101	0
1	C	68/101 (67%)	-1.06	0 100 100	38, 48, 91, 98	0
All	All	204/303 (67%)	-1.05	0 100 100	38, 48, 95, 103	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	C	701	1/1	0.99	0.05	66,66,66,66	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.