



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 01:53 PM EDT

PDB ID : 6JGT / pdb_00006jgt
Title : Crystal structure of barley exohydrolaseI W434Y mutant in complex with methyl 6-thio-beta-gentiobioside.
Authors : Luang, S.; Streltsov, V.A.; Hrmova, M.
Deposited on : 2019-02-14
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	609	3% 88% 9% ..
2	B	2	50% 50%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 5360 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-D-GLUCAN GLUCOHYDROLASE ISOENZYME EXO1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	606	Total	C	N	O	S	109	15	0
			4636	2926	809	872	29			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	HIS	-	expression tag	UNP A0A287SCR5
A	-2	HIS	-	expression tag	UNP A0A287SCR5
A	-1	ALA	-	expression tag	UNP A0A287SCR5
A	0	ALA	-	expression tag	UNP A0A287SCR5
A	320	LYS	ASN	conflict	UNP A0A287SCR5
A	434	TYR	TRP	engineered mutation	UNP A0A287SCR5

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-6)-methyl 6-thio-beta-D-glucopyranoside.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	S	0	0	0
			24	13	10	1			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: C₁₀H₂₂O₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			5	3	2		
5	A	1	Total	C	O	0	0
			10	6	4		
5	A	1	Total	C	O	0	0
			8	5	3		
5	A	1	Total	C	O	0	0
			5	3	2		
5	A	1	Total	C	O	0	0
			5	3	2		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			8	5	3		
5	A	1	Total	C	O	0	0
			5	3	2		
5	A	1	Total	C	O	0	0
			6	4	2		

- Molecule 6 is ACETATE ION (CCD ID: ACT) (formula: $\text{C}_2\text{H}_3\text{O}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is SULFATE ION (CCD ID: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	568	Total	O	0	0
			568	568		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BETA-D-GLUCAN GLUCOHYDROLASE ISOENZYME EXO1



• Molecule 2: beta-D-glucopyranose-(1-6)-methyl 6-thio-beta-D-glucopyranoside



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	100.71 Å 100.71 Å 181.01 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.56 – 1.82 48.56 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.56-1.82) 99.9 (48.56-1.82)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 1.82 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.142 , 0.162 0.152 , 0.173	Depositor DCC
R_{free} test set	4183 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	20.4	Xtriage
Anisotropy	0.142	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5360	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, 1PE, BGC, U1Y, ACT, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	7/4808 (0.1%)	0.99	13/6524 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	496	ASN	C-N	17.17	1.57	1.33
1	A	-2	HIS	C-N	8.31	1.45	1.33
1	A	220	GLU	C-N	-6.44	1.24	1.33
1	A	491	GLU	N-CA	6.03	1.53	1.46
1	A	491	GLU	CD-OE2	5.81	1.36	1.25
1	A	490	THR	C-N	-5.54	1.25	1.33
1	A	-2	HIS	N-CA	5.09	1.52	1.46

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	-2	HIS	O-C-N	-25.01	89.33	122.59
1	A	-2	HIS	CA-C-N	16.30	152.68	121.54
1	A	-2	HIS	C-N-CA	16.30	152.68	121.54
1	A	492	THR	N-CA-C	-10.68	99.61	111.14
1	A	496	ASN	CA-C-N	7.85	136.53	121.54
1	A	496	ASN	C-N-CA	7.85	136.53	121.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	ARG	NE-CZ-NH2	-6.15	113.67	119.20
1	A	495	ASP	N-CA-C	5.95	123.47	110.80
1	A	218	ILE	N-CA-C	5.46	116.10	110.36
1	A	101	ASN	N-CA-C	5.33	119.40	113.01
1	A	581	ASN	N-CA-C	5.17	117.83	109.40
1	A	339[A]	MET	CB-CG-SD	-5.02	97.64	112.70
1	A	339[B]	MET	CB-CG-SD	-5.02	97.64	112.70

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	-2	HIS	Mainchain,Peptide
1	A	-3	HIS	Peptide
1	A	491	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4636	0	4605	46	0
2	B	24	0	10	0	0
3	A	14	0	13	0	0
4	A	30	0	40	1	0
5	A	59	0	67	9	0
6	A	24	0	18	2	0
7	A	5	0	0	0	0
8	A	568	0	0	9	0
All	All	5360	0	4753	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:577:GLN:HB3	1:A:580[B]:MET:HE3	1.47	0.96
1:A:466[A]:ASN:ND2	1:A:508:THR:OG1	1.98	0.94
1:A:394:ASN:HD21	1:A:404:LEU:H	1.23	0.85
1:A:577:GLN:HB3	1:A:580[B]:MET:CE	2.06	0.85
5:A:712:1PE:C15	8:A:1231:HOH:O	2.32	0.77
1:A:577:GLN:NE2	1:A:580[B]:MET:HE1	2.00	0.77
1:A:100:GLN:HE21	1:A:100:GLN:HA	1.54	0.72
1:A:122:TYR:CE2	1:A:126:ARG:HD2	2.25	0.71
1:A:12:VAL:HB	5:A:713:1PE:H121	1.75	0.69
1:A:566[A]:ARG:NH2	8:A:805:HOH:O	2.26	0.68
1:A:262:HIS:HE1	1:A:285:ASP:H	1.40	0.67
1:A:181:GLN:HE21	1:A:203:ALA:H	1.42	0.66
1:A:497:LEU:O	1:A:497:LEU:HD23	2.00	0.61
1:A:331:HIS:HD2	8:A:1274:HOH:O	1.84	0.61
1:A:234:HIS:HD2	8:A:1111:HOH:O	1.84	0.60
1:A:167:ARG:HH21	1:A:171:GLN:HE22	1.48	0.60
1:A:97:VAL:H	1:A:101:ASN:HD21	1.48	0.59
1:A:100:GLN:HE21	1:A:100:GLN:CA	2.16	0.59
1:A:577:GLN:CD	1:A:580[B]:MET:HE1	2.27	0.58
1:A:262:HIS:CE1	1:A:285:ASP:H	2.22	0.57
1:A:58:GLY:H	1:A:102:ASN:ND2	2.04	0.54
1:A:134:GLU:OE2	1:A:377:HIS:HD2	1.93	0.52
1:A:291:ARG:HH22	5:A:708:1PE:C15	2.23	0.52
1:A:580[B]:MET:HE2	1:A:586:HIS:CD2	2.45	0.51
5:A:714:1PE:H151	8:A:1003:HOH:O	2.11	0.51
1:A:234:HIS:HE1	8:A:1277:HOH:O	1.95	0.49
1:A:155:ARG:HD2	1:A:497:LEU:HG	1.96	0.48
1:A:437:ASP:HB3	1:A:441:THR:HG21	1.95	0.48
1:A:440[A]:ARG:NH2	1:A:449:GLU:OE1	2.46	0.48
1:A:97:VAL:H	1:A:101:ASN:ND2	2.10	0.47
1:A:167:ARG:HH21	1:A:171:GLN:NE2	2.11	0.47
1:A:125:LYS:O	1:A:129[A]:GLU:HG3	2.14	0.47
1:A:407:PRO:HB2	5:A:712:1PE:C15	2.45	0.47
1:A:333:ASN:HD22	5:A:714:1PE:C14	2.27	0.47
6:A:721:ACT:H3	8:A:1181:HOH:O	2.16	0.46
1:A:577:GLN:CB	1:A:580[B]:MET:HE3	2.33	0.46
1:A:560:ASP:CG	8:A:1026:HOH:O	2.58	0.46
1:A:181:GLN:HE22	1:A:247:SER:H	1.64	0.46
1:A:301:SER:O	1:A:331:HIS:HE1	1.98	0.45
1:A:167:ARG:NH2	1:A:171:GLN:HE22	2.14	0.45
1:A:265:GLN:H	6:A:720:ACT:H2	1.82	0.45
1:A:374:LYS:HE2	1:A:377:HIS:CE1	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:409:LYS:HB3	5:A:712:1PE:C15	2.49	0.43
1:A:158:ARG:HD3	4:A:704:GOL:H11	2.01	0.43
1:A:169:ILE:O	1:A:173[A]:MET:HG2	2.19	0.42
1:A:482[B]:VAL:HG22	1:A:512:VAL:HG11	2.01	0.42
5:A:713:1PE:OH4	5:A:713:1PE:C22	2.67	0.42
1:A:100:GLN:HA	1:A:100:GLN:NE2	2.31	0.41
5:A:708:1PE:C15	8:A:1208:HOH:O	2.68	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	617/609 (101%)	586 (95%)	21 (3%)	10 (2%)	7 2

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	-2	HIS
1	A	491	GLU
1	A	496	ASN
1	A	584	ASP
1	A	494	GLY
1	A	495	ASP
1	A	505	GLY
1	A	585	ALA
1	A	-1	ALA
1	A	582	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	500/490 (102%)	486 (97%)	14 (3%)	38	21

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	LYS
1	A	100	GLN
1	A	186	LYS
1	A	218	ILE
1	A	226	ASN
1	A	440[A]	ARG
1	A	440[B]	ARG
1	A	482[A]	VAL
1	A	482[B]	VAL
1	A	491	GLU
1	A	495	ASP
1	A	496	ASN
1	A	584	ASP
1	A	592[A]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	GLN
1	A	100	GLN
1	A	101	ASN
1	A	102	ASN
1	A	171	GLN
1	A	181	GLN
1	A	226	ASN
1	A	234	HIS
1	A	262	HIS
1	A	323	GLN
1	A	331	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	333	ASN
1	A	377	HIS
1	A	394	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	701	1	14,14,15	0.62	0	17,19,21	1.64	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	701	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	NAG	C3-C4-C5	3.93	117.36	110.23
3	A	701	NAG	O5-C5-C4	2.99	118.11	110.83
3	A	701	NAG	O5-C1-C2	-2.48	107.46	111.29
3	A	701	NAG	C1-O5-C5	2.36	115.35	112.19

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	701	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	U1Y	B	1	2	13,13,13	0.61	0	17,18,18	0.91	0
2	BGC	B	2	2	11,11,12	0.97	0	15,15,17	1.24	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	U1Y	B	1	2	-	0/4/24/24	0/1/1/1
2	BGC	B	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	BGC	C1-C2-C3	3.35	114.52	109.64


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	ACT	A	719	-	3,3,3	0.86	0	3,3,3	0.63	0
5	1PE	A	710	-	7,7,15	0.48	0	6,6,14	0.26	0
5	1PE	A	709	-	9,9,15	0.40	0	8,8,14	0.49	0
6	ACT	A	720	-	3,3,3	0.79	0	3,3,3	0.57	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	705	-	5,5,5	0.32	0	5,5,5	0.35	0
5	1PE	A	713	-	6,6,15	0.45	0	5,5,14	0.31	0
4	GOL	A	704	-	5,5,5	0.46	0	5,5,5	0.87	0
6	ACT	A	717	-	3,3,3	0.81	0	3,3,3	0.76	0
4	GOL	A	707	-	5,5,5	0.42	0	5,5,5	0.20	0
5	1PE	A	708	-	4,4,15	0.45	0	3,3,14	0.24	0
6	ACT	A	721	-	3,3,3	0.78	0	3,3,3	0.77	0
6	ACT	A	718	-	3,3,3	0.76	0	3,3,3	0.91	0
5	1PE	A	714	-	7,7,15	0.59	0	6,6,14	0.40	0
7	SO4	A	723	-	4,4,4	0.55	0	6,6,6	0.69	0
5	1PE	A	712	-	4,4,15	0.23	0	3,3,14	0.49	0
6	ACT	A	722	-	3,3,3	0.98	0	3,3,3	0.76	0
5	1PE	A	715	-	4,4,15	0.60	0	3,3,14	0.18	0
5	1PE	A	711	-	4,4,15	0.39	0	3,3,14	0.27	0
4	GOL	A	703	-	5,5,5	0.57	0	5,5,5	1.04	0
4	GOL	A	706	-	5,5,5	0.26	0	5,5,5	0.62	0
3	NAG	A	701	1	14,14,15	0.62	0	17,19,21	1.64	4 (23%)
5	1PE	A	716	-	5,5,15	0.57	0	4,4,14	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	1PE	A	711	-	-	1/2/2/13	-
5	1PE	A	712	-	-	0/2/2/13	-
5	1PE	A	714	-	-	3/5/5/13	-
5	1PE	A	715	-	-	0/2/2/13	-
4	GOL	A	707	-	-	4/4/4/4	-
5	1PE	A	710	-	-	5/5/5/13	-
4	GOL	A	703	-	-	2/4/4/4	-
5	1PE	A	708	-	-	2/2/2/13	-
5	1PE	A	709	-	-	2/7/7/13	-
4	GOL	A	706	-	-	0/4/4/4	-
3	NAG	A	701	1	-	1/6/23/26	0/1/1/1
4	GOL	A	705	-	-	0/4/4/4	-
5	1PE	A	713	-	-	2/4/4/13	-
4	GOL	A	704	-	-	2/4/4/4	-
5	1PE	A	716	-	-	1/3/3/13	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	NAG	C3-C4-C5	3.93	117.36	110.23
3	A	701	NAG	O5-C5-C4	2.99	118.11	110.83
3	A	701	NAG	O5-C1-C2	-2.48	107.46	111.29
3	A	701	NAG	C1-O5-C5	2.36	115.35	112.19

There are no chirality outliers.

All (25) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	701	NAG	C3-C2-N2-C7
4	A	703	GOL	C1-C2-C3-O3
4	A	704	GOL	O1-C1-C2-O2
4	A	704	GOL	O1-C1-C2-C3
4	A	707	GOL	C1-C2-C3-O3
4	A	707	GOL	O2-C2-C3-O3
5	A	711	1PE	OH2-C12-C22-OH3
4	A	707	GOL	O1-C1-C2-C3
5	A	708	1PE	OH7-C16-C26-OH6
5	A	713	1PE	OH2-C12-C22-OH3
4	A	703	GOL	O2-C2-C3-O3
4	A	707	GOL	O1-C1-C2-O2
5	A	709	1PE	OH6-C15-C25-OH5
5	A	710	1PE	OH7-C16-C26-OH6
5	A	710	1PE	C15-C25-OH5-C14
5	A	709	1PE	OH7-C16-C26-OH6
5	A	714	1PE	C15-C25-OH5-C14
5	A	714	1PE	OH7-C16-C26-OH6
5	A	713	1PE	C13-C23-OH3-C22
5	A	708	1PE	C16-C26-OH6-C15
5	A	710	1PE	OH6-C15-C25-OH5
5	A	714	1PE	C25-C15-OH6-C26
5	A	716	1PE	C12-C22-OH3-C23
5	A	710	1PE	C25-C15-OH6-C26
5	A	710	1PE	C16-C26-OH6-C15





There are no ring outliers.



7 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	720	ACT	1	0
5	A	713	1PE	2	0
4	A	704	GOL	1	0
5	A	708	1PE	2	0
6	A	721	ACT	1	0
5	A	714	1PE	2	0
5	A	712	1PE	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	592/609 (97%)	-0.31	21 (3%)	47 54	12, 19, 37, 68	13 (2%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	497	LEU	7.5
1	A	582	VAL	7.4
1	A	585	ALA	6.2
1	A	602	THR	4.6
1	A	586	HIS	4.5
1	A	224	ILE	4.3
1	A	583	GLY	4.2
1	A	-1	ALA	4.1
1	A	187	ASP	4.1
1	A	584	ASP	3.9
1	A	225	ILE	3.4
1	A	186	LYS	3.1
1	A	214	THR	3.1
1	A	256	TRP	2.8
1	A	232	ASN	2.8
1	A	228	GLU	2.7
1	A	221	ASN	2.5
1	A	227	ARG	2.2
1	A	257	ASN	2.1
1	A	294	THR	2.1
1	A	576	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	701	14/15	0.67	0.20	54,62,66,71	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	B	2	11/12	0.98	0.04	13,15,16,16	0
2	U1Y	B	1	13/13	0.99	0.04	15,17,21,22	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	A	701	14/15	0.67	0.20	54,62,66,71	0
6	ACT	A	721	4/4	0.67	0.25	65,66,67,67	0
6	ACT	A	717	4/4	0.73	0.23	64,66,67,67	0
5	1PE	A	714	8/16	0.73	0.23	38,53,58,60	0
4	GOL	A	705	6/6	0.74	0.19	60,61,62,63	0
5	1PE	A	709	10/16	0.76	0.28	67,72,73,75	0
6	ACT	A	720	4/4	0.76	0.22	47,50,54,62	0
5	1PE	A	708	5/16	0.76	0.23	54,63,66,67	0
5	1PE	A	715	5/16	0.77	0.21	48,53,55,59	0
6	ACT	A	718	4/4	0.79	0.20	45,46,47,49	0
5	1PE	A	716	6/16	0.80	0.26	46,53,57,59	0
6	ACT	A	722	4/4	0.80	0.18	36,47,52,53	0
4	GOL	A	707	6/6	0.82	0.16	58,62,64,67	0
5	1PE	A	710	8/16	0.85	0.20	55,58,69,71	0
4	GOL	A	703	6/6	0.87	0.16	24,31,33,34	0
6	ACT	A	719	4/4	0.88	0.17	58,58,60,61	0
7	SO4	A	723	5/5	0.88	0.17	27,29,38,40	5
5	1PE	A	713	7/16	0.90	0.13	31,41,52,57	0
5	1PE	A	711	5/16	0.91	0.15	36,46,47,51	0
4	GOL	A	704	6/6	0.91	0.12	23,27,32,33	0
4	GOL	A	706	6/6	0.92	0.14	21,32,35,41	0
5	1PE	A	712	5/16	0.93	0.14	29,30,35,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 1PE A 714:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 1PE A 709:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 1PE A 708:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 1PE A 715:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 1PE A 716:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 1PE A 710:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around 1PE A 713:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 1PE A 711:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.