



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 05:39 PM UTC

PDB ID : 5JHB / pdb_00005jhb
Title : Structure of Phosphoinositide 3-kinase gamma (PI3K) bound to the potent inhibitor PIKin3
Authors : Burke, J.E.; Inglis, A.J.; Williams, R.L.
Deposited on : 2016-04-20
Resolution : 2.48 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	960	5% 52% 32% 16%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6673 atoms, of which 21 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit gamma isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	811	Total	C	N	O	S	0	0	0
			6584	4236	1121	1192	35			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	143	MET	-	initiating methionine	UNP P48736

- Molecule 2 is [1-{4-[6-amino-4-(trifluoromethyl)pyridin-3-yl]-6-(morpholin-4-yl)-1,3,5-triazin-2-yl}-3-(chloromethyl)azetidin-3-yl]methanol (CCD ID: 6K5) (formula: C₁₈H₂₁ClF₃N₇O₂).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total	C	Cl	F	H	N	O	0	0
			52	18	1	3	21	7	2		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	32	Total	O	0	0
			32	32		

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 52% 5% 32% 16%

The sequence logo displays the conservation of amino acids across 250 positions. The y-axis for each position lists the amino acids and their corresponding residue numbers. The x-axis represents the position in the sequence. The logo is color-coded into four regions: green (52%), red (5%), yellow (32%), and grey (16%). Red dots above the bars indicate specific mutations.

Position	Amino Acid	Residue Number
1	Met	1
2	Ala	2
3	Ala	3
4	Leu	4
5	Leu	5
6	Val	6
7	Val	7
8	Thr	8
9	Thr	9
10	Thr	10
11	Thr	11
12	Thr	12
13	Thr	13
14	Thr	14
15	Thr	15
16	Thr	16
17	Thr	17
18	Thr	18
19	Thr	19
20	Thr	20
21	Thr	21
22	Thr	22
23	Thr	23
24	Thr	24
25	Thr	25
26	Thr	26
27	Thr	27
28	Thr	28
29	Thr	29
30	Thr	30
31	Thr	31
32	Thr	32
33	Thr	33
34	Thr	34
35	Thr	35
36	Thr	36
37	Thr	37
38	Thr	38
39	Thr	39
40	Thr	40
41	Thr	41
42	Thr	42
43	Thr	43
44	Thr	44
45	Thr	45
46	Thr	46
47	Thr	47
48	Thr	48
49	Thr	49
50	Thr	50
51	Thr	51
52	Thr	52
53	Thr	53
54	Thr	54
55	Thr	55
56	Thr	56
57	Thr	57
58	Thr	58
59	Thr	59
60	Thr	60
61	Thr	61
62	Thr	62
63	Thr	63
64	Thr	64
65	Thr	65
66	Thr	66
67	Thr	67
68	Thr	68
69	Thr	69
70	Thr	70
71	Thr	71
72	Thr	72
73	Thr	73
74	Thr	74
75	Thr	75
76	Thr	76
77	Thr	77
78	Thr	78
79	Thr	79
80	Thr	80
81	Thr	81
82	Thr	82
83	Thr	83
84	Thr	84
85	Thr	85
86	Thr	86
87	Thr	87
88	Thr	88
89	Thr	89
90	Thr	90
91	Thr	91
92	Thr	92
93	Thr	93
94	Thr	94
95	Thr	95
96	Thr	96
97	Thr	97
98	Thr	98
99	Thr	99
100	Thr	100
101	Thr	101
102	Thr	102
103	Thr	103
104	Thr	104
105	Thr	105
106	Thr	106
107	Thr	107
108	Thr	108
109	Thr	109
110	Thr	110
111	Thr	111
112	Thr	112
113	Thr	113
114	Thr	114
115	Thr	115
116	Thr	116
117	Thr	117
118	Thr	118
119	Thr	119
120	Thr	120
121	Thr	121
122	Thr	122
123	Thr	123
124	Thr	124
125	Thr	125
126	Thr	126
127	Thr	127
128	Thr	128
129	Thr	129
130	Thr	130
131	Thr	131
132	Thr	132
133	Thr	133
134	Thr	134
135	Thr	135
136	Thr	136
137	Thr	137
138	Thr	138
139	Thr	139
140	Thr	140
141	Thr	141
142	Thr	142
143	Thr	143
144	Thr	144
145	Thr	145
146	Thr	146
147	Thr	147
148	Thr	148



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	142.53Å 67.60Å 106.55Å 90.00° 96.23° 90.00°	Depositor
Resolution (Å)	62.08 – 2.48 62.08 – 2.48	Depositor EDS
% Data completeness (in resolution range)	98.0 (62.08-2.48) 98.0 (62.08-2.48)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 2.48Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.231 , 0.277 0.240 , 0.280	Depositor DCC
R_{free} test set	1767 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	70.4	Xtriage
Anisotropy	0.160	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 87.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6673	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6K5, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/6723	0.70	4/9086 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	788	ASP	CA-C-N	6.46	126.15	119.56
1	A	788	ASP	C-N-CA	6.46	126.15	119.56
1	A	988	THR	CA-C-N	5.54	125.37	119.28
1	A	988	THR	C-N-CA	5.54	125.37	119.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6584	0	6625	303	0
2	A	31	21	0	0	0
3	A	5	0	0	0	0
4	A	32	0	0	2	0
All	All	6652	21	6625	303	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 303 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:272:LEU:HB3	1:A:305:VAL:HG11	1.37	1.06
1:A:367:GLY:HA3	1:A:409:LEU:HG	1.39	0.99
1:A:202:VAL:HG21	1:A:285:THR:HG21	1.43	0.99
1:A:1014:VAL:HG11	1:A:1065:LYS:HE3	1.41	0.98
1:A:207:LEU:HD21	1:A:211:LEU:HB3	1.53	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	785/960 (82%)	760 (97%)	24 (3%)	1 (0%)	48 66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1079	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	730/858 (85%)	729 (100%)	1 (0%)	88 95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	211	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	225	HIS
1	A	268	GLN
1	A	389	HIS
1	A	705	GLN
1	A	949	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	1202	-	4,4,4	0.23	0	6,6,6	0.09	0
2	6K5	A	1201	-	33,34,34	2.66	14 (42%)	43,51,51	2.24	14 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6K5	A	1201	-	-	5/21/42/42	0/4/4/4

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	6K5	CAQ-NAR	6.72	1.49	1.35
2	A	1201	6K5	CAV-CBA	5.62	1.59	1.54
2	A	1201	6K5	CAO-NAU	5.28	1.46	1.35
2	A	1201	6K5	CAZ-CBA	4.56	1.58	1.54
2	A	1201	6K5	CAI-CAM	4.46	1.58	1.48

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	6K5	CAO-NAL-CAM	5.02	117.78	114.41
2	A	1201	6K5	NAN-CAQ-NAR	4.65	123.73	117.12
2	A	1201	6K5	CAC-CAD-NAE	-4.63	117.31	121.47
2	A	1201	6K5	CAQ-NAN-CAM	4.39	117.36	114.41
2	A	1201	6K5	CAC-CAH-CAI	4.38	124.97	119.85

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1201	6K5	NAL-CAO-NAU-CAZ
2	A	1201	6K5	NAL-CAO-NAU-CAV
2	A	1201	6K5	NAP-CAO-NAU-CAZ
2	A	1201	6K5	NAP-CAO-NAU-CAV
2	A	1201	6K5	NAN-CAQ-NAR-CAW

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	811/960 (84%)	0.48	45 (5%) 30 27	44, 97, 160, 237	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1044	SER	3.6
1	A	249	PHE	3.5
1	A	365	ILE	3.4
1	A	902	PHE	3.3
1	A	892	GLN	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	1202	5/5	0.79	0.11	120,120,128,141	0
2	6K5	A	1201	31/31	0.91	0.10	60,79,129,139	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.