



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 05:55 PM UTC

PDB ID : 2JIW / pdb_00002jiw
Title : Bacteroides thetaiotaomicron GH84 O-GlcNAcase in complex with 2- Acetylamino-2-deoxy-1-epivalienamine
Authors : Dennis, R.J.; Davies, G.J.
Deposited on : 2007-07-02
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	716	
1	B	716	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 10470 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called O-GLCNACASE BT_4395.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	590	Total	C	N	O	S	0	4	0
			4821	3087	812	904	18			
1	B	590	Total	C	N	O	S	0	9	0
			4853	3109	820	905	19			

- Molecule 2 is N-[(1S,2R,5R,6R)-2-AMINO-5,6-DIHYDROXY-4-(HYDROXYMETHYL)CYCLOHEX-3-EN-1-YL]ACETAMIDE (CCD ID: BEU) (formula: C₉H₁₆N₂O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			15	9	2	4		
2	B	1	Total	C	N	O	0	0
			15	9	2	4		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	363	Total 363	O 363	0	0
3	B	403	Total 403	O 403	0	0

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

[illegible]

Chain B:

6% 71% 10% 18%

MET ASN VAL S4 L5 Q6 Q15 M16 K17 T18 I19 S20 L21 P22 A23 V24 Y25 Q26 L27 N28 L45 S46 G47 K48 Q49 S50 S51 K52 A53 G54 M55 L56 G63 Y81 Y82 L83 E87 K112 L116 E120 I121 D123 A173 P174 L188 R219 G220 L221 M229 S236 F237 A238 L257 N273 V276 W286 M308 W309 T310 V314 I315 R320 S324 Y335 I336 W337 V355 Y356 T371 T399 W400 E416 C420 W423 M441 L451 K452 A453 F454 K455 E456 G457 K458 W459

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	51.40Å 94.13Å 99.81Å 104.97° 94.09° 102.48°	Depositor
Resolution (Å)	95.35 – 1.95 95.35 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.0 (95.35-1.95) 97.0 (95.35-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.206 , 0.248 0.207 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	22.1	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10470	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BEU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	2/4961 (0.0%)	0.91	1/6726 (0.0%)
1	B	0.77	0/5008	0.91	4/6786 (0.1%)
All	All	0.77	2/9969 (0.0%)	0.91	5/13512 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	496	THR	CA-CB	5.97	1.59	1.54
1	A	273	ASN	CA-C	5.71	1.55	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	116	LEU	CA-C-N	-6.50	113.94	120.31
1	B	116	LEU	C-N-CA	-6.50	113.94	120.31
1	B	291	GLY	N-CA-C	-6.17	104.31	112.82
1	B	399	THR	CB-CA-C	-5.86	101.68	110.88
1	A	89	GLU	N-CA-C	5.04	115.97	108.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	22	PRO	Peptide
1	B	22	PRO	Peptide
1	B	4	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4821	0	4721	31	0
1	B	4853	0	4775	45	0
2	A	15	0	16	0	0
2	B	15	0	16	0	0
3	A	363	0	0	4	0
3	B	403	0	0	1	0
All	All	10470	0	9528	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:458:LYS:HB3	1:B:459:ASN:HA	1.25	1.12
1:B:22:PRO:HA	1:B:23:ALA:HB3	1.35	1.08
1:B:22:PRO:HA	1:B:23:ALA:CB	1.85	1.06
1:B:219[A]:ARG:HG3	1:B:219[A]:ARG:HH11	1.32	0.92
1:A:22:PRO:HA	1:A:23:ALA:CB	2.03	0.89
1:A:22:PRO:HA	1:A:23:ALA:HB3	1.55	0.89
1:B:458:LYS:HB3	1:B:459:ASN:CA	2.12	0.77
1:B:22:PRO:CA	1:B:23:ALA:HB3	2.17	0.73
1:B:219[A]:ARG:HG3	1:B:219[A]:ARG:NH1	2.03	0.72
1:B:453:ALA:O	1:B:458:LYS:N	2.26	0.63
1:A:19:ILE:HD13	1:A:55:MET:HE1	1.83	0.60
1:A:532:LYS:O	1:A:536:GLN:HG3	1.99	0.60
1:B:355:VAL:O	1:B:399:THR:HG23	2.01	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:LYS:HG2	3:A:2026:HOH:O	2.01	0.60
1:B:536:GLN:HG2	1:B:590:TYR:CD1	2.38	0.58
1:B:458:LYS:CB	1:B:459:ASN:HA	2.14	0.57
1:A:22:PRO:CA	1:A:23:ALA:HB3	2.33	0.56
1:A:70:TYR:OH	1:A:89:GLU:OE1	2.22	0.56
1:B:22:PRO:HA	1:B:23:ALA:HB2	1.82	0.56
1:B:462:LYS:HD3	1:B:466:GLU:OE2	2.06	0.56
1:A:262:ASP:HA	1:A:266:ALA:HB3	1.88	0.55
1:B:21:LEU:HD23	1:B:48:LYS:HE3	1.89	0.55
1:A:238:ALA:HA	1:A:276:VAL:O	2.07	0.55
1:A:585:ASP:OD1	1:A:587:THR:HG23	2.08	0.54
1:B:19:ILE:HD13	1:B:55:MET:HE1	1.89	0.53
1:B:219[A]:ARG:HH12	1:B:257:LEU:HA	1.73	0.53
1:B:28:ASN:HD22	1:B:56:LEU:HD11	1.75	0.51
1:A:22:PRO:CA	1:A:23:ALA:CB	2.85	0.51
1:A:81:TYR:CZ	1:A:123:ASP:HB3	2.46	0.50
1:A:22:PRO:HA	1:A:23:ALA:HB2	1.93	0.49
1:A:455:LYS:HB3	3:A:2302:HOH:O	2.13	0.49
1:A:293:TYR:CD2	1:A:293:TYR:C	2.91	0.49
1:B:549:PRO:HG2	1:B:550:TYR:CE2	2.48	0.49
1:A:341:PRO:O	1:A:342:VAL:C	2.56	0.49
1:A:70:TYR:O	1:A:73:GLN:HG2	2.14	0.48
1:B:451:LEU:HD13	1:B:564:LEU:HD12	1.96	0.48
1:B:441[A]:MET:HE3	1:B:441[A]:MET:H	1.79	0.47
1:A:451:LEU:HD13	1:A:564:LEU:HD12	1.97	0.47
1:A:188:LEU:HD13	1:A:229:MET:HE1	1.98	0.46
1:B:534:LEU:O	1:B:538:MET:HG3	2.16	0.46
1:B:356:TYR:HB3	1:B:399:THR:HG21	1.97	0.46
1:B:308:MET:HA	1:B:335:TYR:O	2.16	0.45
1:B:83:LEU:HD22	1:B:121:ILE:HD12	1.98	0.45
1:B:173:ALA:HA	1:B:174:PRO:HA	1.67	0.45
1:A:228[A]:LYS:HA	1:A:228[A]:LYS:HD2	1.83	0.45
1:B:585:ASP:OD1	1:B:587:THR:HG23	2.17	0.45
1:A:423:MET:HE2	3:A:2282:HOH:O	2.17	0.44
1:B:238:ALA:HA	1:B:276:VAL:O	2.18	0.44
1:B:420:CYS:HA	1:B:423[A]:MET:HE3	1.99	0.44
1:B:310:THR:HA	1:B:337:TRP:O	2.17	0.44
1:B:540[A]:TYR:CE1	1:B:544:THR:HG21	2.53	0.43
1:A:261:ILE:O	1:A:265:PHE:HB3	2.18	0.43
1:B:236:SER:OG	1:B:273:ASN:HB2	2.19	0.43
1:A:536:GLN:HG2	1:A:590:TYR:CD1	2.53	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:MET:HA	1:A:335:TYR:O	2.18	0.43
1:B:81:TYR:CZ	1:B:123:ASP:HB3	2.52	0.43
1:B:475:MET:HG2	1:B:502:PHE:CZ	2.53	0.43
1:B:188:LEU:HD13	1:B:229:MET:HE1	1.99	0.43
1:B:516:VAL:HG22	1:B:572:VAL:HG11	2.02	0.42
1:A:378:GLU:HG3	1:A:490:PRO:HB2	2.01	0.42
1:B:320:ARG:O	1:B:324:SER:OG	2.32	0.42
1:B:173:ALA:HB3	3:B:2049:HOH:O	2.19	0.42
1:A:303:PRO:HD2	3:A:2208:HOH:O	2.20	0.42
1:A:557:ALA:HB1	1:A:561:ILE:HB	2.01	0.42
1:B:314:VAL:HG12	1:B:315:ILE:HD13	2.02	0.41
1:B:302:ASN:HA	1:B:303:PRO:HD3	1.93	0.41
1:B:83:LEU:HD23	1:B:83:LEU:C	2.46	0.41
1:B:219[A]:ARG:NH1	1:B:257:LEU:HD12	2.35	0.41
1:B:423[B]:MET:HB2	1:B:423[B]:MET:HE2	1.30	0.41
1:A:73:GLN:HG3	1:A:82:TYR:CE2	2.56	0.40
1:B:531:VAL:HG11	1:B:569:PHE:CE1	2.56	0.40
1:A:534:LEU:O	1:A:538:MET:HG3	2.21	0.40
1:A:562:LYS:HB3	1:A:563:PRO:HD3	2.04	0.40
1:B:454:PHE:HZ	1:B:516:VAL:HG13	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	592/716 (83%)	570 (96%)	18 (3%)	4 (1%)	18	10
1	B	597/716 (83%)	577 (97%)	18 (3%)	2 (0%)	36	28
All	All	1189/1432 (83%)	1147 (96%)	36 (3%)	6 (0%)	24	16

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	458	LYS
1	A	23	ALA
1	A	518	GLY
1	B	23	ALA
1	A	49	GLN
1	B	49	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	522/630 (83%)	502 (96%)	20 (4%)	29	19
1	B	527/630 (84%)	504 (96%)	23 (4%)	25	15
All	All	1049/1260 (83%)	1006 (96%)	43 (4%)	28	17

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	SER
1	A	27	LEU
1	A	32	GLU
1	A	43	GLU
1	A	45	LEU
1	A	46	SER
1	A	48	LYS
1	A	53	LYS
1	A	73	GLN
1	A	88	LYS
1	A	178	LEU
1	A	219	ARG
1	A	221	LEU
1	A	371	THR
1	A	452	LYS
1	A	455	LYS
1	A	456	GLU
1	A	459	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	462	LYS
1	A	564	LEU
1	B	6	GLN
1	B	19	ILE
1	B	24	VAL
1	B	26	GLN
1	B	27	LEU
1	B	45	LEU
1	B	49	GLN
1	B	52	LYS
1	B	87	GLU
1	B	120	GLU
1	B	219[A]	ARG
1	B	219[B]	ARG
1	B	221	LEU
1	B	324	SER
1	B	371	THR
1	B	416	GLU
1	B	441[A]	MET
1	B	441[B]	MET
1	B	452	LYS
1	B	455	LYS
1	B	462	LYS
1	B	496	THR
1	B	519	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	156	ASN
1	A	189	GLN
1	A	267	GLN
1	A	274	GLN
1	A	459	ASN
1	A	469	GLN
1	A	593	HIS
1	B	6	GLN
1	B	11	GLN
1	B	28	ASN
1	B	49	GLN
1	B	73	GLN
1	B	252	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	254	GLN
1	B	274	GLN
1	B	306	GLN
1	B	444	GLN
1	B	543	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BEU	A	1594	-	15,15,15	0.75	0	13,21,21	1.08	2 (15%)
2	BEU	B	1594	-	15,15,15	0.99	1 (6%)	13,21,21	1.40	3 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEU	A	1594	-	-	2/6/26/26	0/1/1/1
2	BEU	B	1594	-	-	2/6/26/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1594	BEU	C3-C4	3.36	1.57	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1594	BEU	C5-C6-C1	-2.78	107.29	111.67
2	A	1594	BEU	C5-C6-C1	-2.53	107.69	111.67
2	B	1594	BEU	C4-C3-N3	2.31	115.87	111.71
2	A	1594	BEU	C6-C1-C2	-2.25	118.53	122.23
2	B	1594	BEU	C5-C4-N4	-2.03	106.88	110.62

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1594	BEU	C3-C4-N4-C4'
2	B	1594	BEU	C3-C4-N4-C4'
2	A	1594	BEU	C5-C4-N4-C4'
2	B	1594	BEU	C5-C4-N4-C4'

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	590/716 (82%)	0.21	41 (6%)	23 26	10, 20, 47, 64	4 (0%)
1	B	590/716 (82%)	0.19	42 (7%)	22 25	11, 19, 45, 62	9 (1%)
All	All	1180/1432 (82%)	0.20	83 (7%)	22 26	10, 20, 47, 64	13 (1%)

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	23	ALA	5.1
1	B	453	ALA	4.5
1	B	457	GLY	4.4
1	B	16	ASN	4.4
1	A	461	ASP	4.2
1	A	50	SER	4.2
1	A	453	ALA	4.1
1	B	23	ALA	4.0
1	A	26	GLN	3.9
1	B	451	LEU	3.9
1	A	54	GLY	3.9
1	B	24	VAL	3.9
1	B	54	GLY	3.8
1	B	454	PHE	3.7
1	B	455	LYS	3.6
1	B	458	LYS	3.6
1	A	18	THR	3.4
1	B	583	HIS	3.4
1	B	47	GLY	3.3
1	B	45	LEU	3.3
1	A	51	SER	3.3
1	A	24	VAL	3.3
1	B	50	SER	3.3
1	A	458	LYS	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	457	GLY	3.2
1	A	454	PHE	3.1
1	A	49	GLN	3.1
1	B	52	LYS	3.1
1	B	22	PRO	3.1
1	A	451	LEU	3.0
1	B	461	ASP	3.0
1	A	455	LYS	2.9
1	A	456	GLU	2.9
1	B	53	LYS	2.9
1	A	16	ASN	2.8
1	B	18	THR	2.8
1	B	452	LYS	2.8
1	B	286	TRP	2.8
1	A	47	GLY	2.8
1	B	49	GLN	2.7
1	B	63	GLY	2.7
1	A	27	LEU	2.7
1	B	48	LYS	2.7
1	A	544	THR	2.7
1	A	19	ILE	2.7
1	B	19	ILE	2.7
1	A	113	ASP	2.6
1	A	583	HIS	2.6
1	A	45	LEU	2.6
1	B	87	GLU	2.6
1	A	463	ALA	2.6
1	A	52	LYS	2.6
1	A	87	GLU	2.5
1	B	581	ASN	2.5
1	A	4	SER	2.5
1	B	518	GLY	2.4
1	B	17	LYS	2.4
1	A	28	ASN	2.4
1	A	116	LEU	2.4
1	B	112	LYS	2.4
1	B	56	LEU	2.3
1	A	22	PRO	2.3
1	A	46	SER	2.3
1	B	456	GLU	2.3
1	A	593	HIS	2.3
1	A	14	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	15	GLN	2.3
1	B	584	LEU	2.3
1	A	540[A]	TYR	2.2
1	B	25	TYR	2.2
1	A	17	LYS	2.2
1	B	46	SER	2.2
1	A	25	TYR	2.1
1	B	400	TRP	2.1
1	B	459	ASN	2.1
1	A	15	GLN	2.1
1	A	580	PHE	2.1
1	B	26	GLN	2.1
1	A	57	ILE	2.1
1	B	540[A]	TYR	2.1
1	B	21	LEU	2.0
1	B	582	ALA	2.0
1	A	459	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BEU	A	1594	15/15	0.96	0.06	13,16,18,19	0
2	BEU	B	1594	15/15	0.96	0.05	11,14,15,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.