



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 08:58 PM UTC

PDB ID : 5JIA / pdb_00005jia
Title : The Crystal Structure Of IUS-SPRY Domain From RanBP10
Authors : Hong, S.K.; Kim, K.-H.; Kim, E.E.
Deposited on : 2016-04-22
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	80% 16% ..
1	B	210	4% 86% 11% ..
1	C	210	3% 84% 12% ..
1	D	210	2% 85% 13% .
1	E	210	3% 86% 12% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	210	
1	G	210	
1	H	210	
1	I	210	
1	J	210	
1	K	210	
1	L	210	
1	M	210	
1	N	210	
1	O	210	
1	P	210	

2 Entry composition


There are 3 unique types of molecules in this entry. The entry contains 27983 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ran-binding protein 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	206	Total	C	N	O	S	0	0	0
			1614	1021	288	297	8			
1	B	207	Total	C	N	O	S	0	0	0
			1623	1026	289	300	8			
1	C	205	Total	C	N	O	S	0	0	0
			1602	1012	285	297	8			
1	D	210	Total	C	N	O	S	0	0	0
			1640	1036	292	304	8			
1	E	207	Total	C	N	O	S	0	0	0
			1618	1023	288	299	8			
1	F	207	Total	C	N	O	S	0	0	0
			1615	1020	287	300	8			
1	G	203	Total	C	N	O	S	0	0	0
			1586	1003	280	295	8			
1	H	203	Total	C	N	O	S	0	0	0
			1589	1005	283	293	8			
1	I	206	Total	C	N	O	S	0	0	0
			1614	1021	287	298	8			
1	J	199	Total	C	N	O	S	0	0	0
			1550	980	275	287	8			
1	K	203	Total	C	N	O	S	0	0	0
			1586	1003	280	295	8			
1	L	205	Total	C	N	O	S	0	0	0
			1607	1016	286	297	8			
1	M	208	Total	C	N	O	S	0	0	0
			1624	1026	289	301	8			
1	N	203	Total	C	N	O	S	0	0	0
			1586	1003	280	295	8			
1	O	202	Total	C	N	O	S	0	0	0
			1572	992	278	294	8			
1	P	204	Total	C	N	O	S	0	0	0
			1596	1011	284	293	8			

- Molecule 2 is IMIDAZOLE (CCD ID: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			5	3	2		
2	D	1	Total	C	N	0	0
			5	3	2		
2	E	1	Total	C	N	0	0
			5	3	2		
2	G	1	Total	C	N	0	0
			5	3	2		
2	H	1	Total	C	N	0	0
			5	3	2		
2	J	1	Total	C	N	0	0
			5	3	2		
2	N	1	Total	C	N	0	0
			5	3	2		
2	P	1	Total	C	N	0	0
			5	3	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	169	Total	O	0	0
			169	169		
3	B	130	Total	O	0	0
			130	130		
3	C	141	Total	O	0	0
			141	141		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	137	Total 137	O 137	0	0
3	E	173	Total 173	O 173	0	0
3	F	148	Total 148	O 148	0	0
3	G	153	Total 153	O 153	0	0
3	H	161	Total 161	O 161	0	0
3	I	199	Total 199	O 199	0	0
3	J	155	Total 155	O 155	0	0
3	K	168	Total 168	O 168	0	0
3	L	94	Total 94	O 94	0	0
3	M	165	Total 165	O 165	0	0
3	N	148	Total 148	O 148	0	0
3	O	101	Total 101	O 101	0	0
3	P	79	Total 79	O 79	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ran-binding protein 10




- Molecule 1: Ran-binding protein 10




- Molecule 1: Ran-binding protein 10



- Molecule 1: Ran-binding protein 10




- Molecule 1: Ran-binding protein 10








- Molecule 1: Ran-binding protein 10





- Molecule 1: Ran-binding protein 10




- Molecule 1: Ran-binding protein 10




- Molecule 1: Ran-binding protein 10




- Molecule 1: Ran-binding protein 10




- Molecule 1: Ran-binding protein 10





- Molecule 1: Ran-binding protein 10





- Molecule 1: Ran-binding protein 10





- Molecule 1: Ran-binding protein 10



- Molecule 1: Ran-binding protein 10



- Molecule 1: Ran-binding protein 10



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	146.34Å 79.24Å 157.63Å 90.00° 97.91° 90.00°	Depositor
Resolution (Å)	41.25 – 1.80 41.25 – 1.80	Depositor EDS
% Data completeness (in resolution range)	92.6 (41.25-1.80) 92.6 (41.25-1.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.184 , 0.227 0.188 , 0.189	Depositor DCC
R_{free} test set	15429 reflections (4.67%)	wwPDB-VP
Wilson B-factor (Å ²)	14.3	Xtriage
Anisotropy	0.408	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	27983	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 44.32 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5519e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IMD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1662	0.81	2/2255 (0.1%)
1	B	0.47	0/1671	0.82	3/2267 (0.1%)
1	C	0.61	4/1650 (0.2%)	0.80	0/2239
1	D	0.51	0/1689	0.91	3/2292 (0.1%)
1	E	0.47	0/1666	0.82	0/2260
1	F	0.51	0/1664	0.85	2/2259 (0.1%)
1	G	0.45	1/1634 (0.1%)	0.80	0/2218
1	H	0.47	0/1637	0.78	0/2222
1	I	0.53	1/1662 (0.1%)	0.83	1/2255 (0.0%)
1	J	0.52	1/1596 (0.1%)	0.84	3/2166 (0.1%)
1	K	0.51	1/1634 (0.1%)	0.78	0/2218
1	L	0.40	0/1655	0.75	0/2245
1	M	0.53	0/1673	0.81	0/2270
1	N	0.45	0/1634	0.78	0/2218
1	O	0.44	0/1618	0.83	3/2195 (0.1%)
1	P	0.46	0/1644	0.90	3/2231 (0.1%)
All	All	0.49	8/26389 (0.0%)	0.82	20/35810 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	41	ARG	C-O	-7.10	1.15	1.24
1	C	38	ARG	C-O	-6.21	1.16	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	41	ARG	CZ-NH2	-6.11	1.25	1.33
1	C	37	ARG	C-O	-6.01	1.17	1.24
1	J	41	ARG	C-O	-5.48	1.17	1.24

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	63	TYR	CA-C-N	8.49	132.51	120.28
1	F	63	TYR	C-N-CA	8.49	132.51	120.28
1	J	34	GLU	N-CA-C	-8.17	102.20	112.90
1	B	59	PRO	CA-C-N	7.26	132.64	120.88
1	B	59	PRO	C-N-CA	7.26	132.64	120.88

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	64	ASN	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1614	0	1540	33	0
1	B	1623	0	1546	18	0
1	C	1602	0	1519	20	0
1	D	1640	0	1561	30	0
1	E	1618	0	1541	21	0
1	F	1615	0	1531	17	0
1	G	1586	0	1501	16	0
1	H	1589	0	1510	14	0
1	I	1614	0	1538	36	0
1	J	1550	0	1476	22	0
1	K	1586	0	1501	15	0
1	L	1607	0	1529	9	0
1	M	1624	0	1544	25	0
1	N	1586	0	1501	18	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	O	1572	0	1491	53	0
1	P	1596	0	1524	30	0
2	A	5	0	5	2	0
2	D	5	0	5	0	0
2	E	5	0	5	0	0
2	G	5	0	5	0	0
2	H	5	0	5	3	0
2	J	5	0	5	1	0
2	N	5	0	5	0	0
2	P	5	0	5	1	0
3	A	169	0	0	1	0
3	B	130	0	0	2	0
3	C	141	0	0	1	0
3	D	137	0	0	2	0
3	E	173	0	0	1	0
3	F	148	0	0	0	0
3	G	153	0	0	1	0
3	H	161	0	0	1	0
3	I	199	0	0	4	0
3	J	155	0	0	2	0
3	K	168	0	0	2	0
3	L	94	0	0	1	0
3	M	165	0	0	1	0
3	N	148	0	0	1	0
3	O	101	0	0	2	0
3	P	79	0	0	2	0
All	All	27983	0	24393	368	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 368 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:231:ARG:O	1:O:232:GLU:HG2	1.41	1.17
1:O:231:ARG:O	1:O:232:GLU:CG	1.95	1.14
1:I:90:VAL:HA	1:I:129:MET:CE	1.78	1.12
1:O:40:GLN:HG2	1:O:41:ARG:NH1	1.65	1.12
1:N:137:LYS:HZ3	1:N:138:HIS:CE1	1.67	1.11

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	204/210 (97%)	200 (98%)	4 (2%)	0	100	100
1	B	205/210 (98%)	197 (96%)	8 (4%)	0	100	100
1	C	203/210 (97%)	199 (98%)	4 (2%)	0	100	100
1	D	208/210 (99%)	204 (98%)	4 (2%)	0	100	100
1	E	205/210 (98%)	202 (98%)	3 (2%)	0	100	100
1	F	205/210 (98%)	198 (97%)	7 (3%)	0	100	100
1	G	201/210 (96%)	197 (98%)	4 (2%)	0	100	100
1	H	201/210 (96%)	197 (98%)	4 (2%)	0	100	100
1	I	204/210 (97%)	200 (98%)	4 (2%)	0	100	100
1	J	197/210 (94%)	192 (98%)	5 (2%)	0	100	100
1	K	201/210 (96%)	196 (98%)	5 (2%)	0	100	100
1	L	203/210 (97%)	197 (97%)	6 (3%)	0	100	100
1	M	206/210 (98%)	200 (97%)	5 (2%)	1 (0%)	24	14
1	N	201/210 (96%)	197 (98%)	4 (2%)	0	100	100
1	O	200/210 (95%)	193 (96%)	7 (4%)	0	100	100
1	P	202/210 (96%)	195 (96%)	7 (4%)	0	100	100
All	All	3246/3360 (97%)	3164 (98%)	81 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	M	30	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/173 (98%)	166 (98%)	4 (2%)	43	31
1	B	171/173 (99%)	165 (96%)	6 (4%)	32	19
1	C	168/173 (97%)	162 (96%)	6 (4%)	31	18
1	D	173/173 (100%)	169 (98%)	4 (2%)	44	33
1	E	170/173 (98%)	169 (99%)	1 (1%)	78	77
1	F	170/173 (98%)	166 (98%)	4 (2%)	43	31
1	G	167/173 (96%)	166 (99%)	1 (1%)	78	77
1	H	167/173 (96%)	165 (99%)	2 (1%)	63	57
1	I	170/173 (98%)	166 (98%)	4 (2%)	43	31
1	J	164/173 (95%)	157 (96%)	7 (4%)	26	13
1	K	167/173 (96%)	166 (99%)	1 (1%)	78	77
1	L	169/173 (98%)	166 (98%)	3 (2%)	51	43
1	M	171/173 (99%)	169 (99%)	2 (1%)	63	57
1	N	167/173 (96%)	165 (99%)	2 (1%)	63	57
1	O	166/173 (96%)	163 (98%)	3 (2%)	51	43
1	P	168/173 (97%)	165 (98%)	3 (2%)	51	43
All	All	2698/2768 (98%)	2645 (98%)	53 (2%)	48	38

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	32	GLU
1	J	60	LYS
1	O	232	GLU
1	I	53	LEU
1	J	33	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	138	HIS
1	M	207	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	J	148	HIS
1	M	48	GLN
1	N	187	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	IMD	A	901	-	5,5,5	0.85	0	5,5,5	0.31	0
2	IMD	P	301	-	5,5,5	0.70	0	5,5,5	0.38	0
2	IMD	H	301	-	5,5,5	0.79	0	5,5,5	0.33	0
2	IMD	G	301	-	5,5,5	0.62	0	5,5,5	0.66	0
2	IMD	N	301	-	5,5,5	0.91	0	5,5,5	0.19	0
2	IMD	J	301	-	5,5,5	0.86	0	5,5,5	0.31	0
2	IMD	E	301	-	5,5,5	0.80	0	5,5,5	0.31	0
2	IMD	D	301	-	5,5,5	0.77	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IMD	A	901	-	-	-	0/1/1/1
2	IMD	P	301	-	-	-	0/1/1/1
2	IMD	H	301	-	-	-	0/1/1/1
2	IMD	G	301	-	-	-	0/1/1/1
2	IMD	N	301	-	-	-	0/1/1/1
2	IMD	J	301	-	-	-	0/1/1/1
2	IMD	E	301	-	-	-	0/1/1/1
2	IMD	D	301	-	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	IMD	2	0
2	P	301	IMD	1	0
2	H	301	IMD	3	0
2	J	301	IMD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	206/210 (98%)	-0.46	2 (0%) 79 80	8, 13, 30, 51	0
1	B	207/210 (98%)	-0.16	9 (4%) 40 39	9, 16, 39, 90	0
1	C	205/210 (97%)	-0.19	7 (3%) 48 48	8, 15, 38, 68	0
1	D	210/210 (100%)	-0.22	5 (2%) 59 60	9, 16, 34, 61	0
1	E	207/210 (98%)	-0.31	6 (2%) 53 53	8, 14, 29, 62	0
1	F	207/210 (98%)	-0.13	5 (2%) 59 60	8, 16, 35, 62	0
1	G	203/210 (96%)	-0.30	4 (1%) 65 65	7, 15, 29, 53	0
1	H	203/210 (96%)	-0.30	5 (2%) 58 58	8, 15, 32, 66	0
1	I	206/210 (98%)	-0.47	2 (0%) 79 80	5, 12, 25, 52	0
1	J	199/210 (94%)	-0.23	5 (2%) 58 58	6, 14, 35, 68	0
1	K	203/210 (96%)	-0.34	6 (2%) 52 52	7, 14, 30, 61	0
1	L	205/210 (97%)	0.09	4 (1%) 65 65	14, 21, 37, 60	0
1	M	208/210 (99%)	-0.29	7 (3%) 48 48	7, 13, 31, 71	0
1	N	203/210 (96%)	-0.25	3 (1%) 72 72	7, 15, 34, 61	0
1	O	202/210 (96%)	0.11	12 (5%) 28 27	11, 19, 43, 76	0
1	P	204/210 (97%)	0.74	34 (16%) 4 3	11, 25, 46, 72	0
All	All	3278/3360 (97%)	-0.17	116 (3%) 47 47	5, 16, 37, 90	0

The worst 5 of 116 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	235	ALA	6.9
1	P	52	PRO	6.2
1	C	31	GLY	5.9
1	P	51	THR	5.8
1	P	126	GLY	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	IMD	P	301	5/5	0.85	0.14	27,29,34,39	0
2	IMD	D	301	5/5	0.89	0.11	14,18,20,23	0
2	IMD	J	301	5/5	0.90	0.11	11,12,22,23	0
2	IMD	A	901	5/5	0.90	0.09	9,15,15,17	0
2	IMD	N	301	5/5	0.91	0.11	8,13,18,19	0
2	IMD	G	301	5/5	0.91	0.10	15,16,20,26	0
2	IMD	H	301	5/5	0.92	0.11	13,13,17,20	0
2	IMD	E	301	5/5	0.94	0.09	11,13,15,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.