



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 01:47 PM UTC

PDB ID : 5JIW / pdb_00005jiw
Title : Crystal structure of Thermus aquaticus amylomaltase (GH77) in complex with a 34-meric cycloamylose
Authors : Roth, C.; Bexten, N.; Weizenmann, N.; Saenger, T.; Maier, T.; Zimmermann, W.; Straeter, N.
Deposited on : 2016-04-22
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	500	12% 96% .
2	B	17	47% 47% 6%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8926 atoms, of which 4173 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-alpha-glucanotransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	500	Total	C	H	N	O	S	0	8	0
			8119	2676	4008	718	705	12			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	293	ALA	ASP	engineered mutation	UNP O87172
A	370	SNN	ASP	conflict	UNP O87172
A	395	ASN	ASP	engineered mutation	UNP O87172

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	17	Total	C	H	O	0	0	0
			340	102	153	85			

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 4 is CARBONATE ION (CCD ID: CO3) (formula: CO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	443	Total 443	O 443	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-alpha-glucanotransferase



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	157.60Å 157.60Å 112.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.15 – 1.73 37.15 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.4 (37.15-1.73) 99.3 (37.15-1.73)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 1.73Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.164 , 0.188 0.167 , 0.190	Depositor DCC
R_{free} test set	1946 reflections (2.63%)	wwPDB-VP
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.209	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 62.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8926	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, GLC, CO3, SNN, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.22	0/4267	0.41	0/5797

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4111	4008	4002	12	0
2	B	187	153	154	1	0
3	A	8	12	12	1	0
4	A	4	0	0	0	0
5	A	443	0	0	8	3
All	All	4753	4173	4168	14	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291[B]:ARG:NH1	5:A:701:HOH:O	2.08	0.85
1:A:249:ASP:O	1:A:251:PHE:N	2.22	0.73
5:A:702:HOH:O	2:B:5:GLC:O3	2.10	0.70
1:A:91:PRO:O	5:A:703:HOH:O	2.10	0.68
1:A:395:ASN:OD1	5:A:704:HOH:O	2.13	0.67
1:A:464:ASN:ND2	5:A:704:HOH:O	2.32	0.62
1:A:270:ARG:NH2	5:A:713:HOH:O	2.38	0.47
1:A:76:PRO:O	1:A:80:ARG:NH1	2.45	0.46
1:A:338[B]:LEU:HD21	5:A:701:HOH:O	2.15	0.45
1:A:119:ALA:O	1:A:120:SER:CB	2.65	0.45
1:A:156:TRP:HA	1:A:159:TRP:CD2	2.54	0.43
1:A:291[B]:ARG:NH1	1:A:361[B]:MET:SD	2.93	0.41
1:A:119:ALA:O	1:A:120:SER:OG	2.34	0.41
3:A:618:EDO:O2	5:A:705:HOH:O	2.22	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:751:HOH:O	5:A:760:HOH:O[3_555]	2.10	0.10
5:A:988:HOH:O	5:A:1059:HOH:O[16_444]	2.15	0.05
5:A:1085:HOH:O	5:A:1136:HOH:O[7_555]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	504/500 (101%)	493 (98%)	7 (1%)	4 (1%)	16 4

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	250	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	2	GLU
1	A	249	ASP
1	A	120	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	407/399 (102%)	404 (99%)	3 (1%)	76	67

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	158	ARG
1	A	175	LYS
1	A	296	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	464	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SNN	A	370	1	7,8,8	2.58	3 (42%)	10,11,11	2.55	4 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SNN	A	370	1	-	-	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	370	SNN	C-N1	4.37	1.43	1.37
1	A	370	SNN	C5-N1	3.73	1.43	1.37
1	A	370	SNN	C4-CA	-3.51	1.49	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	370	SNN	C5-N1-C	-5.67	108.26	113.96
1	A	370	SNN	CA-C-N1	3.08	109.72	107.30
1	A	370	SNN	O5-C5-C4	-2.71	123.03	126.48
1	A	370	SNN	C4-CA-C	2.67	107.30	104.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

17 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	B	1	2	11,11,12	0.64	0	15,15,17	0.60	0
2	GLC	B	10	2	11,11,12	0.55	0	15,15,17	0.82	1 (6%)
2	GLC	B	11	2	11,11,12	0.62	0	15,15,17	0.83	0
2	GLC	B	12	2	11,11,12	0.70	0	15,15,17	0.81	0
2	GLC	B	13	2	11,11,12	0.58	0	15,15,17	0.96	1 (6%)
2	GLC	B	14	2	11,11,12	0.49	0	15,15,17	1.17	1 (6%)
2	GLC	B	15	2	11,11,12	0.58	0	15,15,17	0.88	0
2	GLC	B	16	2	11,11,12	0.60	0	15,15,17	0.73	0
2	GLC	B	17	2	11,11,12	0.54	0	15,15,17	0.77	1 (6%)
2	GLC	B	2	2	11,11,12	0.64	0	15,15,17	0.40	0
2	GLC	B	3	2	11,11,12	0.66	0	15,15,17	0.93	1 (6%)
2	GLC	B	4	2	11,11,12	0.54	0	15,15,17	0.96	1 (6%)
2	GLC	B	5	2	11,11,12	0.59	0	15,15,17	0.90	1 (6%)
2	GLC	B	6	2	11,11,12	0.57	0	15,15,17	1.06	1 (6%)
2	GLC	B	7	2	11,11,12	0.67	0	15,15,17	0.96	1 (6%)
2	GLC	B	8	2	11,11,12	0.67	0	15,15,17	0.76	0
2	GLC	B	9	2	11,11,12	0.65	0	15,15,17	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	1	2	-	0/2/19/22	0/1/1/1
2	GLC	B	10	2	-	0/2/19/22	0/1/1/1
2	GLC	B	11	2	-	1/2/19/22	0/1/1/1
2	GLC	B	12	2	-	1/2/19/22	0/1/1/1
2	GLC	B	13	2	-	0/2/19/22	0/1/1/1
2	GLC	B	14	2	-	0/2/19/22	0/1/1/1
2	GLC	B	15	2	-	1/2/19/22	0/1/1/1
2	GLC	B	16	2	-	0/2/19/22	0/1/1/1
2	GLC	B	17	2	-	2/2/19/22	0/1/1/1
2	GLC	B	2	2	-	0/2/19/22	0/1/1/1
2	GLC	B	3	2	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	4	2	-	0/2/19/22	0/1/1/1
2	GLC	B	5	2	-	0/2/19/22	0/1/1/1
2	GLC	B	6	2	-	0/2/19/22	0/1/1/1
2	GLC	B	7	2	-	0/2/19/22	0/1/1/1
2	GLC	B	8	2	-	2/2/19/22	0/1/1/1
2	GLC	B	9	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	14	GLC	C1-O5-C5	3.38	116.72	112.19
2	B	4	GLC	C1-O5-C5	2.81	115.95	112.19
2	B	6	GLC	C1-O5-C5	2.60	115.68	112.19
2	B	7	GLC	C1-O5-C5	2.59	115.66	112.19
2	B	17	GLC	C1-O5-C5	2.43	115.44	112.19
2	B	13	GLC	C1-O5-C5	2.27	115.23	112.19
2	B	10	GLC	C1-O5-C5	2.24	115.19	112.19
2	B	5	GLC	C1-C2-C3	2.20	112.85	109.64
2	B	3	GLC	C1-C2-C3	2.12	112.73	109.64

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	17	GLC	O5-C5-C6-O6
2	B	8	GLC	O5-C5-C6-O6
2	B	17	GLC	C4-C5-C6-O6
2	B	12	GLC	O5-C5-C6-O6
2	B	11	GLC	O5-C5-C6-O6
2	B	8	GLC	C4-C5-C6-O6
2	B	15	GLC	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	5	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	619	-	3,3,3	0.49	0	2,2,2	0.27	0
4	CO3	A	620	-	3,3,3	0.84	0	2,3,3	0.10	0
3	EDO	A	618	-	3,3,3	0.42	0	2,2,2	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	619	-	-	1/1/1/1	-
3	EDO	A	618	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	619	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	618	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	499/500 (99%)	0.27	59 (11%) 9 12	17, 41, 114, 185	8 (1%)

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	251	PHE	7.8
1	A	343	GLY	6.1
1	A	119	ALA	5.1
1	A	87	ASP	5.0
1	A	250	TYR	4.7
1	A	467	GLY	4.7
1	A	121	PRO	4.6
1	A	92	GLN	4.5
1	A	91	PRO	4.3
1	A	344	VAL	4.3
1	A	249	ASP	4.1
1	A	176	SER	3.8
1	A	30	ARG	3.6
1	A	468[A]	ARG	3.5
1	A	252	SER	3.4
1	A	289	LEU	3.4
1	A	84	ARG	3.4
1	A	154	LEU	3.4
1	A	1	MET	3.3
1	A	2	GLU	3.1
1	A	123	GLU	3.1
1	A	85	LEU	3.0
1	A	34	ARG	3.0
1	A	181	GLU	3.0
1	A	254	THR	2.9
1	A	469	PRO	2.9
1	A	60	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	248	PRO	2.8
1	A	281	GLU	2.8
1	A	239	ARG	2.8
1	A	79	GLU	2.8
1	A	90	PHE	2.7
1	A	151	HIS	2.7
1	A	470	SER	2.6
1	A	120	SER	2.5
1	A	326	GLN	2.4
1	A	4	PRO	2.4
1	A	88	PRO	2.4
1	A	99	LEU	2.4
1	A	139	GLU	2.4
1	A	122	GLU	2.3
1	A	150	ALA	2.3
1	A	153	GLY	2.3
1	A	314	GLY	2.3
1	A	350	GLU	2.3
1	A	82	TYR	2.2
1	A	86	GLU	2.2
1	A	311	ALA	2.2
1	A	133	ARG	2.2
1	A	310	THR	2.2
1	A	312	VAL	2.2
1	A	313	GLU	2.1
1	A	113	ARG	2.1
1	A	89	GLY	2.1
1	A	178	LEU	2.1
1	A	322	GLU	2.1
1	A	278	ARG	2.1
1	A	160	PRO	2.1
1	A	445	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SNN	A	370	8/8	0.95	0.11	29,36,42,42	0

6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	B	9	11/12	0.71	0.18	84,109,144,144	0
2	GLC	B	16	11/12	0.71	0.17	102,119,143,152	0
2	GLC	B	10	11/12	0.76	0.18	77,121,152,171	0
2	GLC	B	15	11/12	0.77	0.15	69,102,130,132	0
2	BGC	B	1	11/12	0.78	0.18	52,114,148,156	0
2	GLC	B	14	11/12	0.78	0.18	74,98,118,124	0
2	GLC	B	17	11/12	0.79	0.15	108,127,153,164	0
2	GLC	B	4	11/12	0.80	0.23	51,73,111,116	0
2	GLC	B	11	11/12	0.82	0.24	97,151,182,183	0
2	GLC	B	6	11/12	0.86	0.15	33,43,53,57	0
2	GLC	B	12	11/12	0.86	0.14	55,74,88,92	0
2	GLC	B	8	11/12	0.87	0.15	49,72,91,91	0
2	GLC	B	13	11/12	0.87	0.17	58,72,98,115	0
2	GLC	B	5	11/12	0.87	0.14	38,57,87,105	0
2	GLC	B	7	11/12	0.89	0.16	28,43,122,128	0
2	GLC	B	3	11/12	0.93	0.09	34,42,55,55	0
2	GLC	B	2	11/12	0.96	0.08	34,44,54,59	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CO3	A	620	4/4	0.83	0.19	81,91,99,113	0
3	EDO	A	618	4/4	0.85	0.21	85,102,123,123	0
3	EDO	A	619	4/4	0.88	0.12	48,66,80,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.