



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:44 AM UTC

PDB ID : 7JIU / pdb_00007jiu
Title : HUMAN PI3KDELTA IN COMPLEX WITH COMPOUND 2F
Authors : Lesburg, C.A.; Augustin, M.
Deposited on : 2020-07-23
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8290 (2.14-2.10)
Clashscore	190562	8817 (2.14-2.10)
Ramachandran outliers	187476	8738 (2.14-2.10)
Sidechain outliers	187428	8739 (2.14-2.10)
RSRZ outliers	180081	8294 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	946	13% 92% 5%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 7738 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic sub-unit alpha isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	900	Total	C	N	O	S	274	5	0
			7409	4724	1274	1349	62			

- Molecule 2 is (3S)-3-benzyl-5-[9-ethyl-8-(2-methylpyrimidin-5-yl)-9H-purin-6-yl]-3-methyl-1,3-dihydro-2H-indol-2-one (CCD ID: VBS) (formula: C₂₈H₂₅N₇O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			36	28	7	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	289	Total	O	0	4
			293	293		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.46Å 135.13Å 142.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	97.91 – 2.12 97.91 – 2.12	Depositor EDS
% Data completeness (in resolution range)	99.9 (97.91-2.12) 99.9 (97.91-2.12)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 2.12Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.212 , 0.246 (Not available) , 0.240	Depositor DCC
R_{free} test set	1612 reflections (2.49%)	wwPDB-VP
Wilson B-factor (Å ²)	38.1	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.014 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7738	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: VBS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	4/7570 (0.1%)	0.87	3/10221 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	889	ILE	C-N	-21.76	1.05	1.33
1	A	888	GLU	C-N	-11.27	1.20	1.34
1	A	720	LYS	C-N	6.33	1.42	1.33
1	A	726	GLU	C-N	-5.38	1.26	1.33

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	775	ALA	N-CA-C	13.79	128.01	111.33
1	A	719	LEU	O-C-N	9.10	131.76	122.12
1	A	776	LYS	N-CA-CB	-7.60	98.82	110.44

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	720	LYS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7409	0	7364	12	0
2	A	36	0	0	0	0
3	A	293	0	0	2	0
All	All	7738	0	7364	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:720:LYS:HE2	1:A:803:ASN:O	2.03	0.58
1:A:889:ILE:O	1:A:890:TYR:C	2.53	0.49
1:A:717:ASP:OD2	3:A:1203:HOH:O	2.20	0.47
1:A:194:ILE:HD12	1:A:209:LEU:HD11	1.99	0.45
1:A:812:LEU:HD22	1:A:1010:MET:HE1	1.98	0.44
1:A:446:TRP:CZ3	1:A:679:THR:HG22	2.53	0.43
1:A:802:LYS:NZ	3:A:1207:HOH:O	2.50	0.43
1:A:357:ARG:HD2	1:A:372:ASN:OD1	2.19	0.43
1:A:881:LEU:HD11	1:A:927:GLY:HA2	2.01	0.42
1:A:194:ILE:HD11	1:A:220:VAL:CG1	2.50	0.41
1:A:194:ILE:HD12	1:A:209:LEU:CD1	2.50	0.41
1:A:661:GLN:NE2	1:A:698:TYR:HB2	2.36	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	887/946 (94%)	858 (97%)	28 (3%)	1 (0%)	48	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	774	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	828/860 (96%)	823 (99%)	5 (1%)	78	85

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	330	ILE
1	A	376	VAL
1	A	774	SER
1	A	834	LEU
1	A	937	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	189	GLN
1	A	370	ASN
1	A	510	HIS
1	A	682	GLN
1	A	759	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	VBS	A	1101	-	41,41,41	0.95	1 (2%)	59,61,61	2.25	23 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	VBS	A	1101	-	-	6/15/31/31	0/6/6/6

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	VBS	C1-N27	-2.86	1.34	1.39

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	VBS	C1-N27-C26	6.22	110.34	104.29
2	A	1101	VBS	N3-C2-N25	5.62	134.27	125.09
2	A	1101	VBS	C1-C2-N3	-5.50	119.14	126.72
2	A	1101	VBS	C2-C1-N27	-4.20	106.62	110.91
2	A	1101	VBS	C6-C1-C2	4.06	118.52	116.63
2	A	1101	VBS	C16-C9-C10	-3.58	106.07	109.26
2	A	1101	VBS	N5-C4-N3	-3.48	123.31	128.58
2	A	1101	VBS	C4-N3-C2	3.39	120.11	111.83
2	A	1101	VBS	N25-C26-N27	-3.31	109.65	112.96
2	A	1101	VBS	C28-C33-N32	-3.27	118.78	123.59
2	A	1101	VBS	C9-C16-C14	3.01	103.68	101.83
2	A	1101	VBS	C34-C31-N32	2.97	120.36	117.20
2	A	1101	VBS	C33-N32-C31	2.96	120.93	116.07
2	A	1101	VBS	C29-N30-C31	2.96	120.93	116.07
2	A	1101	VBS	C28-C29-N30	-2.87	119.37	123.59
2	A	1101	VBS	C7-C6-N5	2.83	119.58	115.40
2	A	1101	VBS	C11-C10-C9	-2.77	119.33	121.92
2	A	1101	VBS	C7-C6-C1	-2.22	122.68	124.72
2	A	1101	VBS	C4-N5-C6	2.14	121.19	117.87
2	A	1101	VBS	C8-C9-C16	2.13	133.11	130.58
2	A	1101	VBS	C29-C28-C33	2.12	117.94	115.36
2	A	1101	VBS	N32-C31-N30	-2.09	122.07	125.46
2	A	1101	VBS	C6-C1-N27	2.07	134.86	133.18

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1101	VBS	C36-C35-N25-C26
2	A	1101	VBS	C36-C35-N25-C2
2	A	1101	VBS	N5-C6-C7-C12
2	A	1101	VBS	C1-C6-C7-C12
2	A	1101	VBS	N5-C6-C7-C8
2	A	1101	VBS	C1-C6-C7-C8

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	888:GLU	C	889:ILE	N	1.20
1	A	889:ILE	C	890:TYR	N	1.05

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	900/946 (95%)	0.67	120 (13%) 7 7	17, 44, 77, 118	87 (9%)

All (120) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	872	PHE	5.0
1	A	246	TYR	4.9
1	A	871	GLN	4.2
1	A	198	VAL	4.2
1	A	869	ALA	3.8
1	A	300	ASP	3.6
1	A	338	ILE	3.6
1	A	438	SER	3.6
1	A	377	PRO	3.5
1	A	178	PRO	3.5
1	A	378	CYS	3.5
1	A	353	LYS	3.5
1	A	858	MET	3.4
1	A	384	ASN	3.4
1	A	867	LYS	3.4
1	A	192	VAL	3.3
1	A	742	PRO	3.3
1	A	344	VAL	3.2
1	A	744	PHE	3.2
1	A	356	VAL	3.2
1	A	422	LEU	3.2
1	A	866	LEU	3.2
1	A	450	HIS	3.2
1	A	521	ASN	3.2
1	A	345	ASN	3.1
1	A	375	ARG	3.1
1	A	329	VAL	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	225	ILE	3.1
1	A	381	PRO	3.1
1	A	1008	SER	3.1
1	A	479	TRP	3.0
1	A	331	ASN	3.0
1	A	177	LEU	3.0
1	A	409	VAL	3.0
1	A	449	PRO	3.0
1	A	520	ASP	3.0
1	A	354	ILE	3.0
1	A	459	ILE	3.0
1	A	451	GLY	3.0
1	A	379	SER	3.0
1	A	484	VAL	3.0
1	A	949	ARG	2.9
1	A	862	CYS	2.9
1	A	473	LEU	2.9
1	A	437	VAL	2.9
1	A	322	THR	2.9
1	A	339	LEU	2.8
1	A	870	LEU	2.8
1	A	1052	THR	2.8
1	A	247	GLN	2.8
1	A	452	LEU	2.8
1	A	887	GLY	2.8
1	A	941	LYS	2.8
1	A	376	VAL	2.8
1	A	406	ILE	2.8
1	A	196	VAL	2.7
1	A	340	CYS	2.7
1	A	310	ARG	2.7
1	A	197	ILE	2.7
1	A	346	VAL	2.7
1	A	951	ARG	2.7
1	A	278	MET	2.6
1	A	522	GLU	2.6
1	A	347	ASN	2.6
1	A	475	LEU	2.6
1	A	725	ASP	2.6
1	A	351	ILE	2.6
1	A	863	LYS	2.6
1	A	477	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	374	GLN	2.5
1	A	775	ALA	2.5
1	A	232	MET	2.5
1	A	788	ILE	2.5
1	A	387	LEU	2.5
1	A	383	TRP	2.5
1	A	181	ILE	2.5
1	A	726	GLU	2.5
1	A	408	SER	2.4
1	A	524	ARG	2.4
1	A	721	GLN	2.4
1	A	407	CYS	2.4
1	A	158	SER	2.4
1	A	183	ASN	2.4
1	A	889	ILE	2.4
1	A	229	THR	2.4
1	A	1051	TRP	2.4
1	A	622	TYR	2.3
1	A	350	ASP	2.3
1	A	382	ARG	2.3
1	A	540	LEU	2.3
1	A	155[A]	ASP	2.3
1	A	480	PHE	2.3
1	A	156	LEU	2.3
1	A	860	ILE	2.3
1	A	436	LEU	2.2
1	A	1047	HIS	2.2
1	A	641	TYR	2.2
1	A	174	SER	2.2
1	A	386	TRP	2.2
1	A	441	MET	2.2
1	A	483	VAL	2.2
1	A	1035	ALA	2.2
1	A	295	SER	2.2
1	A	194	ILE	2.1
1	A	216	VAL	2.1
1	A	355	TYR	2.1
1	A	380	ASN	2.1
1	A	180	HIS	2.1
1	A	720	LYS	2.1
1	A	289	ALA	2.1
1	A	173	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	287	LEU	2.1
1	A	107	ASN	2.1
1	A	939	ASP	2.1
1	A	546	GLN	2.1
1	A	519	ARG	2.0
1	A	175	PRO	2.0
1	A	210	LYS	2.0
1	A	1009	GLY	2.0
1	A	1045	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	VBS	A	1101	36/36	0.86	0.18	40,64,96,98	6

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.