



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:10 PM UTC

PDB ID : 2JJP / pdb_00002jjp
Title : Structure of cytochrome P450 EryK in complex with inhibitor ketoconazole (KC)
Authors : Savino, C.; Sciara, G.; Miele, A.E.; Kendrew, S.G.; Vallone, B.
Deposited on : 2008-04-15
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	411	9% 71% 24% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3470 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYTOCHROME P450 113A1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	397	3152	1962	553	626	11	0	22	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	15	LEU	MET	engineered mutation	UNP P48635
A	344	LEU	PHE	engineered mutation	UNP P48635


- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0


- Molecule 3 is 1-ACETYL-4-(4-{{[(2S,4R)-2-(2,4-DICHLOROPHENYL)-2-(1H-IMIDAZOL

-1-YLMETHYL)-1,3-DIOXOLAN-4-YL[METHOXY}PHENYL)PIPERAZINE (CCD ID: KLN) (formula: C₂₆H₂₈Cl₂N₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0
			36	26	2	4	4	

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	S	0
			5	4	1	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	229	Total	O	0	0
			229	229		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYTOCHROME P450 113A1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.24Å 68.04Å 57.69Å 90.00° 100.71° 90.00°	Depositor
Resolution (Å)	56.70 – 2.10 56.68 – 2.10	Depositor EDS
% Data completeness (in resolution range)	91.4 (56.70-2.10) 91.4 (56.68-2.10)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.195 , 0.262 (Not available) , 0.275	Depositor DCC
R_{free} test set	1287 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	24.4	Xtriage
Anisotropy	0.875	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3470	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, SO4, KLN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	1/3332 (0.0%)	1.06	4/4550 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	92	PRO	CA-C	5.09	1.56	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	21	VAL	CA-C-N	-6.66	112.64	119.83
1	A	21	VAL	C-N-CA	-6.66	112.64	119.83
1	A	179	ASP	CA-C-N	5.33	125.14	119.28
1	A	179	ASP	C-N-CA	5.33	125.14	119.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3152	0	3059	85	0
2	A	43	0	30	10	0
3	A	36	0	28	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	10	0	0	0	0
5	A	229	0	0	10	0
All	All	3470	0	3117	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:311[B]:MET:HE3	1:A:313:ASN:HD21	1.03	1.08
1:A:311[B]:MET:HE3	1:A:313:ASN:ND2	1.80	0.95
1:A:291[A]:MET:CE	1:A:316:VAL:HG21	2.06	0.86
1:A:291[A]:MET:HE1	1:A:316:VAL:HG21	1.60	0.82
1:A:20:GLU:HG2	1:A:21:VAL:H	1.47	0.79
1:A:109:ARG:HH11	1:A:109:ARG:HG2	1.49	0.77
1:A:163:GLY:HA3	1:A:166:SER:OG	1.89	0.72
1:A:291[A]:MET:HE2	2:A:412:HEM:CMA	2.23	0.69
1:A:291[A]:MET:HE2	2:A:412:HEM:HMA1	1.76	0.68
1:A:165:TRP:HE1	1:A:240[A]:LEU:HD21	1.57	0.67
1:A:178:THR:HA	1:A:311[A]:MET:HE1	1.77	0.67
1:A:390[B]:GLU:OE1	5:A:2218:HOH:O	2.09	0.67
1:A:165:TRP:HE1	1:A:240[A]:LEU:CD2	2.09	0.64
1:A:17:THR:HG22	1:A:17:THR:O	1.97	0.64
1:A:331:ARG:HD2	5:A:2187:HOH:O	1.97	0.64
1:A:20:GLU:CG	1:A:21:VAL:N	2.62	0.62
2:A:412:HEM:HMB1	2:A:412:HEM:HBB2	1.81	0.62
1:A:109:ARG:CZ	5:A:2077:HOH:O	2.48	0.60
1:A:20:GLU:HG2	1:A:21:VAL:N	2.16	0.59
1:A:226:LEU:HD22	1:A:230:GLU:HB3	1.85	0.59
1:A:336:ARG:O	1:A:336:ARG:HG3	2.03	0.58
1:A:172:ILE:HG23	1:A:174:MET:HB2	1.87	0.56
1:A:150:GLU:HG3	5:A:2107:HOH:O	2.06	0.55
1:A:136:VAL:HA	1:A:140:ALA:HB3	1.89	0.54
1:A:80:ALA:O	1:A:82:PRO:HD3	2.08	0.53
1:A:201:CYS:HB2	1:A:228[A]:ASP:OD1	2.08	0.53
1:A:356:ALA:O	1:A:360:ARG:HG3	2.07	0.53
1:A:96:ARG:NH1	5:A:2072:HOH:O	2.42	0.53
1:A:20:GLU:CG	1:A:21:VAL:H	2.16	0.53
1:A:109:ARG:HH11	1:A:109:ARG:CG	2.21	0.52
1:A:193:LEU:HD23	1:A:236:THR:HG23	1.90	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:200:ARG:HG2	1:A:200:ARG:HH11	1.73	0.52
2:A:412:HEM:HMC1	2:A:412:HEM:HBC2	1.90	0.52
1:A:162:PHE:CE1	1:A:247:THR:HG21	2.44	0.52
1:A:174:MET:HE1	3:A:413:KLN:H263	1.92	0.52
1:A:224:ARG:HH12	1:A:230:GLU:CD	2.18	0.51
1:A:165:TRP:CH2	1:A:243:HIS:CE1	2.99	0.51
1:A:200:ARG:HG2	1:A:200:ARG:NH1	2.26	0.51
1:A:289:PRO:HD2	5:A:2220:HOH:O	2.12	0.50
1:A:331:ARG:HD2	1:A:333:ASP:HB2	1.93	0.49
1:A:109:ARG:HD3	1:A:109:ARG:H	1.78	0.49
1:A:165:TRP:CD1	1:A:165:TRP:C	2.91	0.49
1:A:20:GLU:O	1:A:21:VAL:HG23	2.13	0.49
1:A:389:PHE:CZ	1:A:397:ARG:HG2	2.48	0.49
1:A:165:TRP:CZ3	1:A:243:HIS:CE1	3.03	0.47
1:A:280:GLU:HG3	1:A:332:PHE:CE1	2.49	0.47
1:A:80:ALA:C	1:A:82:PRO:HD3	2.40	0.47
1:A:239:LEU:O	1:A:243:HIS:HD2	1.97	0.47
1:A:291[B]:MET:HE3	2:A:412:HEM:CMA	2.44	0.47
1:A:134:ASP:OD1	1:A:137[A]:ASP:CG	2.58	0.46
2:A:412:HEM:HBB2	2:A:412:HEM:CMB	2.45	0.46
1:A:56:HIS:CG	1:A:325:ALA:HB2	2.50	0.46
1:A:18:ILE:O	1:A:18:ILE:HG23	2.16	0.46
1:A:392:ILE:HD12	3:A:413:KLN:H242	1.98	0.46
1:A:237:ALA:HB1	3:A:413:KLN:CL1	2.52	0.46
1:A:41:GLN:NE2	5:A:2021:HOH:O	2.47	0.46
1:A:148:VAL:HG21	1:A:242:GLY:HA3	1.98	0.45
1:A:86:MET:HE3	1:A:88:HIS:HE1	1.81	0.45
1:A:43:VAL:HB	1:A:304:VAL:HG21	1.99	0.45
2:A:412:HEM:C4C	3:A:413:KLN:H2	2.52	0.44
1:A:201:CYS:CB	1:A:228[A]:ASP:OD1	2.65	0.44
1:A:222:ASP:OD1	1:A:222:ASP:N	2.49	0.44
1:A:227:ASP:OD1	1:A:229[A]:GLU:HB3	2.18	0.44
1:A:255:ARG:NE	5:A:2148:HOH:O	2.49	0.44
2:A:412:HEM:HBC2	2:A:412:HEM:CMC	2.48	0.44
1:A:109:ARG:O	1:A:113:ASP:OD2	2.35	0.44
1:A:21:VAL:O	1:A:22:PRO:C	2.60	0.43
1:A:87:ILE:HD11	1:A:99:ARG:HA	2.00	0.43
1:A:392:ILE:CD1	3:A:413:KLN:H242	2.49	0.43
1:A:280:GLU:HA	1:A:280:GLU:OE1	2.18	0.43
1:A:245:THR:HB	2:A:412:HEM:CAB	2.49	0.42
1:A:106:PHE:HD1	1:A:212:LEU:HD21	1.84	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:280:GLU:HG3	1:A:332:PHE:CD1	2.54	0.42
1:A:380:ASP:OD1	1:A:382:ASP:HB2	2.18	0.42
1:A:86:MET:HE3	1:A:88:HIS:CE1	2.55	0.42
1:A:174:MET:CE	1:A:177:PRO:HG3	2.50	0.41
1:A:60:GLN:NE2	5:A:2040:HOH:O	2.52	0.41
1:A:190:LEU:HD23	1:A:190:LEU:HA	1.79	0.41
1:A:86:MET:HE1	1:A:237:ALA:HB2	2.03	0.41
1:A:270:ASP:HB3	1:A:273:ARG:HD2	2.02	0.41
1:A:291[B]:MET:SD	2:A:412:HEM:HBA2	2.61	0.41
1:A:118:ILE:HG22	1:A:365:VAL:HG11	2.03	0.41
1:A:79:GLY:O	1:A:81:SER:N	2.51	0.41
1:A:203[A]:GLU:HG2	5:A:2129:HOH:O	2.21	0.41
1:A:132:SER:HB2	1:A:402:LEU:HD23	2.04	0.40
1:A:143:LEU:O	1:A:147:ILE:HG12	2.21	0.40
1:A:17:THR:O	1:A:17:THR:CG2	2.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	417/411 (102%)	387 (93%)	26 (6%)	4 (1%)	12 9

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	341	ALA
1	A	80	ALA
1	A	21	VAL
1	A	160	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	353/343 (103%)	326 (92%)	27 (8%)	12 9

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	THR
1	A	18	ILE
1	A	28	THR
1	A	74	THR
1	A	77	ILE
1	A	78[A]	GLU
1	A	78[B]	GLU
1	A	96	ARG
1	A	100	LYS
1	A	109	ARG
1	A	157	MET
1	A	164	ASP
1	A	165	TRP
1	A	169	LEU
1	A	170	VAL
1	A	210[A]	ASP
1	A	210[B]	ASP
1	A	222	ASP
1	A	238	LEU
1	A	244	ILE
1	A	291[A]	MET
1	A	291[B]	MET
1	A	331	ARG
1	A	338	SER
1	A	384	GLU
1	A	386	LEU
1	A	410	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	45	GLN
1	A	60	GLN
1	A	88	HIS
1	A	161	GLN
1	A	243	HIS
1	A	313	ASN
1	A	363	ASN
1	A	391	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	415	-	4,4,4	0.27	0	6,6,6	0.44	0
4	SO4	A	414	-	4,4,4	0.23	0	6,6,6	0.22	0
3	KLN	A	413	2	40,40,40	1.00	2 (5%)	57,57,57	2.05	11 (19%)
2	HEM	A	412	1,3	50,50,50	1.85	9 (18%)	67,82,82	1.44	9 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KLN	A	413	2	-	10/24/45/45	0/5/5/5
2	HEM	A	412	1,3	-	0/14/54/54	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	412	HEM	C3D-C2D	8.12	1.54	1.36
2	A	412	HEM	FE-NB	3.76	2.06	1.94
2	A	412	HEM	FE-ND	3.34	2.05	1.94
2	A	412	HEM	CAC-C3C	3.11	1.55	1.47
3	A	413	KLN	C5-C8	-3.08	1.50	1.53
2	A	412	HEM	FE-NC	3.02	2.05	1.95
3	A	413	KLN	C25-N4	2.93	1.43	1.35
2	A	412	HEM	CAB-C3B	2.88	1.55	1.47
2	A	412	HEM	CMC-C2C	2.35	1.55	1.50
2	A	412	HEM	CAA-C2A	2.23	1.57	1.51
2	A	412	HEM	CMD-C2D	2.17	1.55	1.50

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	413	KLN	C5-C4-N1	-7.99	105.81	112.76
3	A	413	KLN	C22-N4-C25	-5.22	108.91	122.96
3	A	413	KLN	C23-N4-C25	-4.98	109.57	122.96
2	A	412	HEM	C4D-ND-C1D	4.91	111.03	105.21
3	A	413	KLN	C13-C8-C9	4.52	121.32	116.51
3	A	413	KLN	C5-C8-C9	-3.46	121.27	123.72
3	A	413	KLN	O2-C5-C8	-3.43	105.49	111.37
3	A	413	KLN	O3-C14-C6	3.36	115.53	107.79
2	A	412	HEM	CHB-C1B-NB	3.21	128.33	124.37
3	A	413	KLN	C4-C5-C8	3.07	113.90	109.59
2	A	412	HEM	C3B-C2B-C1B	3.02	108.68	106.41
2	A	412	HEM	CHD-C4C-NC	2.88	127.59	124.45
3	A	413	KLN	C13-C8-C5	-2.68	117.41	119.18
2	A	412	HEM	CAD-C3D-C2D	2.55	132.65	127.87
2	A	412	HEM	C1B-NB-C4B	2.52	108.19	105.21
2	A	412	HEM	CMB-C2B-C1B	-2.50	121.12	125.03
3	A	413	KLN	C21-N3-C18	-2.42	111.50	118.11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	413	KLN	C10-C9-C8	-2.42	119.56	121.97
2	A	412	HEM	C3B-C4B-NB	-2.16	107.92	109.47
2	A	412	HEM	C2B-C1B-NB	-2.03	107.50	109.84

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	413	KLN	N1-C4-C5-O1
3	A	413	KLN	N1-C4-C5-C8
3	A	413	KLN	O3-C14-C6-O1
3	A	413	KLN	O4-C25-N4-C22
3	A	413	KLN	C26-C25-N4-C22
3	A	413	KLN	O4-C25-N4-C23
3	A	413	KLN	C26-C25-N4-C23
3	A	413	KLN	C16-C15-O3-C14
3	A	413	KLN	C20-C15-O3-C14
3	A	413	KLN	N1-C4-C5-O2


There are no ring outliers.

2 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	413	KLN	5	0
2	A	412	HEM	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand KLN A 413



Ligand HEM A 412



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	397/411 (96%)	0.33	36 (9%) 15 15	16, 26, 66, 81	22 (5%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	169	LEU	3.8
1	A	162	PHE	3.7
1	A	165	TRP	3.6
1	A	164	ASP	3.6
1	A	181	ALA	3.3
1	A	81	SER	3.2
1	A	383[A]	ASP	3.2
1	A	177	PRO	3.0
1	A	18	ILE	3.0
1	A	77	ILE	3.0
1	A	79	GLY	3.0
1	A	80	ALA	3.0
1	A	175	ASP	2.9
1	A	160	GLU	2.9
1	A	163	GLY	2.9
1	A	21	VAL	2.8
1	A	159	HIS	2.8
1	A	156	PRO	2.8
1	A	170	VAL	2.8
1	A	339	GLY	2.8
1	A	341	ALA	2.8
1	A	340	GLY	2.7
1	A	176	ASP	2.5
1	A	168	ALA	2.5
1	A	173	GLN	2.5
1	A	161	GLN	2.4
1	A	178	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	75	ARG	2.4
1	A	158	ASP	2.3
1	A	157	MET	2.3
1	A	228[A]	ASP	2.3
1	A	338	SER	2.2
1	A	385	ARG	2.2
1	A	15	LEU	2.2
1	A	20	GLU	2.1
1	A	172	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	A	415	5/5	0.77	0.14	66,66,68,69	0
3	KLN	A	413	36/36	0.86	0.13	25,52,56,57	0
4	SO4	A	414	5/5	0.92	0.11	55,55,56,57	0
2	HEM	A	412	43/43	0.98	0.05	14,18,21,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around KLN A 413:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.