



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 06:53 PM UTC

PDB ID : 5JJ4 / pdb_00005jj4
Title : Crystal Structure of a Variant Human Activation-induced Deoxycytidine Deaminase as an MBP fusion protein
Authors : Pedersen, L.C.; Goodman, M.F.; Pham, P.; Afif, S.A.
Deposited on : 2016-04-22
Resolution : 2.81 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	546	 97%
1	B	546	 98%
1	C	546	 96%
2	D	4	 100%
2	E	4	 25% 75%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	4	 A horizontal bar chart showing the quality of chain F. The bar is divided into three segments: green (50%), yellow (25%), and orange (25%). The percentages are labeled below each segment.

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12734 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein, Single-stranded DNA cytosine deaminase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	541	Total	C	N	O	S	0	1	0
			4221	2722	700	784	15			
1	A	542	Total	C	N	O	S	0	0	0
			4187	2696	692	784	15			
1	B	545	Total	C	N	O	S	0	0	0
			4149	2681	680	773	15			

There are 81 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	359	ALA	GLU	conflict	UNP P0AEY0
C	362	ALA	LYS	conflict	UNP P0AEY0
C	363	ALA	ASP	conflict	UNP P0AEY0
C	367	ASN	ARG	conflict	UNP P0AEY0
C	368	ALA	-	linker	UNP P0AEY0
C	369	ALA	-	linker	UNP P0AEY0
C	370	ALA	-	linker	UNP P0AEY0
C	1005	LEU	-	linker	UNP P0AEY0
C	1006	MET	-	linker	UNP P0AEY0
C	1007	ASP	-	linker	UNP P0AEY0
C	1008	PRO	-	linker	UNP P0AEY0
C	1009	HIS	-	linker	UNP P0AEY0
C	1010	ILE	-	linker	UNP P0AEY0
C	1011	PHE	-	linker	UNP P0AEY0
C	1012	THR	-	linker	UNP P0AEY0
C	1013	SER	-	linker	UNP P0AEY0
C	1014	ASN	-	linker	UNP P0AEY0
C	1015	PHE	-	linker	UNP P0AEY0
C	1016	ASN	-	linker	UNP P0AEY0
C	1017	ASN	-	linker	UNP P0AEY0
C	1018	GLY	-	linker	UNP P0AEY0
C	1019	ILE	-	linker	UNP P0AEY0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1025	HIS	ARG	conflict	UNP Q9GZX7
C	1026	LYS	GLU	conflict	UNP Q9GZX7
C	1032	GLU	VAL	conflict	UNP Q9GZX7
C	1034	GLU	LYS	conflict	UNP Q9GZX7
C	1036	LEU	ARG	conflict	UNP Q9GZX7
A	359	ALA	GLU	conflict	UNP P0AEY0
A	362	ALA	LYS	conflict	UNP P0AEY0
A	363	ALA	ASP	conflict	UNP P0AEY0
A	367	ASN	ARG	conflict	UNP P0AEY0
A	368	ALA	-	linker	UNP P0AEY0
A	369	ALA	-	linker	UNP P0AEY0
A	370	ALA	-	linker	UNP P0AEY0
A	1005	LEU	-	linker	UNP P0AEY0
A	1006	MET	-	linker	UNP P0AEY0
A	1007	ASP	-	linker	UNP P0AEY0
A	1008	PRO	-	linker	UNP P0AEY0
A	1009	HIS	-	linker	UNP P0AEY0
A	1010	ILE	-	linker	UNP P0AEY0
A	1011	PHE	-	linker	UNP P0AEY0
A	1012	THR	-	linker	UNP P0AEY0
A	1013	SER	-	linker	UNP P0AEY0
A	1014	ASN	-	linker	UNP P0AEY0
A	1015	PHE	-	linker	UNP P0AEY0
A	1016	ASN	-	linker	UNP P0AEY0
A	1017	ASN	-	linker	UNP P0AEY0
A	1018	GLY	-	linker	UNP P0AEY0
A	1019	ILE	-	linker	UNP P0AEY0
A	1025	HIS	ARG	conflict	UNP Q9GZX7
A	1026	LYS	GLU	conflict	UNP Q9GZX7
A	1032	GLU	VAL	conflict	UNP Q9GZX7
A	1034	GLU	LYS	conflict	UNP Q9GZX7
A	1036	LEU	ARG	conflict	UNP Q9GZX7
B	359	ALA	GLU	conflict	UNP P0AEY0
B	362	ALA	LYS	conflict	UNP P0AEY0
B	363	ALA	ASP	conflict	UNP P0AEY0
B	367	ASN	ARG	conflict	UNP P0AEY0
B	368	ALA	-	linker	UNP P0AEY0
B	369	ALA	-	linker	UNP P0AEY0
B	370	ALA	-	linker	UNP P0AEY0
B	1005	LEU	-	linker	UNP P0AEY0
B	1006	MET	-	linker	UNP P0AEY0
B	1007	ASP	-	linker	UNP P0AEY0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1008	PRO	-	linker	UNP P0AEY0
B	1009	HIS	-	linker	UNP P0AEY0
B	1010	ILE	-	linker	UNP P0AEY0
B	1011	PHE	-	linker	UNP P0AEY0
B	1012	THR	-	linker	UNP P0AEY0
B	1013	SER	-	linker	UNP P0AEY0
B	1014	ASN	-	linker	UNP P0AEY0
B	1015	PHE	-	linker	UNP P0AEY0
B	1016	ASN	-	linker	UNP P0AEY0
B	1017	ASN	-	linker	UNP P0AEY0
B	1018	GLY	-	linker	UNP P0AEY0
B	1019	ILE	-	linker	UNP P0AEY0
B	1025	HIS	ARG	conflict	UNP Q9GZX7
B	1026	LYS	GLU	conflict	UNP Q9GZX7
B	1032	GLU	VAL	conflict	UNP Q9GZX7
B	1034	GLU	LYS	conflict	UNP Q9GZX7
B	1036	LEU	ARG	conflict	UNP Q9GZX7

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	D	4	Total	C	O	0	0	0
			45	24	21			
2	E	4	Total	C	O	0	0	0
			45	24	21			
2	F	4	Total	C	O	0	0	0
			45	24	21			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total 1	Ca 1	0	0
4	A	1	Total 1	Ca 1	0	0
4	B	1	Total 1	Ca 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	12	Total 12	O 12	0	0
5	A	13	Total 13	O 13	0	0
5	B	11	Total 11	O 11	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Maltose-binding periplasmic protein,Single-stranded DNA cytosine deaminase

Chain C:  96%




- Molecule 1: Maltose-binding periplasmic protein,Single-stranded DNA cytosine deaminase

Chain A:  97%




- Molecule 1: Maltose-binding periplasmic protein,Single-stranded DNA cytosine deaminase

Chain B:  98%




- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain D:  100%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain E:  25% 75%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain F:  50% 25% 25%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	106.79Å 96.81Å 108.69Å 90.00° 102.86° 90.00°	Depositor
Resolution (Å)	35.48 – 2.81 35.48 – 2.81	Depositor EDS
% Data completeness (in resolution range)	99.4 (35.48-2.81) 93.1 (35.48-2.81)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.81Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.183 , 0.232 0.184 , 0.232	Depositor DCC
R_{free} test set	1991 reflections (3.76%)	wwPDB-VP
Wilson B-factor (Å ²)	66.5	Xtriage
Anisotropy	0.699	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12734	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ZN, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/4299	0.65	0/5863
1	B	0.31	0/4263	0.65	0/5824
1	C	0.33	0/4333	0.65	0/5898
All	All	0.32	0/12895	0.65	0/17585

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4187	0	3936	5	0
1	B	4149	0	3860	6	0
1	C	4221	0	4017	8	0
2	D	45	0	39	0	0
2	E	45	0	39	0	0
2	F	45	0	39	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	13	0	0	0	0
5	B	11	0	0	0	0
5	C	12	0	0	0	0
All	All	12734	0	11930	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 19 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:130:GLU:N	1:C:130:GLU:OE1	2.38	0.56
1:B:205:ASN:ND2	1:B:207:ASP:OD1	2.39	0.55
1:B:95:ASP:OD1	1:B:98:ARG:NH1	2.36	0.55
1:B:44:GLU:OE2	2:F:4:GLC:O2	2.19	0.51
1:C:245:THR:OG1	1:C:246:VAL:N	2.43	0.51

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	538/546 (98%)	518 (96%)	19 (4%)	1 (0%)	43	72
1	B	543/546 (100%)	520 (96%)	22 (4%)	1 (0%)	43	72
1	C	538/546 (98%)	524 (97%)	14 (3%)	0	100	100
All	All	1619/1638 (99%)	1562 (96%)	55 (3%)	2 (0%)	48	77

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1038	SER
1	A	165	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	412/450 (92%)	409 (99%)	3 (1%)	76	91
1	B	398/450 (88%)	396 (100%)	2 (0%)	81	93
1	C	419/450 (93%)	418 (100%)	1 (0%)	87	96
All	All	1229/1350 (91%)	1223 (100%)	6 (0%)	81	93

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	258	PHE
1	B	258	PHE
1	B	1156	GLU
1	A	119	LYS
1	C	258	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	367	ASN
1	A	1155	HIS
1	B	1135	GLN
1	B	367	ASN
1	C	1168	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	D	1	2	12,12,12	0.68	0	17,17,17	0.91	1 (5%)
2	GLC	D	2	2	11,11,12	1.34	2 (18%)	15,15,17	1.56	2 (13%)
2	GLC	D	3	2	11,11,12	1.15	1 (9%)	15,15,17	1.16	0
2	GLC	D	4	2	11,11,12	0.79	0	15,15,17	1.09	1 (6%)
2	GLC	E	1	2	12,12,12	0.70	0	17,17,17	0.85	0
2	GLC	E	2	2	11,11,12	1.09	0	15,15,17	1.01	1 (6%)
2	GLC	E	3	2	11,11,12	1.18	1 (9%)	15,15,17	1.56	5 (33%)
2	GLC	E	4	2	11,11,12	0.65	0	15,15,17	1.22	2 (13%)
2	GLC	F	1	2	12,12,12	0.77	0	17,17,17	0.88	0
2	GLC	F	2	2	11,11,12	1.21	2 (18%)	15,15,17	0.88	1 (6%)
2	GLC	F	3	2	11,11,12	1.09	0	15,15,17	1.03	0
2	GLC	F	4	2	11,11,12	0.71	0	15,15,17	1.20	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	3	2	-	0/2/19/22	0/1/1/1
2	GLC	D	4	2	-	0/2/19/22	0/1/1/1
2	GLC	E	1	2	-	1/2/22/22	0/1/1/1
2	GLC	E	2	2	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	E	3	2	-	0/2/19/22	0/1/1/1
2	GLC	E	4	2	-	0/2/19/22	0/1/1/1
2	GLC	F	1	2	-	2/2/22/22	0/1/1/1
2	GLC	F	2	2	-	0/2/19/22	0/1/1/1
2	GLC	F	3	2	-	1/2/19/22	0/1/1/1
2	GLC	F	4	2	-	2/2/19/22	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	GLC	C4-C5	-2.63	1.47	1.53
2	E	3	GLC	O5-C1	-2.51	1.39	1.43
2	D	3	GLC	O5-C1	-2.51	1.39	1.43
2	D	2	GLC	O2-C2	-2.46	1.38	1.43
2	F	2	GLC	C4-C5	-2.31	1.48	1.53

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	4	GLC	C6-C5-C4	-2.75	106.27	113.02
2	E	4	GLC	C1-C2-C3	2.67	113.53	109.64
2	D	2	GLC	O3-C3-C2	2.61	115.37	110.05
2	D	2	GLC	C6-C5-C4	-2.60	106.64	113.02
2	E	3	GLC	C1-O5-C5	2.43	115.44	112.19

There are no chirality outliers.

5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	F	4	GLC	O5-C5-C6-O6
2	F	4	GLC	C4-C5-C6-O6
2	F	1	GLC	C4-C5-C6-O6
2	F	1	GLC	O5-C5-C6-O6
2	F	3	GLC	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	4	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	542/546 (99%)	-0.42	0 100 100	39, 73, 108, 129	0
1	B	545/546 (99%)	-0.26	2 (0%) 88 84	48, 81, 117, 133	0
1	C	541/546 (99%)	-0.53	1 (0%) 91 88	27, 64, 91, 121	1 (0%)
All	All	1628/1638 (99%)	-0.40	3 (0%) 91 88	27, 72, 110, 133	1 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1181	LEU	3.8
1	B	347	VAL	2.0
1	B	124	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	E	4	11/12	0.51	0.39	44,56,59,61	11
2	GLC	F	4	11/12	0.71	0.23	59,61,65,67	11
2	GLC	D	4	11/12	0.83	0.22	52,54,58,59	11
2	GLC	D	1	12/12	0.93	0.09	46,57,60,62	0
2	GLC	F	3	11/12	0.95	0.08	55,59,64,66	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	E	1	12/12	0.95	0.11	56,59,66,67	0
2	GLC	F	1	12/12	0.96	0.06	59,63,69,76	0
2	GLC	E	3	11/12	0.96	0.07	55,56,61,67	0
2	GLC	D	3	11/12	0.96	0.08	46,50,57,58	0
2	GLC	F	2	11/12	0.97	0.05	49,54,58,59	0
2	GLC	E	2	11/12	0.97	0.04	47,57,63,67	0
2	GLC	D	2	11/12	0.97	0.05	30,40,44,47	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CA	B	1202	1/1	0.74	0.11	131,131,131,131	0
4	CA	A	1202	1/1	0.92	0.09	101,101,101,101	0
4	CA	C	1202	1/1	0.93	0.12	100,100,100,100	0
3	ZN	B	1201	1/1	0.99	0.02	71,71,71,71	0
3	ZN	C	1201	1/1	1.00	0.02	68,68,68,68	0
3	ZN	A	1201	1/1	1.00	0.01	67,67,67,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.