



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 08:38 PM UTC

PDB ID : 8JJC / pdb_00008jjc
Title : Tubulin-Y62
Authors : Yang, J.
Deposited on : 2023-05-30
Resolution : 2.76 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	4% 89% 8% .
1	C	451	3% 89% 9% .
2	B	431	6% 88% 10% .
2	D	431	9% 85% 13% .
3	E	189	8% 61% 35% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	380	 A horizontal bar chart showing the quality of chain F. The bar is divided into four segments: red (22%), green (81%), yellow (10%), and grey (8%). The percentages are labeled above or below the segments. The red segment is at the left, followed by green, then yellow, and grey at the right.

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 33734 atoms, of which 16410 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha-1B chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	438	Total	C	H	N	O	S	0	2	0
			6653	2146	3260	576	648	23			
1	C	440	Total	C	H	N	O	S	0	5	0
			6779	2183	3331	584	658	23			

- Molecule 2 is a protein called Tubulin beta chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	425	Total	C	H	N	O	S	0	1	0
			6414	2078	3111	562	638	25			
2	D	423	Total	C	H	N	O	S	0	0	0
			6361	2059	3090	556	632	24			


- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	E	123	Total	C	H	N	O	S	0	2	0
			1988	614	995	183	191	5			

- Molecule 4 is a protein called TTL.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	F	349	Total	C	H	N	O	S	0	0	0
			5225	1726	2533	462	490	14			

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total 42	C 10	H 10	N 5	O 14	P 3	42	0
5	C	1	Total 42	C 10	H 10	N 5	O 14	P 3	42	0
5	D	1	Total 42	C 10	H 10	N 5	O 14	P 3	42	0

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total 1	Mg 1	1	0
6	B	1	Total 1	Mg 1	1	0
6	C	1	Total 1	Mg 1	1	0
6	D	1	Total 1	Mg 1	1	0
6	F	1	Total 1	Mg 1	1	0

- Molecule 7 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	2	Total Ca 2 2	2	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	2	Total	Ca	2	0
			2	2		
7	C	1	Total	Ca	1	0
			1	1		
7	D	1	Total	Ca	1	0
			1	1		

- Molecule 8 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	B	1	Total	C	H	N	O	S	
			24	6	12	1	4	1	24
									0

- Molecule 9 is 4-(6,7-dimethoxy-3,4-dihydro-1 {H}-isoquinolin-2-yl)-6-(3-methoxyphenyl)pyrimidin-2-amine (CCD ID: UPO) (formula: $C_{22}H_{24}N_4O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	B	1	Total	C	H	N	O	0	0
			53	22	24	4	3		


- Molecule 10 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
10	B	1	Total	C	H	N	O	P	38	0
			38	10	10	5	11	2		

- Molecule 11 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

- Molecule 12 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).




- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|----------------|---------|---------|
| 13 | A | 1 | Total O
1 1 | 0 | 0 |
| 13 | B | 2 | Total O
2 2 | 0 | 0 |
| 13 | C | 9 | Total O
9 9 | 0 | 0 |
| 13 | D | 1 | Total O
1 1 | 0 | 0 |
| 13 | F | 3 | Total O
3 3 | 0 | 0 |


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha-1B chain




- Molecule 1: Tubulin alpha-1B chain



- Molecule 2: Tubulin beta chain



- Molecule 2: Tubulin beta chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	104.49Å 157.00Å 180.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.74 – 2.76 47.74 – 2.76	Depositor EDS
% Data completeness (in resolution range)	99.3 (47.74-2.76) 99.3 (47.74-2.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 2.77Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.235 , 0.264 0.240 , 0.262	Depositor DCC
R_{free} test set	1143 reflections (1.49%)	wwPDB-VP
Wilson B-factor (Å ²)	70.6	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	33734	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MES, GDP, CL, GTP, ACP, CA, UPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.17	0/3473	0.39	0/4723
1	C	0.18	0/3539	0.38	0/4810
2	B	0.17	0/3380	0.38	0/4588
2	D	0.17	0/3343	0.39	0/4537
3	E	0.17	0/1004	0.30	0/1336
4	F	0.15	0/2754	0.32	0/3741
All	All	0.17	0/17493	0.37	0/23735

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3393	3260	3271	24	0
1	C	3448	3331	3344	27	0
2	B	3303	3111	3141	26	0
2	D	3271	3090	3130	33	1
3	E	993	995	998	5	0
4	F	2692	2533	2545	27	1
5	A	32	10	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	32	10	12	0	0
5	D	32	10	12	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
6	F	1	0	0	0	0
7	A	2	0	0	0	0
7	B	2	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
8	B	12	12	12	0	0
9	B	29	24	0	0	0
10	B	28	10	12	0	0
11	D	1	0	0	0	0
12	F	31	14	14	0	0
13	A	1	0	0	0	0
13	B	2	0	0	0	0
13	C	9	0	0	0	0
13	D	1	0	0	1	0
13	F	3	0	0	0	0
All	All	17324	16410	16503	139	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 139 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:36:MET:HE3	1:C:39:ASP:HB2	1.62	0.79
3:E:116:LEU:O	3:E:120:LEU:HD12	1.88	0.72
2:B:293:MET:HE3	2:B:373:ALA:HB1	1.74	0.69
2:D:145:SER:OG	2:D:188:SER:OG	2.07	0.67
2:B:229:VAL:HG12	2:B:233:MET:HE2	1.78	0.66

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:213:ARG:HH22	4:F:324:GLU:OE1[3_545]	1.56	0.04

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/451 (97%)	429 (98%)	9 (2%)	0	100	100
1	C	443/451 (98%)	432 (98%)	11 (2%)	0	100	100
2	B	422/431 (98%)	409 (97%)	13 (3%)	0	100	100
2	D	419/431 (97%)	405 (97%)	13 (3%)	1 (0%)	43	64
3	E	121/189 (64%)	120 (99%)	1 (1%)	0	100	100
4	F	343/380 (90%)	326 (95%)	14 (4%)	3 (1%)	14	28
All	All	2186/2333 (94%)	2121 (97%)	61 (3%)	4 (0%)	43	64

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	246	LEU
4	F	144	GLY
4	F	243	HIS
4	F	258	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	362/379 (96%)	360 (99%)	2 (1%)	78	88
1	C	372/379 (98%)	371 (100%)	1 (0%)	86	93
2	B	355/370 (96%)	353 (99%)	2 (1%)	78	88

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	353/370 (95%)	349 (99%)	4 (1%)	65	79
3	E	104/171 (61%)	103 (99%)	1 (1%)	68	81
4	F	273/338 (81%)	269 (98%)	4 (2%)	57	74
All	All	1819/2007 (91%)	1805 (99%)	14 (1%)	73	84

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	306	ARG
2	D	355	ASP
4	F	243	HIS
4	F	241	THR
4	F	242	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
4	F	183	GLN
4	F	380	HIS
2	B	298	ASN
1	C	85	GLN
1	C	300	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 19 ligands modelled in this entry, 12 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
8	MES	B	502	-	12,12,12	2.30	1 (8%)	15,16,16	1.65	1 (6%)
5	GTP	D	503	6	33,34,34	0.94	1 (3%)	50,54,54	1.58	8 (16%)
9	UPO	B	505	-	32,32,32	1.66	7 (21%)	44,45,45	1.74	8 (18%)
12	ACP	F	402	6	31,33,33	1.62	7 (22%)	47,52,52	1.92	12 (25%)
5	GTP	C	501	6	33,34,34	0.95	1 (3%)	50,54,54	1.63	8 (16%)
10	GDP	B	506	6	29,30,30	1.15	3 (10%)	45,47,47	1.74	7 (15%)
5	GTP	A	501	6	33,34,34	0.92	1 (3%)	50,54,54	1.57	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	MES	B	502	-	-	1/6/14/14	0/1/1/1
5	GTP	D	503	6	-	4/22/38/38	0/3/3/3
9	UPO	B	505	-	-	4/14/23/23	0/4/4/4
12	ACP	F	402	6	-	6/19/38/38	0/3/3/3
5	GTP	C	501	6	-	7/22/38/38	0/3/3/3
10	GDP	B	506	6	-	0/16/32/32	0/3/3/3
5	GTP	A	501	6	-	4/22/38/38	0/3/3/3

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	502	MES	C8-S	-7.70	1.66	1.77
9	B	505	UPO	C10-N11	5.30	1.49	1.37
12	F	402	ACP	C5-C4	4.65	1.47	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	B	505	UPO	C12-N11	3.44	1.52	1.46
10	B	506	GDP	C5-C4	3.08	1.47	1.38

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	F	402	ACP	C5-C4-N3	-5.90	118.59	126.72
10	B	506	GDP	C5-C4-N3	-5.85	119.08	128.39
5	C	501	GTP	C5-C4-N3	-5.27	120.00	128.39
5	A	501	GTP	C5-C4-N3	-5.14	120.21	128.39
9	B	505	UPO	N25-C10-N11	5.10	123.00	116.56

There are no chirality outliers.



5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	501	GTP	C5'-O5'-PA-O3A
5	A	501	GTP	C5'-O5'-PA-O2A
5	C	501	GTP	C5'-O5'-PA-O3A
5	C	501	GTP	C5'-O5'-PA-O2A
12	F	402	ACP	PG-C3B-PB-O1B


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand ACP F 402



Ligand GTP C 501





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	438/451 (97%)	0.53	18 (4%)	41	39	33, 69, 88, 107	2 (0%)
1	C	440/451 (97%)	0.20	13 (2%)	52	50	28, 59, 76, 94	5 (1%)
2	B	425/431 (98%)	0.63	27 (6%)	25	25	37, 69, 98, 112	1 (0%)
2	D	423/431 (98%)	0.80	39 (9%)	14	14	58, 82, 101, 113	0
3	E	123/189 (65%)	0.87	16 (13%)	7	6	35, 80, 113, 126	2 (1%)
4	F	349/380 (91%)	1.31	82 (23%)	2	2	65, 93, 140, 151	0
All	All	2198/2333 (94%)	0.68	195 (8%)	15	15	28, 73, 113, 151	10 (0%)

The worst 5 of 195 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	F	135	TYR	5.8
4	F	169	LEU	5.5
2	B	438	ALA	5.0
2	B	247	ASN	4.9
4	F	165	GLU	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GTP	A	501	32/32	-	-	52,53,64,67	42
5	GTP	C	501	32/32	-	-	45,48,58,64	42
5	GTP	D	503	32/32	-	-	72,78,91,91	42
6	MG	A	502	1/1	-	-	57,57,57,57	1
6	MG	B	501	1/1	-	-	56,56,56,56	1
6	MG	C	502	1/1	-	-	50,50,50,50	1
6	MG	D	501	1/1	-	-	85,85,85,85	1
6	MG	F	401	1/1	-	-	93,93,93,93	1
7	CA	A	503	1/1	-	-	77,77,77,77	1
7	CA	A	504	1/1	-	-	64,64,64,64	1
7	CA	B	503	1/1	-	-	67,67,67,67	1
7	CA	B	504	1/1	-	-	69,69,69,69	1
7	CA	C	503	1/1	-	-	63,63,63,63	1
7	CA	D	502	1/1	-	-	76,76,76,76	1
8	MES	B	502	12/12	-	-	56,59,69,69	24
9	UPO	B	505	29/29	0.88	0.17	58,75,94,96	0
10	GDP	B	506	28/28	-	-	49,53,64,65	38
11	CL	D	504	1/1	-	-	66,66,66,66	1
12	ACP	F	402	31/31	-	-	89,100,121,125	45




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GTP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GTP C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP D 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UPO B 505:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP B 506:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ACP F 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.