



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 04:01 AM UTC

PDB ID : 4JKP / pdb_00004jkip
Title : Restricting HIV-1 Pathways for Escape using Rationally-Designed Anti-HIV-1 Antibodies
Authors : Diskin, R.; Bjorkman, P.J.
Deposited on : 2013-03-11
Resolution : 2.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4591 (2.84-2.80)
Clashscore	190562	5010 (2.84-2.80)
Ramachandran outliers	187476	4916 (2.84-2.80)
Sidechain outliers	187428	4918 (2.84-2.80)
RSRZ outliers	180081	4594 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	G	361	2% 78% 16% 6%
2	H	229	83% 14% 3%
3	L	210	1% 79% 19% 1%
4	A	6	33% 50% 17%
5	B	5	40% 60%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	C	3	33% 67%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 6263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called gp120.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	G	340	Total	C	N	O	S	0	1	0
			2669	1674	463	508	24			


- Molecule 2 is a protein called Heavy chain of antibody 45-46M2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	222	Total	C	N	O	S	0	1	0
			1728	1093	304	321	10			

- Molecule 3 is a protein called Light chain of antibody 45-46M2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	207	Total	C	N	O	S	0	0	0
			1601	1003	273	321	4			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	A	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	B	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	C	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	G	1	Total	C	N	O	0	0
			14	8	1	5		
7	G	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	G	1	Total	C	N	O	0	0
			14	8	1	5		
7	G	1	Total	C	N	O	0	0
			14	8	1	5		
7	L	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	G	7	Total	O	0	0
			7	7		
8	H	13	Total	O	0	0
			13	13		
8	L	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: gp120




- Molecule 2: Heavy chain of antibody 45-46M2




- Molecule 3: Light chain of antibody 45-46M2





- Molecule 4: alpha-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain A:  33% 50% 17%




- Molecule 5: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain B:  40% 60%



- Molecule 6: β -D-mannopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose-(1-4)-2-acetamido-2-deoxy- β -D-glucopyranose

Chain C:  33% 67%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.33Å 70.49Å 232.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.85 – 2.82 34.85 – 2.82	Depositor EDS
% Data completeness (in resolution range)	98.5 (34.85-2.82) 92.5 (34.85-2.82)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 2.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1026)	Depositor
R, R_{free}	0.193 , 0.231 0.195 , 0.234	Depositor DCC
R_{free} test set	1390 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	65.1	Xtriage
Anisotropy	0.530	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.034 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6263	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, NAG, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	G	0.41	0/2728	0.81	2/3703 (0.1%)
2	H	0.35	0/1772	0.74	0/2411
3	L	0.38	0/1637	0.87	2/2223 (0.1%)
All	All	0.39	0/6137	0.81	4/8337 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	26	SER	N-CA-C	-6.14	105.30	114.64
1	G	211	ASP	CA-C-N	5.58	125.50	119.76
1	G	211	ASP	C-N-CA	5.58	125.50	119.76
3	L	89	TYR	CB-CA-C	-5.30	110.03	117.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	2669	0	2598	28	0
2	H	1728	0	1683	20	0
3	L	1601	0	1538	23	0
4	A	72	0	61	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	61	0	52	1	0
6	C	39	0	34	0	0
7	G	56	0	52	0	0
7	L	14	0	13	0	0
8	G	7	0	0	0	0
8	H	13	0	0	1	0
8	L	3	0	0	0	0
All	All	6263	0	6031	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:207:LYS:NZ	5:B:4:MAN:O4	2.09	0.86
2:H:103:ASP:OD2	8:H:312:HOH:O	2.03	0.76
1:G:97:LYS:NZ	2:H:99(C):ASP:OD2	2.15	0.74
1:G:333:ILE:HD12	1:G:390:LEU:HD11	1.72	0.71
2:H:195:THR:OG1	2:H:196:GLN:N	2.30	0.64
3:L:104:ARG:NH1	3:L:105:THR:O	2.31	0.63
3:L:37:ARG:HH22	3:L:43:ARG:NH1	1.97	0.62
2:H:99(C):ASP:OD1	2:H:99(D):TYR:N	2.36	0.59
2:H:99(B):ARG:HH11	2:H:99(B):ARG:HB2	1.66	0.59
2:H:123:PRO:HB3	2:H:149:TYR:HB3	1.85	0.58
2:H:44:ARG:HH22	3:L:94:GLN:HE21	1.53	0.56
1:G:101:VAL:HG21	1:G:480:ARG:HG2	1.88	0.55
3:L:104:ARG:HD2	3:L:167:SER:HB2	1.89	0.55
2:H:5:SER:HB2	2:H:23:ARG:HB3	1.89	0.54
1:G:231:LYS:HD2	1:G:267:GLU:HG3	1.89	0.54
1:G:104:MET:HE3	1:G:483:LEU:HD11	1.93	0.51
1:G:50:THR:OG1	1:G:51:THR:N	2.43	0.51
1:G:265:LEU:HD21	1:G:291:SER:HB3	1.93	0.51
2:H:99(B):ARG:HB2	2:H:99(B):ARG:NH1	2.26	0.50
3:L:193:THR:HG22	3:L:200:PRO:HG3	1.94	0.50
1:G:93:PHE:HB2	1:G:233:PHE:HZ	1.76	0.49
1:G:425:ASN:HB2	2:H:54:TRP:HZ3	1.77	0.49
2:H:172:ALA:HB2	2:H:182:LEU:HD23	1.95	0.48
1:G:91:GLU:OE1	1:G:487:LYS:NZ	2.45	0.48
3:L:143:GLN:HB3	3:L:191:GLU:HB3	1.96	0.48
3:L:87:GLN:NE2	3:L:89:TYR:O	2.42	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:141:LYS:HB3	3:L:193:THR:OG1	2.14	0.47
1:G:298:ARG:NH2	1:G:441:GLY:O	2.47	0.47
1:G:120:VAL:HG22	1:G:434:MET:HB3	1.96	0.47
3:L:5:THR:HA	3:L:94:GLN:HE22	1.79	0.47
2:H:47:TRP:CZ2	2:H:49:GLY:HA2	2.50	0.46
1:G:208:ILE:HG13	1:G:380:GLY:HA3	1.98	0.46
1:G:277:LEU:HB2	4:A:1:NAG:H82	1.98	0.46
1:G:280:ASN:HD21	3:L:90:GLU:HG2	1.81	0.46
2:H:37:ILE:HD11	2:H:104:PHE:CE2	2.52	0.45
3:L:37:ARG:NH2	3:L:43:ARG:NH1	2.63	0.45
1:G:93:PHE:HB2	1:G:233:PHE:CZ	2.53	0.44
1:G:104:MET:HE2	1:G:217:TYR:CE1	2.53	0.44
3:L:33:TRP:CZ3	3:L:86:CYS:HB3	2.52	0.44
3:L:90:GLU:H	3:L:90:GLU:HG3	1.47	0.44
3:L:5:THR:OG1	3:L:24:ARG:HB3	2.17	0.44
1:G:270:ILE:HD13	1:G:345:VAL:HG12	1.98	0.44
2:H:172:ALA:HA	2:H:182:LEU:HB3	2.01	0.43
2:H:150:PHE:HA	2:H:151:PRO:HA	1.81	0.43
1:G:346:THR:HG22	1:G:361:PHE:HE2	1.83	0.43
2:H:48:MET:HG2	2:H:63:PHE:CE2	2.53	0.43
3:L:136:TYR:CG	3:L:137:PRO:HA	2.54	0.43
1:G:471:GLY:HA2	1:G:472:GLY:HA3	1.73	0.43
3:L:159:VAL:HG12	3:L:160:THR:O	2.19	0.43
1:G:457:ASP:OD2	1:G:469:ARG:NH2	2.52	0.42
1:G:469:ARG:HA	1:G:470:PRO:HD3	1.89	0.42
2:H:121:LYS:NZ	2:H:148:ASP:O	2.47	0.42
3:L:139:GLU:O	3:L:194:HIS:HD2	2.03	0.42
1:G:78:ASP:HA	1:G:79:PRO:HD2	1.89	0.42
1:G:341:VAL:O	1:G:345:VAL:HG13	2.19	0.42
2:H:28:GLU:HB3	2:H:31:ASN:ND2	2.34	0.42
1:G:49:ASP:OD2	1:G:99:ASN:HB2	2.20	0.42
3:L:25:THR:O	3:L:67:ALA:HB1	2.20	0.41
2:H:87:THR:HG23	2:H:114:THR:HA	2.02	0.41
1:G:270:ILE:O	1:G:348:LYS:HE2	2.20	0.41
2:H:18:MET:SD	2:H:113:VAL:HG21	2.60	0.41
3:L:17:GLU:O	3:L:76:LEU:HD13	2.21	0.41
3:L:45:VAL:HG12	3:L:46:ILE:HG13	2.03	0.41
3:L:59:ARG:HH22	3:L:80:ASP:CG	2.29	0.41
1:G:379:ARG:HG3	1:G:443:ILE:HG23	2.03	0.40
3:L:23:CYS:HB2	3:L:33:TRP:CH2	2.56	0.40
3:L:108:ALA:HA	3:L:109:PRO:HD3	1.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	335/361 (93%)	318 (95%)	17 (5%)	0	100	100
2	H	219/229 (96%)	208 (95%)	10 (5%)	1 (0%)	24	53
3	L	205/210 (98%)	198 (97%)	6 (3%)	1 (0%)	24	53
All	All	759/800 (95%)	724 (95%)	33 (4%)	2 (0%)	36	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	104	PHE
3	L	30	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	305/318 (96%)	287 (94%)	18 (6%)	18	46
2	H	189/195 (97%)	186 (98%)	3 (2%)	55	82
3	L	178/181 (98%)	173 (97%)	5 (3%)	38	71
All	All	672/694 (97%)	646 (96%)	26 (4%)	28	62

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	84	ILE
1	G	91	GLU
1	G	208	ILE
1	G	259	LEU
1	G	261	LEU
1	G	297	THR
1	G	342	LEU
1	G	343	LYS
1	G	345	VAL
1	G	349	LEU
1	G	371	ILE
1	G	390	LEU
1	G	415	THR
1	G	416	LEU
1	G	445	CYS
1	G	462	ASN
1	G	464	SER
1	G	492	GLU
2	H	46	GLU
2	H	80	LEU
2	H	195	THR
3	L	3	VAL
3	L	83	VAL
3	L	90	GLU
3	L	133	ASN
3	L	138	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	99	ASN
1	G	301	ASN
2	H	31	ASN
3	L	194	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PCA	H	1	2	7,8,9	1.95	1 (14%)	9,10,12	2.26	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	H	1	2	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	1	PCA	CD-N	5.04	1.47	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	1	PCA	OE-CD-CG	-3.03	121.31	126.72
2	H	1	PCA	CA-N-CD	-2.99	103.33	113.58
2	H	1	PCA	CB-CA-C	-2.99	108.55	112.66
2	H	1	PCA	CB-CA-N	2.68	110.61	103.24
2	H	1	PCA	CG-CD-N	2.61	114.79	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAG	A	1	1,4	14,14,15	0.53	0	17,19,21	0.88	1 (5%)
4	NAG	A	2	4	14,14,15	0.56	0	17,19,21	1.17	2 (11%)
4	BMA	A	3	4	11,11,12	2.21	3 (27%)	15,15,17	1.04	1 (6%)
4	MAN	A	4	4	11,11,12	0.62	0	15,15,17	1.23	1 (6%)
4	MAN	A	5	4	11,11,12	0.61	0	15,15,17	0.56	0
4	MAN	A	6	4	11,11,12	0.59	0	15,15,17	0.57	0
5	NAG	B	1	1,5	14,14,15	0.54	0	17,19,21	1.04	2 (11%)
5	NAG	B	2	5	14,14,15	0.64	0	17,19,21	0.56	0
5	BMA	B	3	5	11,11,12	2.14	3 (27%)	15,15,17	1.20	2 (13%)
5	MAN	B	4	5	11,11,12	0.57	0	15,15,17	0.59	0
5	MAN	B	5	5	11,11,12	0.67	0	15,15,17	0.66	0
6	NAG	C	1	1,6	14,14,15	0.46	0	17,19,21	0.87	1 (5%)
6	NAG	C	2	6	14,14,15	0.58	0	17,19,21	0.71	0
6	BMA	C	3	6	11,11,12	2.12	3 (27%)	15,15,17	1.28	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	1,4	-	1/6/23/26	0/1/1/1
4	NAG	A	2	4	-	3/6/23/26	0/1/1/1
4	BMA	A	3	4	-	2/2/19/22	0/1/1/1
4	MAN	A	4	4	-	1/2/19/22	0/1/1/1
4	MAN	A	5	4	-	0/2/19/22	0/1/1/1
4	MAN	A	6	4	-	0/2/19/22	0/1/1/1
5	NAG	B	1	1,5	-	1/6/23/26	0/1/1/1
5	NAG	B	2	5	-	4/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BMA	B	3	5	-	1/2/19/22	0/1/1/1
5	MAN	B	4	5	-	1/2/19/22	0/1/1/1
5	MAN	B	5	5	-	0/2/19/22	0/1/1/1
6	NAG	C	1	1,6	-	0/6/23/26	0/1/1/1
6	NAG	C	2	6	-	0/6/23/26	0/1/1/1
6	BMA	C	3	6	-	2/2/19/22	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	3	BMA	C4-C3	-5.13	1.39	1.52
6	C	3	BMA	C4-C3	-4.90	1.39	1.52
5	B	3	BMA	C4-C3	-4.83	1.39	1.52
4	A	3	BMA	C2-C3	-3.78	1.46	1.52
5	B	3	BMA	C2-C3	-3.58	1.47	1.52
6	C	3	BMA	C2-C3	-3.56	1.47	1.52
5	B	3	BMA	O5-C1	-2.87	1.38	1.43
6	C	3	BMA	O5-C1	-2.50	1.39	1.43
4	A	3	BMA	O5-C1	-2.45	1.39	1.43

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	4	MAN	C1-C2-C3	3.75	115.10	109.64
6	C	3	BMA	O3-C3-C2	-2.91	104.11	110.05
4	A	2	NAG	C3-C4-C5	2.84	115.38	110.23
5	B	1	NAG	C1-O5-C5	2.76	115.88	112.19
5	B	3	BMA	O4-C4-C3	-2.40	104.72	110.38
4	A	2	NAG	O5-C1-C2	-2.39	107.59	111.29
6	C	1	NAG	C1-O5-C5	2.37	115.36	112.19
4	A	1	NAG	C1-O5-C5	2.30	115.26	112.19
5	B	1	NAG	C6-C5-C4	-2.15	107.73	113.02
4	A	3	BMA	C1-O5-C5	2.09	114.98	112.19
5	B	3	BMA	C1-C2-C3	-2.00	106.73	109.64

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2	NAG	C3-C2-N2-C7
4	A	2	NAG	C8-C7-N2-C2

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	2	NAG	O7-C7-N2-C2
5	B	2	NAG	C8-C7-N2-C2
5	B	2	NAG	O7-C7-N2-C2
6	C	3	BMA	C4-C5-C6-O6
6	C	3	BMA	O5-C5-C6-O6
4	A	3	BMA	C4-C5-C6-O6
4	A	4	MAN	O5-C5-C6-O6
5	B	4	MAN	O5-C5-C6-O6
5	B	3	BMA	O5-C5-C6-O6
4	A	1	NAG	O5-C5-C6-O6
4	A	3	BMA	O5-C5-C6-O6
5	B	1	NAG	O5-C5-C6-O6
5	B	2	NAG	O5-C5-C6-O6
5	B	2	NAG	C4-C5-C6-O6

There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1	NAG	1	0
5	B	4	MAN	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



Oligosaccharide Chain B



Oligosaccharide Chain C



5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	NAG	G	618	1	14,14,15	0.46	0	17,19,21	1.26	1 (5%)
7	NAG	G	615	1	14,14,15	0.54	0	17,19,21	1.17	1 (5%)
7	NAG	G	616	1	14,14,15	0.50	0	17,19,21	0.56	0
7	NAG	G	617	1	14,14,15	0.55	0	17,19,21	0.65	0
7	NAG	L	501	3	14,14,15	0.69	1 (7%)	17,19,21	0.72	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	G	618	1	-	3/6/23/26	0/1/1/1
7	NAG	G	615	1	-	4/6/23/26	0/1/1/1
7	NAG	G	616	1	-	0/6/23/26	0/1/1/1
7	NAG	G	617	1	-	3/6/23/26	0/1/1/1
7	NAG	L	501	3	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	L	501	NAG	C1-C2	2.05	1.55	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	G	618	NAG	C1-O5-C5	3.79	117.26	112.19
7	G	615	NAG	C1-O5-C5	3.24	116.53	112.19
7	L	501	NAG	C1-O5-C5	2.20	115.13	112.19

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	G	615	NAG	O5-C5-C6-O6
7	G	617	NAG	C8-C7-N2-C2
7	G	618	NAG	C8-C7-N2-C2
7	G	618	NAG	O7-C7-N2-C2
7	G	617	NAG	O7-C7-N2-C2
7	G	615	NAG	C4-C5-C6-O6
7	G	617	NAG	O5-C5-C6-O6
7	G	618	NAG	O5-C5-C6-O6
7	G	615	NAG	C1-C2-N2-C7
7	G	615	NAG	C3-C2-N2-C7
7	L	501	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	340/361 (94%)	-0.13	8 (2%) 59 49	39, 67, 125, 174	1 (0%)
2	H	221/229 (96%)	-0.25	0 100 100	42, 69, 98, 142	1 (0%)
3	L	207/210 (98%)	0.11	3 (1%) 73 64	56, 89, 121, 147	0
All	All	768/800 (96%)	-0.10	11 (1%) 73 64	39, 73, 119, 174	2 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	207	LYS	4.6
1	G	396	ILE	3.5
1	G	472	GLY	3.5
1	G	473	GLY	3.5
1	G	409	GLY	3.4
3	L	154	ASN	3.2
3	L	81	PHE	3.0
3	L	3	VAL	2.9
1	G	206	PRO	2.7
1	G	471	GLY	2.6
1	G	48	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	H	1	8/9	0.71	0.11	74,83,88,90	0

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	1	14/15	-	-	79,99,112,122	0
4	NAG	A	2	14/15	-	-	138,146,153,158	0
4	BMA	A	3	11/12	-	-	155,162,165,169	0
4	MAN	A	4	11/12	-	-	147,150,153,154	0
4	MAN	A	5	11/12	-	-	144,150,151,153	0
4	MAN	A	6	11/12	-	-	163,172,173,175	0
5	NAG	B	1	14/15	-	-	46,59,68,71	0
5	NAG	B	2	14/15	-	-	75,79,85,101	0
5	BMA	B	3	11/12	-	-	113,118,127,127	0
5	MAN	B	4	11/12	-	-	116,128,130,130	0
5	MAN	B	5	11/12	-	-	121,126,127,129	0
6	NAG	C	1	14/15	-	-	68,81,87,95	0
6	NAG	C	2	14/15	-	-	104,112,118,125	0
6	BMA	C	3	11/12	-	-	121,129,131,132	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain A:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain B:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain C:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	NAG	G	618	14/15	0.67	0.15	89,107,113,113	0
7	NAG	G	617	14/15	0.74	0.14	91,105,112,116	0
7	NAG	L	501	14/15	0.75	0.15	119,128,133,134	0
7	NAG	G	616	14/15	0.85	0.09	69,79,85,90	0
7	NAG	G	615	14/15	0.86	0.11	74,94,97,97	0

6.5 Other polymers [i](#)

There are no such residues in this entry.