



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 05:11 AM UTC

PDB ID : 5JK9 / pdb_00005jk9
Title : Crystal structure of human IZUMO1
Authors : Ohto, U.; Ishida, H.; Shimizu, T.
Deposited on : 2016-04-26
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	2% 91% 6% ..
1	B	246	2% 86% 6% 6%
1	C	246	3% 87% 10% .
1	D	246	11% 74% 11% 14%
1	E	246	6% 83% 9% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	246	11% 90% 6% . .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Izumo sperm-egg fusion protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	241	Total	C	N	O	S	0	0	0
			1925	1221	323	363	18			
1	B	231	Total	C	N	O	S	0	0	0
			1839	1167	309	345	18			
1	C	237	Total	C	N	O	S	0	1	0
			1896	1200	321	357	18			
1	D	212	Total	C	N	O	S	0	0	0
			1703	1078	291	319	15			
1	E	231	Total	C	N	O	S	0	0	0
			1838	1167	308	345	18			
1	F	238	Total	C	N	O	S	0	0	0
			1895	1200	317	360	18			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	ARG	-	expression tag	UNP Q8IYV9
A	19	SER	-	expression tag	UNP Q8IYV9
A	20	PRO	-	expression tag	UNP Q8IYV9
A	21	TRP	-	expression tag	UNP Q8IYV9
A	256	GLU	-	expression tag	UNP Q8IYV9
A	257	PHE	-	expression tag	UNP Q8IYV9
A	258	LEU	-	expression tag	UNP Q8IYV9
A	259	GLU	-	expression tag	UNP Q8IYV9
A	260	VAL	-	expression tag	UNP Q8IYV9
A	261	LEU	-	expression tag	UNP Q8IYV9
A	262	PHE	-	expression tag	UNP Q8IYV9
A	263	GLN	-	expression tag	UNP Q8IYV9
B	18	ARG	-	expression tag	UNP Q8IYV9
B	19	SER	-	expression tag	UNP Q8IYV9
B	20	PRO	-	expression tag	UNP Q8IYV9
B	21	TRP	-	expression tag	UNP Q8IYV9
B	256	GLU	-	expression tag	UNP Q8IYV9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	257	PHE	-	expression tag	UNP Q8IYV9
B	258	LEU	-	expression tag	UNP Q8IYV9
B	259	GLU	-	expression tag	UNP Q8IYV9
B	260	VAL	-	expression tag	UNP Q8IYV9
B	261	LEU	-	expression tag	UNP Q8IYV9
B	262	PHE	-	expression tag	UNP Q8IYV9
B	263	GLN	-	expression tag	UNP Q8IYV9
C	18	ARG	-	expression tag	UNP Q8IYV9
C	19	SER	-	expression tag	UNP Q8IYV9
C	20	PRO	-	expression tag	UNP Q8IYV9
C	21	TRP	-	expression tag	UNP Q8IYV9
C	256	GLU	-	expression tag	UNP Q8IYV9
C	257	PHE	-	expression tag	UNP Q8IYV9
C	258	LEU	-	expression tag	UNP Q8IYV9
C	259	GLU	-	expression tag	UNP Q8IYV9
C	260	VAL	-	expression tag	UNP Q8IYV9
C	261	LEU	-	expression tag	UNP Q8IYV9
C	262	PHE	-	expression tag	UNP Q8IYV9
C	263	GLN	-	expression tag	UNP Q8IYV9
D	18	ARG	-	expression tag	UNP Q8IYV9
D	19	SER	-	expression tag	UNP Q8IYV9
D	20	PRO	-	expression tag	UNP Q8IYV9
D	21	TRP	-	expression tag	UNP Q8IYV9
D	256	GLU	-	expression tag	UNP Q8IYV9
D	257	PHE	-	expression tag	UNP Q8IYV9
D	258	LEU	-	expression tag	UNP Q8IYV9
D	259	GLU	-	expression tag	UNP Q8IYV9
D	260	VAL	-	expression tag	UNP Q8IYV9
D	261	LEU	-	expression tag	UNP Q8IYV9
D	262	PHE	-	expression tag	UNP Q8IYV9
D	263	GLN	-	expression tag	UNP Q8IYV9
E	18	ARG	-	expression tag	UNP Q8IYV9
E	19	SER	-	expression tag	UNP Q8IYV9
E	20	PRO	-	expression tag	UNP Q8IYV9
E	21	TRP	-	expression tag	UNP Q8IYV9
E	256	GLU	-	expression tag	UNP Q8IYV9
E	257	PHE	-	expression tag	UNP Q8IYV9
E	258	LEU	-	expression tag	UNP Q8IYV9
E	259	GLU	-	expression tag	UNP Q8IYV9
E	260	VAL	-	expression tag	UNP Q8IYV9
E	261	LEU	-	expression tag	UNP Q8IYV9
E	262	PHE	-	expression tag	UNP Q8IYV9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	263	GLN	-	expression tag	UNP Q8IYV9
F	18	ARG	-	expression tag	UNP Q8IYV9
F	19	SER	-	expression tag	UNP Q8IYV9
F	20	PRO	-	expression tag	UNP Q8IYV9
F	21	TRP	-	expression tag	UNP Q8IYV9
F	256	GLU	-	expression tag	UNP Q8IYV9
F	257	PHE	-	expression tag	UNP Q8IYV9
F	258	LEU	-	expression tag	UNP Q8IYV9
F	259	GLU	-	expression tag	UNP Q8IYV9
F	260	VAL	-	expression tag	UNP Q8IYV9
F	261	LEU	-	expression tag	UNP Q8IYV9
F	262	PHE	-	expression tag	UNP Q8IYV9
F	263	GLN	-	expression tag	UNP Q8IYV9

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	E	1	Total	C	N	O	0	0
			14	8	1	5		
2	F	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	166	Total	O	0	0
			166	166		
3	B	132	Total	O	0	0
			132	132		
3	C	78	Total	O	0	0
			78	78		
3	D	65	Total	O	0	0
			65	65		
3	E	39	Total	O	0	0
			39	39		
3	F	57	Total	O	0	0
			57	57		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Izumo sperm-egg fusion protein 1




- Molecule 1: Izumo sperm-egg fusion protein 1



- Molecule 1: Izumo sperm-egg fusion protein 1




- Molecule 1: Izumo sperm-egg fusion protein 1





• Molecule 1: Izumo sperm-egg fusion protein 1

Chain E:  6% 83% 9% • 6%



• Molecule 1: Izumo sperm-egg fusion protein 1

Chain F:  11% 90% 6% • •



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	64.94Å 75.07Å 108.89Å 77.65° 79.11° 70.66°	Depositor
Resolution (Å)	50.00 – 2.10 50.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.4 (50.00-2.10) 97.4 (50.00-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.194 , 0.231 (Not available) , 0.232	Depositor DCC
R_{free} test set	5249 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	38.0	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11717	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/1968	0.96	5/2661 (0.2%)
1	B	0.91	0/1880	0.93	2/2541 (0.1%)
1	C	0.78	0/1941	0.90	1/2625 (0.0%)
1	D	0.78	0/1739	0.93	1/2346 (0.0%)
1	E	0.75	0/1879	0.92	3/2541 (0.1%)
1	F	0.82	0/1937	0.95	5/2620 (0.2%)
All	All	0.81	0/11344	0.93	17/15334 (0.1%)

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	120	GLU	CB-CG-CD	8.00	126.20	112.60
1	F	164	ASP	CA-C-N	-6.70	110.25	122.38
1	F	164	ASP	C-N-CA	-6.70	110.25	122.38
1	E	174	GLN	N-CA-C	-5.91	101.53	110.28
1	B	154	LYS	N-CA-C	-5.84	99.22	108.73

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1925	0	1885	5	0
1	B	1839	0	1802	11	0
1	C	1896	0	1859	15	0
1	D	1703	0	1657	20	0
1	E	1838	0	1799	15	0
1	F	1895	0	1852	5	0
2	A	14	0	13	0	0
2	B	14	0	13	0	0
2	C	14	0	13	1	0
2	D	14	0	13	0	0
2	E	14	0	13	0	0
2	F	14	0	13	0	0
3	A	166	0	0	1	0
3	B	132	0	0	0	0
3	C	78	0	0	1	0
3	D	65	0	0	0	0
3	E	39	0	0	0	0
3	F	57	0	0	0	0
All	All	11717	0	10932	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 69 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:93:ASP:OD1	1:D:96:ARG:NH1	1.85	1.09
1:B:24:ILE:HG21	1:B:61:VAL:HG13	1.58	0.84
1:D:180:LEU:O	1:D:217:THR:HG22	1.80	0.81
1:D:185:ASN:ND2	1:D:185:ASN:H	1.80	0.78
1:D:24:ILE:HG21	1:D:61:VAL:CG1	2.15	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/246 (97%)	237 (99%)	2 (1%)	0	100	100
1	B	227/246 (92%)	224 (99%)	3 (1%)	0	100	100
1	C	236/246 (96%)	229 (97%)	7 (3%)	0	100	100
1	D	206/246 (84%)	201 (98%)	3 (2%)	2 (1%)	12	9
1	E	227/246 (92%)	218 (96%)	7 (3%)	2 (1%)	14	10
1	F	236/246 (96%)	228 (97%)	7 (3%)	1 (0%)	30	28
All	All	1371/1476 (93%)	1337 (98%)	29 (2%)	5 (0%)	30	28

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	240	SER
1	D	166	GLY
1	D	214	LYS
1	F	255	LYS
1	E	166	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/220 (98%)	208 (97%)	7 (3%)	33	37
1	B	205/220 (93%)	195 (95%)	10 (5%)	22	22
1	C	212/220 (96%)	208 (98%)	4 (2%)	50	58
1	D	189/220 (86%)	184 (97%)	5 (3%)	40	46
1	E	205/220 (93%)	200 (98%)	5 (2%)	43	49
1	F	212/220 (96%)	203 (96%)	9 (4%)	26	28
All	All	1238/1320 (94%)	1198 (97%)	40 (3%)	34	38

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	162	SER
1	F	165	CYS
1	E	163	TYR
1	F	30	VAL
1	F	191	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	130	GLN
1	E	118	GLN
1	C	174	GLN
1	D	239	ASN
1	C	116	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	300	1	14,14,15	0.85	1 (7%)	17,19,21	1.93	3 (17%)
2	NAG	B	300	1	14,14,15	0.47	0	17,19,21	1.14	1 (5%)
2	NAG	E	300	1	14,14,15	0.71	0	17,19,21	1.23	2 (11%)
2	NAG	D	300	1	14,14,15	0.65	0	17,19,21	1.96	4 (23%)
2	NAG	A	300	1	14,14,15	0.41	0	17,19,21	1.51	2 (11%)
2	NAG	F	300	1	14,14,15	0.82	0	17,19,21	1.46	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	300	1	-	2/6/23/26	0/1/1/1
2	NAG	B	300	1	-	2/6/23/26	0/1/1/1
2	NAG	E	300	1	-	2/6/23/26	0/1/1/1
2	NAG	D	300	1	-	2/6/23/26	0/1/1/1
2	NAG	A	300	1	-	1/6/23/26	0/1/1/1
2	NAG	F	300	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	300	NAG	C1-C2	2.01	1.55	1.52

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	300	NAG	C1-O5-C5	6.53	120.94	112.19
2	C	300	NAG	C1-O5-C5	5.85	120.03	112.19
2	A	300	NAG	O5-C1-C2	-4.00	105.10	111.29
2	C	300	NAG	C1-C2-N2	3.37	115.74	110.43
2	F	300	NAG	C1-O5-C5	3.23	116.52	112.19

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	300	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	E	300	NAG	O5-C5-C6-O6
2	C	300	NAG	O5-C5-C6-O6
2	D	300	NAG	C4-C5-C6-O6
2	C	300	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	300	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	241/246 (97%)	-0.20	4 (1%) 69 71	20, 33, 55, 80	0
1	B	231/246 (93%)	-0.05	5 (2%) 62 65	22, 35, 58, 97	0
1	C	237/246 (96%)	0.15	7 (2%) 52 55	25, 43, 67, 96	1 (0%)
1	D	212/246 (86%)	0.63	27 (12%) 8 8	25, 48, 101, 140	0
1	E	231/246 (93%)	0.65	14 (6%) 27 29	33, 54, 87, 118	0
1	F	238/246 (96%)	0.65	27 (11%) 10 10	29, 52, 92, 125	0
All	All	1390/1476 (94%)	0.30	84 (6%) 27 29	20, 43, 86, 140	1 (0%)

The worst 5 of 84 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	163	TYR	8.3
1	D	250	VAL	4.9
1	E	88	TRP	4.8
1	E	67	LEU	4.3
1	D	170	VAL	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	F	300	14/15	0.77	0.14	94,108,113,119	0
2	NAG	C	300	14/15	0.80	0.15	80,86,95,96	0
2	NAG	D	300	14/15	0.82	0.14	76,105,111,111	0
2	NAG	E	300	14/15	0.86	0.11	73,79,89,93	0
2	NAG	A	300	14/15	0.90	0.10	43,53,65,69	0
2	NAG	B	300	14/15	0.92	0.09	43,53,64,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.