



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 09:28 AM UTC

PDB ID : 6JKR / pdb_00006jkr
Title : Crystal structure of aspartate transcarbamoylase from Trypanosoma cruzi in complex with carbamoyl phosphate (CP)
Authors : Matoba, K.; Shiba, T.; Nara, T.; Aoki, T.; Nagasaki, S.; Hayamizu, R.; Honma, T.; Tanaka, A.; Inoue, M.; Matsuoka, S.; Balogun, E.O.; Inaoka, D.K.; Kita, K.; Harada, S.
Deposited on : 2019-03-01
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	88% 9% ...
1	B	328	86% 7% 6%
1	C	328	2% 87% 5% 8%
1	D	328	2% 87% 7% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	328	4% 87% 6% 6%
1	F	328	4% 93% 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 16448 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aspartate carbamoyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	14	0
			2591	1632	460	481	18			
1	B	307	Total	C	N	O	S	0	13	0
			2448	1545	428	456	19			
1	C	303	Total	C	N	O	S	0	9	0
			2395	1514	419	444	18			
1	D	309	Total	C	N	O	S	0	14	0
			2472	1562	434	457	19			
1	E	308	Total	C	N	O	S	0	13	0
			2449	1548	426	456	19			
1	F	323	Total	C	N	O	S	0	12	0
			2567	1620	449	481	17			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP O15636
A	0	SER	-	expression tag	UNP O15636
B	-1	GLY	-	expression tag	UNP O15636
B	0	SER	-	expression tag	UNP O15636
C	-1	GLY	-	expression tag	UNP O15636
C	0	SER	-	expression tag	UNP O15636
D	-1	GLY	-	expression tag	UNP O15636
D	0	SER	-	expression tag	UNP O15636
E	-1	GLY	-	expression tag	UNP O15636
E	0	SER	-	expression tag	UNP O15636
F	-1	GLY	-	expression tag	UNP O15636
F	0	SER	-	expression tag	UNP O15636

- Molecule 2 is PHOSPHORIC ACID MONO(FORMAMIDE)ESTER (CCD ID: CP) (formula: $\text{CH}_4\text{NO}_5\text{P}$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			8	1	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			8	1	1	5	1		
2	C	1	Total	C	N	O	P	0	0
			8	1	1	5	1		
2	D	1	Total	C	N	O	P	0	0
			8	1	1	5	1		
2	E	1	Total	C	N	O	P	0	0
			8	1	1	5	1		
2	F	1	Total	C	N	O	P	0	0
			8	1	1	5	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	238	Total	O	0	0
			238	238		
4	B	243	Total	O	0	0
			243	243		
4	C	248	Total	O	0	0
			248	248		


Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	252	Total 252	O 252	0	0
4	E	220	Total 220	O 220	0	0
4	F	223	Total 223	O 223	0	0

- Molecule 1: Aspartate carbamoyltransferase






● Molecule 1: Aspartate carbamoyltransferase



● Molecule 1: Aspartate carbamoyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	88.41Å 158.38Å 89.00Å 90.00° 119.66° 90.00°	Depositor
Resolution (Å)	29.60 – 1.60 29.60 – 1.60	Depositor EDS
% Data completeness (in resolution range)	96.4 (29.60-1.60) 96.3 (29.60-1.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.167 , 0.190 0.164 , 0.188	Depositor DCC
R_{free} test set	13742 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	16.2	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	0.000 for -h-l,k,h 0.000 for l,k,-h-l 0.056 for h,-k,-h-l 0.000 for -h-l,-k,l 0.001 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16448	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	11/2648 (0.4%)	0.92	4/3573 (0.1%)
1	B	0.76	0/2517	0.86	2/3400 (0.1%)
1	C	0.71	0/2456	0.85	0/3317
1	D	0.76	0/2543	0.85	0/3431
1	E	0.70	0/2522	0.85	0/3404
1	F	0.74	0/2640	0.85	0/3565
All	All	0.77	11/15326 (0.1%)	0.87	6/20690 (0.0%)

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	202	MET	C-O	-7.33	1.15	1.24
1	A	203	PRO	C-O	-7.15	1.15	1.23
1	A	13	LYS	C-O	-6.93	1.15	1.24
1	A	11	LYS	C-O	-6.92	1.15	1.23
1	A	5	PRO	C-O	-6.76	1.17	1.24

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3	GLU	N-CA-C	8.38	120.50	111.36
1	A	4	LEU	CA-C-N	-5.76	114.44	120.38
1	A	4	LEU	C-N-CA	-5.76	114.44	120.38
1	A	13	LYS	N-CA-C	5.19	117.95	109.59
1	B	5	PRO	CA-C-N	-5.14	114.65	119.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2591	0	2659	17	0
1	B	2448	0	2517	21	0
1	C	2395	0	2467	12	0
1	D	2472	0	2558	20	0
1	E	2449	0	2535	23	0
1	F	2567	0	2627	15	0
2	A	8	0	2	0	0
2	B	8	0	2	0	0
2	C	8	0	2	0	0
2	D	8	0	2	0	0
2	E	8	0	2	0	0
2	F	8	0	2	0	0
3	A	12	0	16	1	0
3	B	6	0	8	1	0
3	C	12	0	16	2	0
3	D	12	0	16	3	0
3	E	6	0	8	2	0
3	F	6	0	8	0	0
4	A	238	0	0	2	0
4	B	243	0	0	4	0
4	C	248	0	0	1	0
4	D	252	0	0	8	0
4	E	220	0	0	3	0
4	F	223	0	0	1	0
All	All	16448	0	15447	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 100 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:57:LEU:HD21	1:B:103:LEU:HD12	1.28	1.15
1:C:14:SER:OG	3:C:403:GOL:H2	1.65	0.96
1:D:281[B]:LYS:HE3	4:D:704:HOH:O	1.65	0.96

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:84:LYS:HE3	1:E:87:ALA:HB2	1.56	0.86
1:B:57:LEU:CD2	1:B:103:LEU:HD12	2.06	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/328 (101%)	327 (98%)	5 (2%)	0	100	100
1	B	316/328 (96%)	310 (98%)	6 (2%)	0	100	100
1	C	308/328 (94%)	302 (98%)	6 (2%)	0	100	100
1	D	319/328 (97%)	313 (98%)	6 (2%)	0	100	100
1	E	317/328 (97%)	313 (99%)	4 (1%)	0	100	100
1	F	331/328 (101%)	326 (98%)	5 (2%)	0	100	100
All	All	1923/1968 (98%)	1891 (98%)	32 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	284/275 (103%)	278 (98%)	6 (2%)	47	23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	273/275 (99%)	272 (100%)	1 (0%)	84	75
1	C	264/275 (96%)	263 (100%)	1 (0%)	84	75
1	D	274/275 (100%)	273 (100%)	1 (0%)	84	75
1	E	273/275 (99%)	272 (100%)	1 (0%)	84	75
1	F	282/275 (102%)	281 (100%)	1 (0%)	84	75
All	All	1650/1650 (100%)	1639 (99%)	11 (1%)	78	63

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	3	GLU
1	D	276	ARG
1	F	201	GLN
1	E	242	ARG
1	A	255[A]	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	129	HIS
1	F	225	HIS
1	D	291	ASN
1	F	82	ASN
1	D	212	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	D	402	-	5,5,5	0.42	0	5,5,5	0.24	0
3	GOL	F	402	-	5,5,5	0.38	0	5,5,5	0.47	0
2	CP	C	401	-	6,7,7	1.39	1 (16%)	7,10,10	1.06	0
2	CP	F	401	-	6,7,7	1.12	0	7,10,10	1.28	1 (14%)
3	GOL	B	402	-	5,5,5	0.43	0	5,5,5	0.34	0
2	CP	E	401	-	6,7,7	0.98	0	7,10,10	1.38	0
3	GOL	A	402	-	5,5,5	0.40	0	5,5,5	0.44	0
2	CP	B	401	-	6,7,7	1.22	0	7,10,10	1.15	0
3	GOL	C	403	-	5,5,5	0.45	0	5,5,5	0.57	0
3	GOL	C	402	-	5,5,5	0.47	0	5,5,5	0.31	0
2	CP	A	401	-	6,7,7	1.18	1 (16%)	7,10,10	1.66	1 (14%)
3	GOL	D	403	-	5,5,5	0.28	0	5,5,5	0.68	0
3	GOL	A	403	-	5,5,5	0.28	0	5,5,5	0.80	0
3	GOL	E	402	-	5,5,5	0.38	0	5,5,5	0.52	0
2	CP	D	401	-	6,7,7	0.61	0	7,10,10	1.33	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	402	-	-	0/4/4/4	-
3	GOL	F	402	-	-	0/4/4/4	-
2	CP	C	401	-	-	0/3/5/5	-
2	CP	F	401	-	-	0/3/5/5	-
3	GOL	B	402	-	-	0/4/4/4	-
2	CP	E	401	-	-	0/3/5/5	-
3	GOL	A	402	-	-	0/4/4/4	-
2	CP	B	401	-	-	0/3/5/5	-
3	GOL	C	403	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	402	-	-	0/4/4/4	-
2	CP	A	401	-	-	0/3/5/5	-
3	GOL	D	403	-	-	4/4/4/4	-
3	GOL	A	403	-	-	3/4/4/4	-
3	GOL	E	402	-	-	2/4/4/4	-
2	CP	D	401	-	-	0/3/5/5	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	CP	P-O4P	2.87	1.64	1.59
2	A	401	CP	P-O4P	2.27	1.63	1.59

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	CP	O4P-P-O1P	-3.67	97.72	109.47
2	D	401	CP	O4P-P-O1P	-2.76	100.63	109.47
2	F	401	CP	O2P-P-O4P	-2.28	98.69	105.32

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	GOL	O1-C1-C2-O2
3	A	403	GOL	O1-C1-C2-C3
3	D	403	GOL	C1-C2-C3-O3
3	D	403	GOL	O2-C2-C3-O3
3	A	403	GOL	C1-C2-C3-O3

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	GOL	1	0
3	C	403	GOL	1	0
3	C	402	GOL	1	0
3	D	403	GOL	3	0
3	A	403	GOL	1	0
3	E	402	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/328 (98%)	-0.25	4 (1%) 76 80	6, 14, 26, 42	14 (4%)
1	B	307/328 (93%)	-0.24	4 (1%) 75 79	5, 15, 26, 36	13 (4%)
1	C	303/328 (92%)	-0.12	8 (2%) 57 60	8, 16, 29, 41	9 (2%)
1	D	309/328 (94%)	-0.23	8 (2%) 57 60	6, 14, 26, 47	14 (4%)
1	E	308/328 (93%)	-0.07	13 (4%) 40 42	5, 17, 30, 47	13 (4%)
1	F	323/328 (98%)	-0.13	13 (4%) 42 44	7, 16, 31, 40	12 (3%)
All	All	1872/1968 (95%)	-0.17	50 (2%) 56 59	5, 15, 28, 47	75 (4%)

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	264	ALA	6.3
1	F	2	LEU	5.7
1	D	266	ALA	5.6
1	E	90	ILE	4.5
1	D	244	GLN	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	E	402	6/6	0.64	0.22	61,62,62,62	0
3	GOL	C	403	6/6	0.65	0.21	42,43,44,44	0
3	GOL	A	403	6/6	0.78	0.17	34,41,41,43	0
3	GOL	D	403	6/6	0.80	0.16	37,41,43,46	0
3	GOL	B	402	6/6	0.85	0.11	30,32,33,34	0
3	GOL	A	402	6/6	0.90	0.09	25,28,29,29	0
3	GOL	C	402	6/6	0.91	0.10	29,30,31,31	0
3	GOL	F	402	6/6	0.91	0.09	26,28,29,31	0
3	GOL	D	402	6/6	0.92	0.08	25,28,29,29	0
2	CP	D	401	8/8	0.99	0.03	9,10,11,11	0
2	CP	E	401	8/8	0.99	0.03	10,11,12,12	0
2	CP	F	401	8/8	0.99	0.03	9,10,10,11	0
2	CP	A	401	8/8	0.99	0.04	9,10,10,10	0
2	CP	B	401	8/8	0.99	0.03	9,10,11,11	0
2	CP	C	401	8/8	0.99	0.03	11,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.