



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:43 PM UTC

PDB ID : 7JL4 / pdb_00007jl4
Title : Crystal structure of TRIM65 PSpry domain
Authors : Kato, K.; Ahmad, S.; Hur, S.
Deposited on : 2020-07-29
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	191	
1	B	191	
1	C	191	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4846 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tripartite motif-containing protein 65.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	190	Total	C	N	O	S	0	4	0
			1502	960	268	268	6			
1	C	176	Total	C	N	O	S	0	4	0
			1413	892	263	251	7			
1	B	187	Total	C	N	O	S	0	5	0
			1484	949	265	264	6			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	LEU	CYS	conflict	UNP Q6PJ69
A	407	SER	CYS	conflict	UNP Q6PJ69
A	421	SER	CYS	conflict	UNP Q6PJ69
C	356	LEU	CYS	conflict	UNP Q6PJ69
C	407	SER	CYS	conflict	UNP Q6PJ69
C	421	SER	CYS	conflict	UNP Q6PJ69
B	356	LEU	CYS	conflict	UNP Q6PJ69
B	407	SER	CYS	conflict	UNP Q6PJ69
B	421	SER	CYS	conflict	UNP Q6PJ69

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	185	Total	O	0	0
			185	185		
3	C	127	Total	O	0	0
			127	127		
3	B	111	Total	O	0	0
			111	111		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Tripartite motif-containing protein 65



- Molecule 1: Tripartite motif-containing protein 65



- Molecule 1: Tripartite motif-containing protein 65



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	76.27Å 76.27Å 82.72Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.13 – 1.92 38.13 – 1.92	Depositor EDS
% Data completeness (in resolution range)	99.9 (38.13-1.92) 99.9 (38.13-1.92)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.22 (at 1.92Å)	Xtriage
Refinement program	PHENIX 1.18.2-3874	Depositor
R, R_{free}	0.168 , 0.209 0.169 , 0.211	Depositor DCC
R_{free} test set	2213 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å ²)	27.1	Xtriage
Anisotropy	0.570	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 59.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.008 for -h,-k,l 0.035 for h,-h-k,-l 0.018 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4846	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1563	0.61	0/2138
1	B	0.39	0/1549	0.93	11/2118 (0.5%)
1	C	0.42	0/1464	1.14	6/1989 (0.3%)
All	All	0.41	0/4576	0.91	17/6245 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	2
All	All	0	3

There are no bond length outliers.

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	470	GLN	CG-CD-NE2	-28.72	73.32	116.40
1	C	470	GLN	CG-CD-OE1	20.90	162.60	120.80
1	C	470	GLN	OE1-CD-NE2	-16.78	105.82	122.60
1	B	493	GLU	OE1-CD-OE2	-14.42	88.29	122.90
1	B	348	ARG	CD-NE-CZ	13.39	143.15	124.40
1	B	493	GLU	CG-CD-OE1	13.36	149.13	118.40
1	B	348	ARG	CG-CD-NE	11.65	137.63	112.00
1	B	493	GLU	CG-CD-OE2	-10.32	94.67	118.40
1	C	348	ARG	CD-NE-CZ	7.70	135.17	124.40
1	B	493	GLU	CB-CA-C	-7.67	95.88	109.62
1	C	348	ARG	NE-CZ-NH2	-7.41	112.53	119.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	470	GLN	N-CA-C	-7.09	97.94	108.79
1	B	348	ARG	CA-CB-CG	7.03	128.16	114.10
1	B	470	GLN	CA-CB-CG	6.65	127.39	114.10
1	B	348	ARG	NE-CZ-NH2	-6.10	113.71	119.20
1	B	493	GLU	N-CA-CB	5.79	118.56	110.04
1	B	493	GLU	N-CA-C	-5.34	103.08	110.35

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	493	GLU	Sidechain
1	C	348	ARG	Sidechain
1	C	470	GLN	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1502	0	1431	14	0
1	B	1484	0	1394	24	0
1	C	1413	0	1346	17	1
2	A	18	0	24	2	1
2	C	6	0	8	0	0
3	A	185	0	0	5	0
3	B	111	0	0	6	0
3	C	127	0	0	2	0
All	All	4846	0	4203	56	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:351:GLN:NE2	3:B:601:HOH:O	1.88	1.06

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:500:CYS:SG	3:B:709:HOH:O	2.28	0.90
1:A:380:GLN:O	3:A:701:HOH:O	1.92	0.86
1:B:348:ARG:NH2	3:B:602:HOH:O	2.15	0.79
1:A:372:GLN:HE22	1:A:415:ASN:H	1.26	0.79
1:A:433[B]:GLN:NE2	3:A:702:HOH:O	2.17	0.78
1:B:354:LYS:HD3	1:B:356:LEU:HD21	1.69	0.75
1:B:413:THR:HG23	1:B:490:TRP:HE1	1.51	0.75
1:B:372:GLN:HE22	1:B:415:ASN:H	1.36	0.71
1:C:319:VAL:N	1:C:471:THR:H	1.88	0.70
1:A:343:HIS:CE1	1:A:359:SER:HB2	2.31	0.64
1:A:372:GLN:NE2	1:A:415:ASN:H	1.95	0.63
1:B:413:THR:HG23	1:B:490:TRP:NE1	2.15	0.61
1:C:348:ARG:HB2	1:C:348:ARG:HH11	1.66	0.60
1:C:325:LYS:O	1:C:328:GLN:HG3	2.00	0.60
1:C:468:GLU:OE2	3:C:701:HOH:O	2.17	0.60
1:C:372:GLN:HE22	1:C:415[B]:ASN:H	1.49	0.59
1:B:493:GLU:HG2	1:B:494:GLY:N	2.18	0.58
1:C:372:GLN:HE22	1:C:415[A]:ASN:H	1.49	0.58
1:B:372:GLN:HE21	1:B:414:ASP:HA	1.68	0.58
1:C:372:GLN:NE2	1:C:415[A]:ASN:H	2.02	0.56
1:C:372:GLN:NE2	1:C:415[B]:ASN:H	2.02	0.56
1:B:372:GLN:NE2	1:B:415:ASN:H	2.04	0.55
1:A:372:GLN:HE21	1:A:414:ASP:HA	1.73	0.54
1:A:380:GLN:HE21	1:A:484:PRO:HG3	1.73	0.53
1:A:393:HIS:CE1	1:A:429:GLU:HG3	2.45	0.52
1:C:348:ARG:HB2	1:C:348:ARG:NH1	2.26	0.51
1:C:432:LEU:HB2	1:C:443:LEU:HB2	1.91	0.51
1:B:461:CYS:SG	3:B:708:HOH:O	2.58	0.51
1:B:366:GLY:O	1:B:406:ARG:NH1	2.37	0.50
1:A:376:ALA:C	2:A:603:GOL:H11	2.37	0.50
1:A:396:THR:HB	1:A:490:TRP:HB3	1.94	0.50
1:A:315:VAL:HG11	3:B:693:HOH:O	2.14	0.48
1:B:413:THR:HG22	3:B:698:HOH:O	2.13	0.47
1:A:318:THR:HG22	3:A:709:HOH:O	2.14	0.47
1:C:372:GLN:HB3	1:C:489:PHE:O	2.15	0.47
2:A:601:GOL:H12	3:A:704:HOH:O	2.13	0.46
1:B:400[B]:SER:OG	1:B:404:LEU:HB2	2.15	0.46
1:A:409[A]:LEU:HD23	3:A:736:HOH:O	2.15	0.46
1:B:492:LEU:O	1:B:493:GLU:C	2.59	0.46
1:C:348:ARG:NH1	1:C:352:GLN:OE1	2.47	0.46
1:B:492:LEU:HB3	1:B:495:ARG:HD3	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:359:SER:OG	1:B:361:GLY:O	2.33	0.45
1:B:468:GLU:HA	1:B:469:PRO:C	2.41	0.45
1:B:492:LEU:HG	1:B:493:GLU:H	1.80	0.45
1:A:323:ARG:HD2	1:A:467:LEU:HD12	1.98	0.45
1:C:481:PHE:CD1	1:C:485:LEU:HD11	2.51	0.44
1:C:445:GLY:HA3	3:C:766:HOH:O	2.18	0.44
1:B:371:TRP:CG	1:B:413:THR:HG1	2.36	0.44
1:B:428:GLN:NE2	1:B:433[B]:GLN:OE1	2.51	0.43
1:B:492:LEU:HD23	1:B:495:ARG:NH1	2.33	0.43
1:C:404:LEU:HD22	1:C:488:VAL:HG11	2.01	0.42
1:C:435:TRP:CE2	1:C:440:ALA:HB2	2.54	0.42
1:B:432:LEU:HB2	1:B:443:LEU:HB2	2.02	0.42
1:C:348:ARG:O	1:C:349:GLN:C	2.62	0.42
1:B:323:ARG:HD2	1:B:467:LEU:HD12	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:342:ARG:NH2	2:A:601:GOL:O1[2_665]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/191 (100%)	187 (97%)	5 (3%)	0	100	100
1	B	188/191 (98%)	186 (99%)	2 (1%)	0	100	100
1	C	172/191 (90%)	170 (99%)	2 (1%)	0	100	100
All	All	552/573 (96%)	543 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	161/166 (97%)	161 (100%)	0	100	100
1	B	158/166 (95%)	158 (100%)	0	100	100
1	C	151/166 (91%)	151 (100%)	0	100	100
All	All	470/498 (94%)	470 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	343	HIS
1	A	372	GLN
1	A	380	GLN
1	C	329	ASN
1	C	372	GLN
1	C	380	GLN
1	C	470	GLN
1	C	472	GLN
1	C	478	HIS
1	C	482	ASN
1	B	343	HIS
1	B	349	GLN
1	B	372	GLN
1	B	428	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	603	-	5,5,5	0.84	0	5,5,5	0.87	0
2	GOL	A	602	-	5,5,5	0.93	0	5,5,5	1.06	0
2	GOL	A	601	-	5,5,5	0.63	0	5,5,5	1.38	1 (20%)
2	GOL	C	601	-	5,5,5	0.36	0	5,5,5	1.32	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	603	-	-	4/4/4/4	-
2	GOL	A	602	-	-	0/4/4/4	-
2	GOL	A	601	-	-	4/4/4/4	-
2	GOL	C	601	-	-	0/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	601	GOL	C3-C2-C1	-2.53	102.53	111.80
2	A	601	GOL	C3-C2-C1	-2.46	102.79	111.80

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	GOL	C1-C2-C3-O3
2	A	603	GOL	O1-C1-C2-C3
2	A	603	GOL	C1-C2-C3-O3
2	A	601	GOL	O1-C1-C2-C3
2	A	601	GOL	O1-C1-C2-O2
2	A	601	GOL	O2-C2-C3-O3
2	A	603	GOL	O1-C1-C2-O2
2	A	603	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	603	GOL	1	0
2	A	601	GOL	1	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	190/191 (99%)	-0.30	4 (2%) 63 70	14, 29, 65, 98	4 (2%)
1	B	187/191 (97%)	0.03	4 (2%) 63 70	18, 38, 78, 106	5 (2%)
1	C	176/191 (92%)	0.08	6 (3%) 48 53	18, 38, 72, 102	4 (2%)
All	All	553/573 (96%)	-0.07	14 (2%) 58 64	14, 35, 72, 106	13 (2%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	404	LEU	4.1
1	A	363	GLY	3.6
1	A	313	ALA	3.1
1	B	362	PRO	3.1
1	C	320	CYS	2.8
1	A	409[A]	LEU	2.8
1	C	319	VAL	2.8
1	B	327[A]	TRP	2.6
1	C	405	PRO	2.4
1	B	364	GLY	2.4
1	C	370	LEU	2.3
1	A	318	THR	2.2
1	B	360	ARG	2.2
1	C	371	TRP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	602	6/6	0.65	0.12	70,77,80,80	0
2	GOL	A	603	6/6	0.78	0.14	48,51,54,62	0
2	GOL	A	601	6/6	0.83	0.11	46,49,57,61	0
2	GOL	C	601	6/6	0.92	0.10	33,44,46,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.